

Adiposity and the Brain

The adiposity-brain-axis in mice and men



Ilse Arnoldussen

ISBN: 978-94-6284164-2

© Ilse Arnoldussen, Nijmegen, The Netherlands

Cover design: Ilse Arnoldussen

Lay-out: Theo Hafmans Fotografie & Ilse Arnoldussen

Printed by: Ipskamp, Enschede, The Netherlands

The publication of this thesis was financially supported by the Department of Anatomy, Radboud university medical center, and Donders Institute for Brain Cognition & Behaviour. Financial support by Reckitt Benckiser / Mead Johnson Pediatric Nutrition Institute and Alzheimer Nederland for the publication of this thesis was also gratefully acknowledged.



**alzheimer
nederland**

All right reserved. No parts of this document may be photocopied, reproduced, stored, in a retrieval system, or transmitted, in any form or by any means whether, electronic, mechanical, or otherwise without the prior written permission of the author.

Adiposity and the Brain

The adiposity-brain-axis in mice and men

Proefschrift
ter verkrijging van de graad van doctor
aan de Radboud Universiteit Nijmegen
op gezag van de rector magnificus prof. dr. J.H.J.M. van Krieken,
volgens besluit van het college van decanen
in het openbaar te verdedigen op dinsdag 20 november 2018
om 16.30 uur precies

door

Ilse Adriana Christina Arnoldussen
geboren op 9 november 1988
te Gennep

Promotoren:

Prof. dr. O.C. Boerman

Prof. dr. D.R. Gustafson (State University New York, Verenigde Staten)

Copromotoren:

Dr. A.J. Kiliaan

Dr. M. Wiesmann

Manuscriptcommissie:

Prof. dr. L.A.B. Joosten

Dr. E. Aarts

Prof. dr. S.E. La Fleur (UvA)

‘Omdat je altijd moet doen wat je leuk vindt’

Contents

Chapter 1	9
General introduction	
Thesis overview	
Chapter 2	25
Early intake of long chain polyunsaturated fatty acids preserve brain structure and function in diet induced obesity	
Chapter 3	49
Butyrate restores HFD induced adaptations in brain function and metabolism in mid-adult obese mice	
Chapter 4	67
Adiposity is differentially associated with vascular versus volumetric brain outcomes and by sex in cerebral small vessel disease. The RUN DMC study	
Chapter 5	87
A 10-year follow-up of adiposity and dementia in Swedish adults aged 70-years and older	
Chapter 6	103
General discussion and concluding remarks	
Concluding remarks of this thesis research	
Chapter 7	113
Appendices	
Summary	114
Nederlandse Samenvatting	117
List of Abbreviations	121
References	125
Dankwoord Acknowledgements	157
List of Publications	162
Curriculum Vitae	164
Radboud Alzheimer Centrum Series	165
Donders Graduate School for Cognitive Neuroscience	168

1

GENERAL INTRODUCTION

Parts of the general introduction are based on:

Adipokines: a link between obesity and dementia? Kiliaan A.J., Arnoldussen I.A.C. and Gustafson D.R. (2014).
The Lancet Neurology 13(9): 913-923.

Obesity and dementia: adipokines interact with the brain. Arnoldussen I.A.C., Kiliaan A.J., and
Gustafson D.R. (2014). European Neuropsychopharmacol 24 (12): 1982-99.

Impact of DHA on metabolic diseases from womb to tomb. Arnoldussen I.A.C. and Kiliaan A.J. (2014).
Marine Drugs 12 (12):6190-212.

General introduction

Worldwide, obesity has reached epidemic proportions, and each year obesity or obesity-related-diseases lead to the death of 2.8 million adults (1). Obesity has an undefined aetiology, but is generally caused by imbalance of energy intake versus energy output. This imbalance results in excessive, mainly white adipose tissue (WAT) accumulation, adiposity. Adiposity may eventually have an adverse effect on human health, which can evolve in adiposity related diseases such as cardiovascular disease, type 2 diabetes mellitus (T2DM) and even cognitive impairment and dementia (1-3). Individuals are considered obese when their body mass index (BMI) exceeds 30.0 kg/m². BMI grossly reflects total WAT during adulthood (4). Although, particularly in older individuals, BMI may not be an accurate marker as aging is characterized by lean body mass loss and WAT increase without gaining weight, a phenomenon that is not captured by BMI (5). In addition, the amount of visceral adipose tissue is associated with higher waist circumference (WC) or waist hip ratio (WHR), and therefore this has been another rationale for determining central body adiposity (6-8).

Fatty acids

Intake of diets with a high content of saturated fat (HFD) or high fat and high carbohydrates (HFHC) are considered to be important contributing factors of adiposity. Moreover, high intake of saturated fatty acids is associated with development of non-alcoholic fatty liver disease, hyperlipidemia, and a low state of chronic adipose inflammation characterized by a high level of macrophages and pro-inflammatory cytokines (9-11). Hyperlipidemia, reinforced by inflammatory processes, disrupts vascular function resulting in for instance hypertension, atherosclerosis, and cardiovascular disease (12, 13). Thus, high intake of HFD and HFHC can contribute to the development of adiposity and cardiovascular disease.

On the other hand, uptake of adequate levels of saturated and essential unsaturated fatty acids from dietary sources is indispensable as they serve as building blocks for hormones and phospholipids. For instance, the long-chain polyunsaturated fatty acids (LCPUFAs), docosahexaenoic acid (DHA) and arachidonic acid (ARA) are essential components of phospholipids, can have anti-inflammatory effects, and improve vascular function. Phospholipids form the lipid bilayer of neuronal membranes therewith maintaining optimal membrane fluidity and signal transmission (Table 1)(14, 15). These LCPUFAs have potency in the prevention and treatment of adiposity and adiposity-related-diseases as for instance DHA and ARA can modulate WAT storage, reduce inflammatory processes, improve endothelial function, cerebrovascular reactivity and memory consolidation (16-19). In addition, promising results are found in examining the short chain fatty acid, butyrate as treatment for adiposity and adiposity-related-diseases (20). Butyrate

is a product of bacterial fermentation of mainly indigestible plant polysaccharides and resistant starch in the colon. Obese individuals have a decreased concentration of butyrate-producing bacteria and dietary butyrate intake may protect against diet-induced adiposity, improve insulin sensitivity and increases energy expenditure (21-24). Moreover, butyrate has anti-inflammatory properties and is involved in the regulation of liver lipogenesis, adipocyte differentiation, and energy intake (23). Thus, dietary interventions such as LCPUFAs, butyrate or even indigestible fiber rich diets, could contribute to the prevention and decline of adiposity-related-diseases and adiposity.

	Year	N	Age (Years)	DHA	Outcome
Childhood					
Desci (25)	2002	80 ♂ & ♀	12	Plasma phospholipids, ARA & DHA	Values of arachidonic acid and docosahexaenoic acid were significantly lower in diabetic children than in controls.
Burrows (26)	2011	48 ♂ & ♀	Non-obese: 9.0±0.9 / Obese: 8.9±1.2	Erythrocyte fatty acid; the Omega-3 index (O3I) composition	Obese children had altered erythrocyte fatty acid composition unrelated to reported dietary intake. A greater proportion of obese children had an omega-3 index of < 4.0 (high risk) compared with non-obese children.
Vasickova (27)	2011	120 ♂ & ♀ obese	10.0±1.9	300mg/d DHA + 42 mg/d EPA for 3weeks	Daily consumption of 300mg DHA and 42 mg EPA for three weeks leads to an improvement of the anthropometric and lipid parameters in obese children (27).
Juarez Lopez (28)	2013	201 ♂ & ♀ obese and insulin resistant	11.6±0.7	12 weeks LC-PUFA supplementation, 360mg EPA & 240 mg DHA daily	LC-PUFA supplementation for 12 weeks decreased the concentrations of glucose, insulin, triglyceride-levels and BMI.
Damsgaard (29)	2013	73 ♂ & ♀	10.29±0.58	Plasma DHA & EPA concentrations	DHA was positively associated with mean arterial pressure in boys.
Adolescence					
Dangardt (30)	2012	25 ♂ & ♀	15.6±0.9 ♀ 15.7±1.0 ♂	1.2 g/d LC-PUFAs (DHA & EPA) for 3 months	Three months of supplementation of omega-3 LCPUFA improved glucose and insulin homeostasis in obese girls without influencing body weight.
Adulthood					
Rivellese (31)	1997	16 ♂ & ♀ NIDDM patients with hypertriglyceridemia	56.0±3.0	First two months: 0.96g EPA and 1.59 g DHA per day Last four months: 0.64 gr EPA and 1.06 gr DHA per day	DHA and EPA significantly reduced plasma triglycerides and VLDL- triglycerides without significant changes in blood glucose control.
Mori(32)	1999	56 ♂ overweight & hyperlipidemic	49.1±1.2	4 g/day DHA, EPA or olive oil (placebo) for 6 weeks	Purified DHA but not EPA reduced ambulatory BP and HR in mildly hyperlipidemic men.
Mori(33)	2000	59 ♂ overweight & hyperlipidemic	50.6±1.4	4 g/day DHA, EPA or olive oil (placebo) for 6 weeks	DHA enhances vasodilator mechanisms and attenuates constrictor responses in the forearm microcirculation.
Woodman(34)	2003	♂ & ♀ (Hypertensive and diabetic)	40-75	4 g/day DHA, EPA or olive oil (placebo) for 6 weeks	DHA increased low density lipoprotein particle size
Kelley (35)	2007	34 ♂	55.0±2.0	7.5 g DHA-oil for 90 days	DHA supplementation for 45 d significantly decreased concentrations of fasting triacylglycerol, large VLDL, and intermediate-density lipoproteins and the mean diameter of VLDL particles.
Sneddon(36)	2008	69 ♂	32.4±2.3	3 g/day CLA + 3 g/day omega-3 LC-PUFAs	Supplementation with conjugated linoleic acids (CLAs) plus omega-3 LC-PUFAs prevents increased abdominal fat mass and raises fat-free mass and adiponectin levels in obese adults
Micallef (37)	2009	124 ♂ & ♀	43.79±2.22	Plasma levels of DHA & EPA	BMI, waist circumference and hip circumference were inversely correlated with n-3 PUFA, EPA and DHA (P < 0.05 for all) in the obese group. Obese individuals had significantly lower plasma concentrations of total n-3 PUFA, compared with healthy-weight individuals.
Stirban (38)	2010	34 ♂ & ♀ (T2DM)	56.8±8.3	2 g/d EPA & DHA for 6 weeks	6 wk of supplementation with LC-PUFAs reduced the postprandial decrease in macrovascular function relative to placebo. LC-PUFAs supplementation improved postprandial microvascular function.
Itariu (17)	2012	55 ♂ & ♀ (obese)	39±2.0	3,36 g/d EPA & DHA for 8 weeks	n-3 PUFAs, which was well tolerated, decreased the gene expression of most analyzed inflammatory genes in subcutaneous adipose tissue (P < 0.05) and increased production of anti-inflammatory eicosanoids in visceral adipose tissue and subcutaneous adipose tissue (P < 0.05).
Labonte (39)	2013	12 ♂ (obese +T2DM)	54.1±7.2	3 g/d EPA & DHA for 8 weeks	In obese patients with T2DM, EPA&DHA supplementation did not affect the gene expression of pro-inflammatory cytokines in duodenal cells.
Singhal (40)	2013	328 ♂ & ♀	28.± 4.8	1,6 gr/d DHA	DHA supplementation did not improve endothelial function in healthy adolescents. Only triglyceride and very low-density lipoprotein concentrations were significant lower in DHA-supplemented individuals compared with controls.

McDonald (41)	2013	22 ♂ & ♀ Hypertensive and T2DM	58.6±8.8	Daily supplementation of 1.8 g EPA and 1.5 g DHA for 8 weeks	LC-PUFAs diminish platelet superoxide production in T2DM hypertensive patients <i>in vivo</i> .
Virtanen (42)	2013	2122 ♂	53.1±5.1	Serum levels DHA, EPA, DPA	Men with higher serum level of EPA+DHA+DPA had a 33% lower multivariate-adjusted risk for T2DM. (Trend: P=0.01)
Elderly					
Woodman(43)	2003	51 (39-♂ & 12-♀) Hypertensive and diabetic 54: Ischemic heart disease	61.2±1.2	4 g/day DHA, EPA or olive oil (placebo) for 6 weeks	DHA supplementation significantly reduced collagen aggregation and collagen-stimulated thromboxane release.
Lemaitre (44)	2003	125: non-fatal myocardial infarction	79.1±7.5	DHA & EPA plasma phospholipids	Higher combined dietary intake of DHA and EPA, and possibly α -linolenic acid, may lower the risk of fatal ischemic heart disease in older adults.
Tsitouras (45)	2008	12 ♂ & ♀	66.1±4.5	Supplemented with 4 g/day EPA and DHA	Insulin sensitivity increased significantly after 8 weeks on the EPA- and DHA-diet, and serum C-reactive protein was significantly reduced.
Heine-Börling(46)	2010	1570 686-♂ & 884-♀	64.0±5.42 - ♂ 64.0±5.6 - ♀	Food intake questionnaire; Dutch food composition table (DHA & EPA levels)	Subjects with a fish intake > 19 g/d had a significantly lower prevalence of mild/moderate calcification. EPA plus DHA intake showed no significant associations.
Djousse (47)	2011	3088 ♂ & ♀	75.0	Plasma phospholipids DHA and EPA	DHA is not associated with a higher incidence of T2DM, and individuals with higher EPA and DHA plasma concentrations had lower risk on T2DM.
Mozaffarian (48)	2011	3630 ♂ & ♀	75.0±5.0	Plasma phospholipids: EPA, DHA & DPA	Circulating individual and total omega-3 fatty acid concentrations are associated with lower incidence of congestive heart failure (CHF) in older adults.
Virtanen (49)	2012	768 396-♂ & 372-♀	60.7±6.5	Serum levels DHA, EPA, DPA	Serum EPA+DHA+DPA was associated with lower systolic blood and pulse pressure, not with diastolic blood pressure. Similar results when DHA, EPA and DPA were investigated individually.

Table 1. DHA and metabolic diseases through lifetime. DHA: docosahexaenoic acid; EPA: eicosapentaenoic acid; DPA: docosapentaenoic acid; CLA: conjugated linoleic acids; CHF: congestive heart failure; T2DM: type 2 diabetes mellitus; NIDDM: non-insulin-dependent diabetes mellitus; VLDL: very-low-density-lipoproteinen; N: number of participants; Age is represented in years and in Mean±SD. Mean age >4 and ≤12 Year → childhood; Mean age >12 and ≤21 → adolescence; Mean age >21 and ≤60 → adulthood; Mean age >60 → elderly.

Adiposity and brain function

Adiposity has been associated with alterations in brain structure and function, cognitive deficits, and even dementia and Alzheimer's disease (AD) (50-55). For instance, loss of gray and white matter integrity, functional connectivity, and cortical thickness are observed in children, adolescents and adults (52, 54, 56-62). Moreover, abnormalities in functional brain regions found in obese adults such as in the amygdala, hippocampus, and frontal cortex (63, 64). Age is crucial in examining the association between adiposity and the brain (Fig. 1 and 2) as individuals with midlife obesity have a higher risk of developing dementia in late life (65-70), whereas late life obesity seems to reduce dementia risk (71-74). In addition, adiposity may lead to damage manifested as structural changes similar to those seen in normal aging (75, 76). Exact biological pathways linking adiposity to brain processes are multi-factorial and still unrevealed (77-81). In this thesis we will discuss two pathways via which adiposity might affect brain structure, function and cognition with emphasis on the (cerebro)vascular pathway and adipokines.




Figure 1. Possible life course trajectory of BMI in association with dementia MCP-1=monocyte chemotactic protein-1. PAI-1=plasminogen activator inhibitor-1. Reviewed adipokines regulate: energy balance and metabolism; thrombosis and hypertension; inflammation.

(Cerebro) vascular pathway

Subsequent to the dramatic rise of adiposity, the prevalence of insulin resistance and premature cardiovascular disease increased (82). Insulin resistance plays a central role in the pathogenesis of T2DM and develops long before the onset of the disease (83). Individuals with peripheral insulin resistance often show reduced insulin sensitivity in the brain, which can affect cognitive functioning (84, 85). Two-third of adults in the U.S. are diagnosed as overweight or obese of which 26 million have T2DM (86). Moreover, two-third of severely obese children younger than 12-years of age, develop cardiovascular risk factors (87), and obese individuals show a significantly increased risk of cardiovascular disease as well (86). Importantly, particularly clustering of adiposity, T2DM and cardiovascular risk factors has been linked to alterations in vasculature (e.g. arterial stiffness, hypertension and atherosclerosis), hypoperfusion, structural brain changes, cognitive impairment, and dementia (Fig. 1 and 2)(85, 88).

We hypothesize that adiposity can induce (cerebro)vascular changes such as arterial stiffness, inflammation and hypoperfusion, which can subsequently disrupt brain structure and function, and increase the risk for cognitive impairment and dementia. In more detail, the brain relies on perfusion as a constant cerebral blood flow (CBF) ensures constant oxygen and glucose supply and drainage of ‘waste products’. In humans, overweight and obese individuals often have a decreased CBF (18). A decreased CBF, or hypoperfusion may be a result of reduced small vessel density, inflammation, arterial stiffness, atherosclerosis of the large and small vessels around/within the brain, and/or impaired vasodilation. Hypoperfusion can result in reduced gray and white matter volumes, which subsequently may evolve in cognitive impairment (18, 89, 90). Neuroimaging in obese individuals revealed structural brain changes like white and gray matter abnormalities, already at a very young age (89-92). These structural brain abnormalities may evolve in functional alternations as adiposity is associated with poorer executive functioning skills such as inhibitory control and working memory, and subsequently increases the risk to develop cognitive impairment and dementia (3, 93-95).




Figure 2. The trajectory of body mass index over the life course by chronological age and potential roles of adipokines. PAI-1=plasminogen activator inhibitor-1. HGF=hepatocyte growth factor. NGF=nerve growth factor. MCP-1=monocyte chemotactic protein-1. BMI=body mass index. ADSC=adipose-derived stem cells.

Adipokines

WAT is a complex tissue consisting of multiple cell types with multiple cellular phenotypes depending on cell composition and location (96). WAT secretes hundreds of polypeptides, adipokines, and includes cytokines, acute phase reactants, growth factors and other inflammatory mediators, adipose tissue hormones such as leptin, and other chemical messengers (51, 77, 97). The adipokines secreting profile is fat

depot-specific, which is coherent with differences in adipocyte morphology and the local milieu (98-100). Adipokines act in autocrine, paracrine, or endocrine ways (101), and many affect processes in the periphery and the central nervous system. In addition, brown, epicardial and pancreatic adipose tissue appear to have unique adipokine profiles (102-104). Adipokine release may be dysregulated in obesity, possibly because the adipose tissue may be 'diseased'. Since neurodegenerative and vascular processes observed in dementia affect several brain regions and nuclei (105), the action of adipokines might be altered during neurodegenerative and vascular events, and may even feedback to contribute to neurodegeneration.

Over 700 adipokines have been reported in response to specific chronic or acute stimuli or at rest (106). In table 2 and 3 we present adipokines and their primary known function such as metabolism (e.g. leptin, adiponectin, adipisin), vascular function (e.g. C-reactive protein, serum amyloid A, plasminogen activator inhibitor-1), and inflammatory processes (e.g. interleukins, monocyte chemotactic protein-1) (77, 88). Each may possess more than one function, and these functions do overlap (107). Figure 3 represents an overview of possible pathways of adipokines affecting the brain. Adipokines are extensively reviewed in *Kiliaan et al.* and *Arnoldussen et al.*, and therefore we will shortly discuss the most studied in humans (77,88).

Leptin, a satiety hormone regulates food intake and energy expenditure (108, 109). Observations from human population studies in late life suggest that high leptin levels and high BMI are associated with lower dementia risk when measured within 10 years of clinical dementia diagnosis (110, 111). Leptin can affect hippocampal memory formation as it enhances N-methyl-D-aspartate receptor function and subsequently enhances long term potentiation formation at hippocampal CA1 synapses and rapidly remodels dendrites (112-114). Moreover, leptin can affect vascular function by enhancing platelet aggregation and arterial thrombosis, promoting angiogenesis and inducing proliferation, and migration of vascular smooth muscle cells (115-117). Leptin seems to enhance the calcification of vascular cells, making also the arterial wall a target of leptin (117). While leptin levels may trend with BMI in relation to dementia, given the strong correlation between them, leptin may also have an independent role in health of the ageing brain.

Adiponectin modulates inflammatory responses, energy expenditure, food intake, and a number of metabolic processes including glucose regulation and fatty acid catabolism (118). Adiponectin protects plaque rupture by the inhibition of matrix metalloproteinase function (117). Moreover, endothelium-dependent vasoreactivity is impaired in people with hypoadiponectinemia, which might be at least one cause of hypertension in visceral obesity, probably by inhibiting the expression of adhesion molecules (117, 119).

Energy balance and metabolism

Adiponectin

Adipsin (complement Factor D)

Apelin

Chemerin

Dipeptidyl peptidase-4 (DPP-4) adenosine deaminase complexing protein 2 or CD26

Leptin

Lipocalin

Omentin

Resistin

Retinol binding protein-4

Vaspin

Visfatin, also pre-B cell enhancing factor (PBEF)

Inflammation

IL-6

IL-1

IL-10

IL-8

Monocyte chemotactic protein-1 (MCP-1)

TNF-alpha

Thrombosis & hypertension

Serum amyloid A (SAA)

CRP

PAI-1, total, active

Proteins of the Renin Angiotensin System

Growth factors

NGF

HGF

Brown fat

Fibroblast growth factor-21

Interleukin-6

Insulin-like growth factor-1

Table 2. A sampling of adipokines involved in energy balance and metabolism, inflammation, and thrombosis and hypertension.

Adipokine	Obesity	BBB	Role in Periphery	Role in non-Obese brain	Role in Obese brain
<i>Leptin</i>	↑	Y	Long term regulation food intake (120)	Inhibition of food intake (120)	Leptin resistance (121-127)
			Body weight (120)	Memory formation (112-114, 128)	Deficits in negative feedback loop food intake
			Energy expenditure (120)	Neurogenesis (114, 129)	Impairs memory
				Neuroprotective (130, 131)	Impairs neurogenesis
				Brain structure (132-135)	Impairs neuroprotection Abnormal brain structure
<i>Adiponectin</i>	↓	N	Glucose regulation (136-138)	Neuroprotective (139)	Decreases neuroprotection
			Fatty acid metabolism (136-138) Less inhibition of inflammatory signals (136-138)	Energy expenditure (140)	Decreases energy expenditure
<i>Angiotensinogen</i>	↑	N	Blood pressure regulation (141-146)	Protects BBB (147-149)	Alters CBF (due to high blood pressure)(146, 150)
				Regulates arterial Enhances learning & memory(155)	

Plasminogen activator inhibitor-1	↑	N	Fibrinolysis (158)	Regulates apoptosis (159, 160)	Inhibit BBB fibrinolysis
			Cell migration (158)	Survival of neurites (159)	Increase energy expenditure
			Angiogenesis (158)	Microglial migration (161)	Increase energy intake
				Neuroinflammation (161-166)	
				Brain immune system (161-166)	
				Energy intake (167)	
Interleukin-6	↑	Y	Inflammation (168)	Neuroprotection(169)	Inhibits neurogenesis (170-173)
				NeuroInflammation (170-173)	Decreases synaptic plasticity (170-173)
				Synaptic formation (174)	Disrupts learning & memory processes (170-173)
				Differentiation of neurons and astrocytes (175)	
				Leptin signalling (176, 177)	
Tumor necrosis factor -α	↑	Y	Inflammation (168)	Synaptic transmission (178, 179)	Inhibits neurogenesis (180)
				Synaptic plasticity (181)	Disrupts learning & memory processes (182, 183)
				Neuroinflammation	Increases glutamate excitotoxicity(181)
				Leptin signalling (184)	Damages myelin & oligodendrocytes (185)

Table 3. Overview of reviewed adipokines, their level in obesity, ability to cross the BBB, and roles in the periphery, non-obese and obese brain. ↑; Plasma levels of adipokine are upregulated in obesity. ↓; Plasma levels of adipokine are downregulated.

A chronic low inflammation state in the adipose tissue is established in case of excessive WAT accumulation (adiposity), partly mediated via production of pro-inflammatory adipokines such as tumor necrosis factor alpha (TNF- α) and interleukin-6 (IL-6) (11). Higher plasma IL-6 levels are associated with lower hippocampal gray matter volume (186), and excessive production of TNF- α by WAT could decrease adult neurogenesis by halting cell division of hippocampal progenitor cells, impair long term potentiation facilitation with subsequent memory and learning deficits (77). The release of pro-inflammatory cytokines such as TNF- α and IL-6 might have a pivotal role in the relationship between oxidative stress and endothelial dysfunction (117). Inflammation might have an essential role in the development of insulin resistance, the initiation and progression of atherosclerotic lesions and plaque disruption. Moreover, IL-6 and TNF- α are the main inducers of C-reactive protein (CRP) secretion in the liver. CRP is a marker of low-grade inflammation, and it has been suggested that this protein has a role in the pathogenesis of atherosclerotic lesions in humans (187).

Mouse models

Studies in humans can illustrate an association between adiposity, cognitive function, and related biological variables. However, the exact mechanism behind adiposity related to cognitive impairment cannot be clarified, and therefore translational animal studies are needed. Animal models provide the opportunity to test the efficacy of potential innovative supplementary dietary components in adiposity, anatomical, metabolic, and neurochemical detail. The low-density lipoprotein receptor knock-out Leiden (LDLr^{-/-}) and apolipoprotein E3 Leiden (ApoE3*Leiden) mouse models were chosen in our research because of their adiposity mimicking features when exposed to a diet containing high levels of fat and/or sucrose, such as hyperlipidemia. Secondly, both mouse models were chosen for their vulnerability to develop cardiovascular risk factors often found in obese individuals. ApoE3*Leiden transgenic mice have a rare dominant-negative mutation in the human APOE3 gene and therefore a humanized lipoprotein metabolism. ApoE3*Leiden mice are highly responsive to fat-, sugar- and cholesterol-containing diets, resulting in strongly elevated plasma cholesterol and triglyceride levels with a prominent increase in very low density lipoprotein- and low-density lipoprotein (LDL)-sized lipoprotein fractions (188). LDLr^{-/-} mice lack the functional low-density lipoprotein receptor (LDLr). The LDLr mediates the clearance of cholesterol transported through the bloodstream in low-density lipoproteins by endocytosis (189). LDLr^{-/-} Leiden mice develop obese features on a HFD including hypertension, hyperlipidemia, hypercholesterolemia, increased body weight, and atherosclerosis (190, 191).




Figure 3. Effects of adipokines in the periphery and brain in adiposity. Fat cells produce and secrete adipokines like, leptin (red), PAI-1 (blue), angiotensinogen (green), adiponectin (pink), TNF- α (turquoise) and IL-6 (purple). In the periphery, TNF- α and IL-6 stimulate inflammation; these adipokines trigger the liver to produce acute phase proteins. Adiponectin attenuates the inflammatory response by inhibiting the production of TNF- α and IL-6. Furthermore, adiponectin modulates inflammatory responses, energy expenditure (CNS and periphery), food intake (CNS) and a number of metabolic processes, including glucose regulation and fatty acid catabolism. Angiotensinogen increases blood pressure. PAI-1 inhibits fibrinolysis and also stimulates the inflammatory response. Leptin increases energy expenditure and decreases food intake. Only of leptin, TNF- α and IL-6 it is known they are able to cross the BBB and affect, positively or negatively depending on concentration and environment, brain processes such as food intake, synaptic plasticity, learning and memory, and development of dementia. Furthermore, angiotensinogen, PAI-1, IL-6, TNF- α are also produced by neurons, astrocytes and microglia, and several theories exist that leptin and adiponectin might also be produced in the central nervous system. Within the brain these adipokines can be involved in the regulation of food intake, neurogenesis, neuroprotection, neuroinflammation, synaptic plasticity, memory and learning processes and the onset and progression of dementia. Abbreviations: PAI-1: plasminogen activator inhibitor-1; IL-6: interleukin-6; TNF- α : tumor necrosis factor- α ; CNS: central nervous system; BBB: blood brain barrier; +: increases/stimulates; -: decreases/inhibit.

Thesis overview

In this thesis we aim to elucidate the ‘adiposity-brain-axis’ involving dietary fatty acids (ARA, DHA and butyrate), (cerebro)vasculature, adipokines, and the effect of aging on these processes. In detail the aims of this thesis are:

- Investigate the ability and mechanisms by which specific dietary fatty acids can protect against and counteract HFD induced obesity in mouse models.
- Indicate adiposity associations in aging.
- Examine adiposity in relation to changes in brain structure and dementia in humans.

In **chapter 2** we studied the effect of early life dietary LCPUFAs (ARA and DHA) intake in mild-obese in transgenic ApoE3*Leiden mice. A broad and multidisciplinary experimental approach was used to examine energy metabolism, brain structure, cerebral circulation, and cognition. Therefore, we assessed immunohistochemical studies on fat depots, liver, and brain tissue. Moreover, we tested these mice in behavioral experiments like the open field test and Morris water maze. Neuroimaging was used to determine CBF (using arterial spin labeling (ASL)), functional connectivity (using resting state fMRI), and gray and white matter integrity (using diffusion tensor imaging (DTI)). We hypothesized that the intake of ARA and DHA could attenuate the high fat diet induced effects on metabolism, cerebral blood flow, brain structure and cognition.

The aim of the study in **chapter 3** was to obtain detailed insights in the effects of the short chain fatty acid, butyrate on mid- and late life adiposity in LDLr-/- Leiden mouse model. We hypothesized that the effects of a HFD and butyrate would differ in mid- and late life, according to findings in humans. Metabolism, inflammatory processes, and brain function were extensively examined using blood and microbiome analyses, immunohistochemistry at liver, adipose and brain tissue, the cognitive test the Morris water maze to indicate learning and memory processes, and MRI determining CBF, functional connectivity and brain structure. In the RUN DMC study (**chapter 4**) we examined adiposity indicators (body mass index (BMI), waist circumference (WC), and blood leptin and total adiponectin levels) in association with brain volumes and components of cerebral Small Vessel Disease (cSVD) in adults age ≥ 50 years enrolled in the Radboud University Nijmegen Diffusion tensor and Magnetic resonance imaging Cohort (RUN DMC study). The RUN DMC study provides a nine year follow-up starting in 2006 of Dutch men and women diagnosed with cerebral small vessel disease.

In the study described in **chapter 5** we analyzed in Gothenburg studies if anthropometric parameters as BMI and waist-hip ratio, and serum adipokines

levels like adiponectin and leptin could predict dementia risk in older Swedish adults over a 10 year period. In addition, we examined whether these associations are different between older men and women. These Gothenburg studies provided us a 10-year prospective population-based study of 70 years and older inhabitants of Gothenburg, Sweden.

Chapter **6 and 7** are dedicated to the general discussion and a summary of all study findings, and we propose suggestions for future studies to gain a better understanding of adiposity and brain function.

2

EARLY INTAKE OF LONG CHAIN POLYUNSATURATED FATTY ACIDS PRESERVE BRAIN STRUCTURE AND FUNCTION IN DIET INDUCED OBESITY

Arnoldussen I.A.C.

Zerbi V.

Wiesmann M.

Noordman R.H.J.

Bolijn S.

Mutsaers M.P.C.

Dederen P.J.W.C.

Kleemann R.

Kooistra T.

van Tol E.A.F.

Gross G.

Schoemaker M.H.

Heerschap A.

Wielinga P.Y.

Kiliaan A.J.

Abstract

Worldwide, the incidence of obesity is increasing at an alarming rate, and the number of children with obesity is especially worrisome. These developments raise concerns about the physical, psychosocial and cognitive consequences of obesity. It was shown that early dietary intake of arachidonic acid (ARA) and docosahexaenoic acid (DHA) can reduce the detrimental effects of later obesogenic feeding on lipid metabolism and adipogenesis in an animal model of mild obesity.

In the present study, the effects of early dietary ARA and DHA on cognition and brain structure were examined in mildly obesogenic ApoE*3Leiden mouse model. We used cognitive tests and neuroimaging during early and later life. During their early development after weaning (4-13 weeks of age), mice were fed a chow diet or ARA and DHA-diet for eight weeks, and then switched to a high fat and high carbohydrates (HFHC) diet for 12 weeks (14-26 weeks of age).

A HFHC-diet led to increased energy storage in white adipose tissue, increased cholesterol levels, decreased triglycerides levels, increased cerebral blood flow, and decreased functional connectivity between brain regions as well as cerebrovascular and gray matter integrity. ARA and DHA intake reduced the HFHC diet-induced increase in body weight, attenuated plasma triglycerides levels and improved cerebrovasculature, gray matter integrity and functional connectivity in later life.

In conclusion, a HFHC diet causes adverse structural brain and metabolic adaptations, most of which can be averted by dietary ARA and DHA intake early in life supporting metabolic flexibility and cerebral integrity later in life.

Introduction

Each year, obesity or obesity-related conditions lead to the death of 2.8 million people around the world (1). Besides the well-known metabolic complications, obesity is associated with cognitive dysfunction and changes in the central nervous system particularly in the areas of executive function, attention and concentration (77, 88, 94, 192-194). For example, visuospatial organization is affected in overweight children, and in later life these children often suffer from emotional problems (95, 195). It has been shown that obese children also demonstrate an impaired neural activity (196). Remarkably, obese children and adolescents have a decreased gray matter volume of frontal and limbic lobe regions (197). Therefore, novel approaches related to healthy weight development are urgently required.

Animal research has revealed that exposure to high fat diets and high caloric diets increases inflammatory signals within the brain and decreases neurogenesis (198-200). Diets with a high content of fat and carbohydrates (HFHC) are considered to be important contributing factors of overweight and obesity. Promising results in animal and human studies show the potency of long-chain polyunsaturated fatty acids (LCPUFAs) in the prevention and treatment of obesity and overweight (17, 26, 28, 29, 201, 202). The amount and ratio of omega-6 and omega-3 LCPUFAs that are consumed during childhood may determine the adult metabolic profile and risk factors of disease (203). For instance, supplementation with a specific ARA and DHA mixture in early life reduces body weight gain, adiposity and adipose tissue inflammation later in life in mildly obese mice with a lipoprotein profile comparable to humans (202). Moreover, LCPUFAs are essential to support optimal brain and visual system development. Docosahexaenoic acid (DHA) and arachidonic acid (ARA), belonging to the omega-3 and omega-6 LCPUFAs respectively, are the two most abundant LCPUFAs in the brain (reviewed by (204, 205)).

In this study, we therefore studied the effect of early LCPUFA dietary-intake on cognition, brain function and structure in early and later life in ApoE3*Leiden transgenic mice in the context of mild diet-induced obesity. These mice have a humanized lipoprotein metabolism and develop hyperlipidemia and mild obesity on HFHC diets (188, 202, 206). In addition, this mouse model of mild obesity development has not been used yet to evaluate brain development and function earlier. We used a multi-disciplinary experimental approach in order to examine the programming impact of combined dietary ARA and DHA intake in early life on a HFHC diet in later life. Furthermore, we performed *in vivo* and *ex vivo* experiments that cover diet-induced adaptations in energy metabolism, cerebral circulation and cognition. For this purpose, translational cognitive tests and read-outs were applied that are relevant for the human context. The results of this

study are of scientific and societal importance as they further substantiate and elucidate the importance of early programming effects of healthy balanced diets containing LCPUFAs, capable of maintaining metabolic and cognitive flexibility during obesogenic challenges later in life.

Material and Methods

Animals, diets and study design

Male heterozygous ApoE*3Leiden-transgenic mice were housed in individually-ventilated cages (IVC) in conventional animal rooms (relative humidity 50–60%, temperature 21°C, light cycle 7 a.m.–7 p.m. (5 mice per cage)) in the preclinical imaging centre (PRIME) at the central animal laboratory, Radboudumc Nijmegen, the Netherlands. Mice had ad libitum access to acidified tap water and food. These mice originated from the SPF breeding stock at TNO Metabolic Health Research, Leiden, the Netherlands. The mice carry the mutated form of the human apolipoprotein E3 gene, and are considered as a humanized animal model for hyperlipidemia, atherosclerosis and mild obesity. The ApoE*3Leiden mutation is a rare dominant-negative mutation in the human APOE3 gene, and it is characterized by a tandem duplication of codons 120 to 126 and associated with familial dysbetalipoproteinemia. Besides the APOE*3-Leiden gene, this construct consists of the ApoC1 gene and a promoter element that regulates the expression of ApoE and ApoC1 genes (188, 207, 208). ApoE*3Leiden transgenic mice have been generated by introducing a human APOE*3Leiden gene construct into C57Bl/6 mice. ApoE*3Leiden mice are highly responsive to fat-, sugar-, and cholesterol-containing diets resulting in strongly elevated plasma cholesterol and triglyceride levels with a prominent increase in VLDL- and LDL-sized lipoprotein fractions (188). Of note, female mice are prone to develop high plasma cholesterol levels on atherogenic cholesterol-containing Western type diets, while male mice are less susceptible and show plasma cholesterol levels < 8 mM (209). Under the experimental conditions conducted in our current study, the mice do not develop sufficiently high plasma VLDL/LDL cholesterol for development of atherosclerosis.

The total group of 45 ApoE*3Leiden mice was divided into subgroups. During the early life period, from 4 to 13 weeks, 30 mice in the first group received a standardized chow diet (referred to as: '**Chow**'; Sniff R/M diet V1530, Bio Services BV, Uden, The Netherlands). 15 mice were fed a chow diet containing with 0.129% (w/w) ARA and 0.088% (w/w) DHA in their early life (referred to as: '**ARA&DHA**'; DSM Nutritional Products North America, Columbia, MD, USA). In later life, from 14 to 26 weeks, the group on the chow diet was split into a control group (N=15), which continued on the chow diet (referred to as: '**Chow→Chow**'), and

an intervention group (N=15) which switched to a HFHC-diet (referred to as: '**Chow→HFHC**'). The ARA&DHA-group (N=15) also switched to the HFHC-diet later in life (referred to as: '**ARA&DHA→HFHC**'). This HFHC-diet is based on the diets used by Madsen et al. (210). Table 1 provides the composition of the three diets used.

	Chow		ARA&DHA		HFHC	
	% (w/w)	kcal%	% (w/w)	kcal%	% (w/w)	kcal%
Crude protein	19.0	33	19.0	33	16.4	19
Crude fat	3.3	9	3.3	9	28.8	32
Crude fibre	46.1	58	46.1	58	42.6	49
Total		100		100		100
kcal/g		3.8		3.8		4.8
Starch/Sugar	41.2		41.2		36.9	
C14:0	0.01		0.01		0.82	
C16:0	0.47		0.47		6.17	
C16:1	0.01		0.01		0.77	
C18:0	0.08		0.08		5.09	
C18:1	0.62		0.62		8.88	
C18:2	1.80		1.80		5.17	
C18:3	0.23		0.23		0.63	
C20:4	-		0.129		-	
C20-C22	0.01		0.01		0.26	
C22:6	-		0.088		-	

Table 1. Diet composition. Chow is a conventional diet (Sniff R/M diet V1530, Uden, The Netherlands). In the ARA&DHA diet, the conventional diet is enriched with 0,129% w/w C20:4 (n-6) and 0,088% w/w C22:6 (n-3) fatty acids. The high fat and high carbohydrate (HFHC) diet is a purified diet with a high concentration of carbohydrates and fat, like for instance dextrose, casein and beef tallow (Arie Blok Diets, Woerden, the Netherlands) (50% saturated fat, 46% mono-unsaturated fat and 2 % poly-unsaturated fat). ARA = arachidonic acid; DHA = docosahexaenoic acid; HFHC = high fat & high carbohydrate; (w/w) = weight per weight; kcal% = percent of kilo calorie.

Blood samples were taken after 5 hours of fasting (8 a.m.–1 p.m.) at 4, 10, 12, 20 and 26 weeks of age. These blood samples were tested for plasma triglycerides, cholesterol, insulin and leptin levels. Individual body weight and food intake (at cage level) were monitored over time. To investigate possible changes in cognition and brain structure induced by diet in early life, the mice, aged 12–13 weeks, were assessed in several cognitive tests and MRI experiments. The following experiments were performed: open field, rotarod, object recognition test (ORT), resting state functional MRI (rsfMRI), arterial spin labeling (ASL), and diffusion tensor imaging (DTI). In early life, all mice were assessed in cognitive tests, and 20 chow and 10 ARA&DHA fed mice were examined in MRI experiments. Thereafter, the mice switched to a HFHC diet. In later life these mice, aged 26 weeks, were assessed in a second run of the same cognitive tests and MRI experiments, but only the ORT was replaced for the long term memory test, Morris water maze (MWM). Ten mice per diet group were assessed for MRI experiments, and fifteen mice per diet group were examined during the cognitive tests. At 26 weeks of age, all mice were anesthetized and sacrificed by transcardial perfusion with 0.1M phosphate-buffered saline (PBS), and thereafter organs like brain, liver and adipose tissue were harvested. An overview of the study design is provided in figure 1.




Figure 1. Study design. In early life, ApoE3*Leiden mice received either a chow diet or an ARA&DHA-diet. The first run of cognitive tests was assessed in all mice aged 12 weeks, i.e. open field, rotarod and object recognition test. The following week, these mice were assessed in MRI-experiments (Chow N=20 and ARA&DHA N=10). At 14 weeks of age, the mice switched from diet. All 25 week-old mice were assessed for the second run of cognitive testing, and MRI-experiments (N=10 per diet group). All mice aged 26 weeks were anesthetized and sacrificed by transcardial perfusion. HFHC: high fat and high carbohydrate diet; rsfMRI: resting state functional magnetic resonance imaging; ASL: arterial spin labelling and DTI: diffusion tensor imaging.

Plasma analyses

Total plasma cholesterol and triglyceride levels were measured enzymatically using kits number 11489437 and 11488872 (Roche Diagnostics, Almere, The Netherlands) following a standardized protocol of *Wielinga et al.* (202).

Open field

The open field test (OF) was performed to analyze locomotion and exploration. We followed a protocol previously described by *Janssen et al.* (211-213).

Rotarod

To test sensory-motor integration, all mice were placed on a rotating drum to assess in the rotarod test. The rotarod was performed and analyzed following a standardized protocol described by *Janssen et al.* (213).

Object recognition test

ApoE3*Leiden male mice were assessed in the object recognition test (ORT) to examine short term memory. This method was based on literature of *Brooks et al.*, *Bevins and Besheer* (56, 214). In short, subsequent to habituation, two similar objects were placed in the box and a mouse was allowed to explore the objects after a 30, 60 and 120 minutes delay, respectively. All short term memory trials were recorded, and afterwards blind and randomized scored using Ethovision (EthoVision XT 8.5, Noldus Information Technology, Wageningen, The Netherlands). A mouse was considered as an outlier, if it had no interest in the objects during the two memory trials, or if it sat at the objects. Preference for the novel object was expressed in discrimination ratio, which can be defined as: the time spent exploring the novel object minus the familiar object divided by the total object explorer-time.

Morris water maze

Acquisition and probe

The Morris water maze (MWM) was assessed to examine long term memory and spatial learning abilities of the ApoE3*Leiden mice aged 25 weeks. Mice had to seek a submerged platform located in the North-East (NE) quadrant in a circular pool (104 cm diameter) filled with water (21–22°C; made opaque by the addition of milk powder), as previously described by *Janssen et al.* (213).

Search strategies

During the MWM acquisition phase, mice use different search strategies in context of spatial learning. During subsequent trials mice proceed from random towards hippocampus-dependent, and more direct and precise search strategies (215, 216). These swimming pathways were analyzed, and classified into seven search strategies

in order to indicate hippocampus-dependent and -independent search strategies (Ethovision XT 8.5, Noldus Information Technology, Wageningen, Netherlands).

Immunohistochemistry

The left hemisphere of the brain was postfixed in 4% paraformaldehyde for 24 hours for immunohistochemical purposes. Afterwards the hemispheres were stored in 0.1M phosphate buffered saline (PBS) with 0.01% sodium azide (NaN_3). Coronal brain sections were sectioned using a sliding microtome (Microm HM 440, Walldorf, Germany) equipped with an object table for freeze sectioning at -60°C gaining 8 series of 30- μm thick sections. The right hemispheres were divided into 4 sections and immediately snap frozen and stored in a -80°C freezer. Adipose tissue was also divided and stored for either immunohistochemical or biochemical purposes.

Adipose tissue analysis

Adipose tissues were fixed in formalin, embedded in paraffin and sliced in 5 μm sections. The slices were stained with hematoxylin. Briefly, were photographed with a Zeiss Imager A2 microscope (Zeiss, Germany) at 20x magnification with Ludl 3-axis controlled stepping motor system with a colour digital camera. Quantification was performed with ImageJ software (JAVA based public domain, National institutes of health, Maryland, USA) which calculated the size of the adipocytes (μm^2) for three sections of each fat depots, and around the 30-40 adipocytes per section.

Glucose transporter-1 fluorescent immunohistochemistry

To measure the amount of glucose transporters-1 (GLUT-1) in the blood vessel walls in the brain indication for blood vessel integrity, an intensity staining was used. We used a free-floating fluorescent staining of GLUT-1 on hippocampal sections of 30 μm . Sections reserved for the fluorescence GLUT-1 staining were pre-incubated for half an hour in PBS-BT followed by the primary incubation overnight with rabbit anti-GLUT-1 (1:5000; Millipore, Billerica, Massachusetts, USA) at room temperature. After the primary incubation, the sections were rinsed and then incubated with the second antibody, donkey anti-rabbit-Alexa-445 conjugated (1:100 in PBS-BT; Jackson Immuno Research, Suffolk, UK), for three hours in the dark. Subsequently, sections were rinsed, mounted on gelatin coated slides and air-dried in the dark for two hours. After embedding the sections in Fluorsave the sections were stored at 7°C . Pictures were taken at a 5x magnification with an Axio Imager A2 (Zeiss Germany) and the sections were stored for a longer period at -20°C . Sections were photographed using a shutter speed of 500 ms at 470 nm to excite the fluorescence. ImageJ was used for quantification of amount of positive glucose

transporters in blood vessels in the cortex, thalamus and hippocampus (bregma -1.7 and -2.46). In addition, the surface and intensity were measured.

Quantitative real-time polymerase chain reaction

The snap-frozen midbrains were used for the analysis of mRNA levels using quantitative real-time polymerase chain reaction (qRT-PCR) of: glucose transporters-1 (GLUT-1, (217)), preproinsulin (218), insulin receptor (219), leptin receptor A (220) and B (221), vascular endothelial growth factor (VEGF, (222)), occludin (223). Forward and reverse primer sequences are represented in table 2. A standardized protocol was used for these analyses as previously described by Janssen *et al.* (213). Per primer a negative control consisting of water instead of cDNA was added and per primer one plate was used consisting of both the glyceraldehyd 3-phosphate dehydrogenase (GAPDH) and the primer of interest. GAPDH is a reference gene serving as a control for general cell activity. The quality of the samples was evaluated using the melt curve and amplification plot. The StepOne Software version 2.2.2 was used to determine the threshold cycle (CT) values. To analyze possible differences between dietgroups CT values were normalized by GAPDH and transformed to the comparative CT value via the $2^{-\Delta CT}$ method (224).

Primer	Sequence
GAPDH	FW 5'-AGGTCGGTGTGAACGGATTTG-3'
	REV 5'-TGTAGACCATGTAGTTGAGGTCA-3'
GLUT-1	FW 5'-TCAACACGGCCTTCACTG-3'
	REV 5'-CACGATGCTCAGATAGGACATC-3'
Occludin	FW 5'-AGACTACACGACAGGTGGGG-3'
	REV 5'-CTGCAGACCTGCATCAAAAT-3'
VEGF-A	FW 5'-GGAGATCCTTCGAGGAGCACTT-3'
	REV 5'-GGCGATTTAGCAGCAGATATAAGAA-3'
ObRA	FW 5'-ATGGGCTGTGATCGGAACCTG-3'
	REV 5'-GTCTTCCCAATAAGCATGTCTCC-3'
ObRB	FW 5'-GTTCCAAACCCCAAGAATTG-3'
	REV 5'-TGCTCAAATGTTTCAGGCTTT-3'
Preproinsulin	FW 5'-ATAAAGCTGGTGGGCATCCA-3'
	REV 5'-GCACCAACAGGGCCATGT-3'
InsulinR	FW 5'-ATGGGCTTCGGGAGAGGAT-3'
	REV 5'-GGATGTCCATACCAGGGCAC-3'

Table 2. Forward and reverse sequences of the primers. GAPDH: glyceraldehyde-3-phosphate dehydrogenase; GLUT-1: glucose transporters-1; Occludin: tight junction protein; VEGF: vascular endothelial growth factor; ObRA: leptin receptor A; ObRB: leptin receptor B and InsulinR: insulin receptor; FW: forward; REV: reverse.

MRI-experiments

MRI measurements were performed with a 11.7 T Biospec Avance III small animal MR system (Bruker BioSpin, Ettlingen, Germany) equipped with an actively shielded gradient set of 600 mT/m, and operated by a Paravision 5.1 software platform. We used a circular polarized volume resonator for signal transmission and an actively decoupled mouse brain quadrature surface coil for signal reception (Bruker BioSpin). During the experiment, isoflurane (Nicholas Primal (I) limited, London, United Kingdom) was used for anaesthesia (3.5% for induction for 2 minutes and approximately 1.8% for maintenance in a mixture of oxygen and nitrous oxide (1:2)). Respiration of the animal was monitored using a pneumatic cushion respiratory monitoring system (Small Animal Instruments Inc., NY, USA). Mice were placed in a stereotactic device to immobilize the head. Body temperature was measured using a rectal thermometer and maintained at 37 °C using a heated air flow device. First, gradient echo images were acquired using previously described image parameters (225). Resting state fMRI (rsfMRI) acquisition and analysis were performed to assess functional connectivity within specific brain regions following a protocol of *Zerbi et al.* (226). Arterial spin labeling (ASL) was assessed to indicate cerebral blood flow (CBF) levels using an established ASL method with flow-sensitive alternating inversion recovery (FAIR) technique (227-229). ALS method was performed as previously described by (225, 226). Measurements were assessed in two standardized gas concentrations, first: 2 O₂ : 1 N₂O, and the second, vasoconstriction inducing concentration, 3 O₂ : 0 N₂O. CBF values from both conditions were used to examine the ability of the cerebrovasculature to adapt from a normal condition into a vasoconstrictive condition, which is referred to as vasoactivity. To calculate vasoactivity, the normal CBF values were abstracted from the vasoconstriction CBF values, and this was then divided by the sum of the normal and vasoconstrictive CBF values. To calculate regional CBF we used the same protocol as previously described (225). Finally, diffusion tensor imaging (DTI) was performed to determine gray and white matter integrity following a modified protocol of *Zerbi et al.* (230). All imaging parameters are represented in table 3.

Statistical analysis

A random and double-blind selection procedure was maintained throughout the experiment. The results are expressed as means \pm standard error of mean (SEM). Means were analyzed using repeated measures, univariate or multivariate analysis of Analysis of variance (ANOVAs) with Bonferroni correction for multiple testing in a statistical program, SPSS20 (IBM SPSS Statistics 20, IBM Corporation, Armonk, New York, USA). Diet effects were referred to as significant if the P-value was lower than 0.050, and a trend was implied if the P-value was below 0.080 and above 0.050.

Statistical analysis were only performed on the basis of tests for normality (231) and homogeneity of variance (232). All Data are presented in Mean \pm SEM.

	Imaging sequences			
	Anatomical T2*w	ASL	DTI	rsfMRI
Imaging method	GE	FAIR-ASL	4-shot spin-echo PI	Spin-echo EPI
Echo time (ms)	5	11.8	20	16.9
Repetition time (s)	0.63	13.75	7.55	1.8
Image matrix (pixel x pixel)	512x512	128x128	128x128	96x96
Field-of-view (mm)	40x40	30x30	20x20	25x25
Spatial resolution (μ m/pixel)	78x78x340	234x234x 1000	156x156x500	260x260x500
Number of slices	20x3	1	22	9
Total acquisition time (min)	8	12	18	18

Table 3. List of parameters used in MRI experiments

Results

Metabolic parameters

Body weight and food intake

At the start of the experiment, when the mice were four weeks of age, the averaged body weight was 14.9 ± 0.36 g. Up to 13 weeks, body weight measurements revealed that ApoE3*Leiden mice on ARA&DHA-diet had a significantly ($P=0.028$, $F(1,43)=5.16$) lower body weight compared to the mice on chow-diet (Fig. 2A). Food intake was measured per cage ($N=3$ per diet group), and in early life, from 4 to 13 weeks of age, these mice consumed about the same amount of chow or ARA&DHA-diet (Chow: 4.1 ± 0.052 g diet/mouse/day; ARA&DHA: 3.84 ± 0.052 g diet/mouse/day).

After the diet switch at 14 weeks, mice fed the ARA&DHA \rightarrow HFHC diets tended to maintain a lower body weight when compared to Chow \rightarrow HFHC fed mice ($P=0.080$, $F(1,28)=3.14$) (Fig. 2). The switch to a HFHC-diet caused a significantly reduced food intake in these mice in later life, possibly because rodents often maintain an iso-caloric eating pattern (Chow \rightarrow Chow: 3.13 ± 0.33 ; Chow \rightarrow HFHC: 1.54 ± 0.33 ; ARA&DHA \rightarrow HFHC: 1.50 ± 0.22 ; $P=0.007$, $F(1,4)=12.42$).

Plasma analysis

Moreover, as reported previously by *Wielinga et al.*, the ARA&DHA treated mice showed a decreased plasma cholesterol level compared to chow fed mice (Chow: 3.01 ± 0.08 mM; ARA&DHA: 2.71 ± 0.07 mM; $P=0.02$, $F(1,43)=5.92$). Also plasma triglyceride concentrations were significantly decreased in the ARA&DHA fed mice (Chow: 2.32 ± 0.09 mmol/L; ARA&DHA: 2.01 ± 0.01 mmol/L; $P=0.03$, $F(1,43)=5.20$). Plasma insulin levels were below 0.5 ug/mL in early life and comparable in both groups (Insulin: Chow: 0.019 ± 0.00 ug/mL; ARA&DHA: 0.031 ± 0.00 ug/mL), whereas plasma leptin concentrations were below detectable for both diets.

At 26 weeks of life, plasma cholesterol levels remained stable in the Chow→Chow diet group up to 20 weeks, whereas the plasma cholesterol concentrations increased in the Chow→HFHC and ARA&DHA→HFHC diet group (Chow→Chow: 2.92 ± 0.08 mM; Chow→HFHC: 6.23 ± 0.01 mM; ARA&DHA→HFHC: 6.45 ± 0.12 mM; $P=0.00$, $F(1,28)=476.25$). Plasma triglycerides concentrations decreased in all HFHC fed mice, compared to chow fed mice (Chow→Chow: 2.19 ± 0.30 mmol/L; Chow→HFHC: 1.07 ± 0.02 mmol/L; ARA&DHA→HFHC: 1.12 ± 0.02 mmol/L; $P=0.00$, $F(1,28)=184.07$). Moreover, plasma insulin levels remained low for all diet groups (Chow/Chow: 0.63 ± 0.06 ug/mL; Chow→HFHC: 0.46 ± 0.12 ug/mL; ARA&DHA→HFHC: 0.50 ± 0.11 ug/mL). Leptin plasma levels increased in the mice fed a HFHC diet (Chow→Chow: 1.00 ± 0.30 ng/mL; Chow→HFHC: 2.08 ± 0.58 ng/mL; ARA&DHA→HFHC: 3.14 ± 0.46 ng/mL; $P=0.10$, $F(1,28)=2.87$).

Liver, white adipose tissue and brain tissue analysis

At 26 weeks all mice were sacrificed, and organs harvested and weighed. Liver weight was significantly decreased in the Chow→HFHC-diet group compared to the Chow→Chow-group ($P<0.001$, $F(1,28)=32.63$; Fig. 2B), whereas epididymal and inguinal fat weights were significantly increased in the Chow→HFHC-group compared to the Chow→Chow-group ($P=0.001$, $F(1,28)=12.97$; $P=0.010$, $F(1,28)=7.55$; Fig. 2B).

In order to determine whether the storage of lipids was indeed increased in adipocytes due to HFHC feeding, we measured the size of adipocytes in omental, epididymal and inguinal fat depots. In later life, mice fed a HFHC diet had an increased adipocyte size in all three analyzed fat depots, versus chow fed mice (Omental: $P=0.01$, $F(1,28)=6.84$; Epididymal: $P<0.001$, $F(1,28)=14.88$; Inguinal: $P=0.03$, $F(1,28)=5.41$; Fig. 2C). Only in the inguinal fat depot, the adipocyte size of mice fed ARA&DHA was reduced, when compared to HFHC fed mice ($P=0.05$, $F(1,28)=4.27$; Fig. 2C).

In order to reveal changes in insulin and leptin metabolism in the brain, we performed biochemical analysis of the mice brain tissue. No significant diet effects were found in mRNA levels of the leptin receptors A and B nor in mRNA levels of preproinsulin and the insulin receptor (data not shown).




Figure 2. Metabolic parameters. A) Body weight. In early life, the ARA&DHA-group (N=15) showed a reduced body weight when compared to the Chow-group (N=30). In later life, mice on the ARA&DHAHFHC diets had a decreased body weight compared to the Chow→HFHC-group (N=15 per diet group). B) Organ weights at 26 weeks of life. Liver weight was reduced in the Chow→HFHC-dietgroup. Epididymal and inguinal fat weight was increased in the Chow→HFHC-mice (N=15 per diet group). C) Adipocyte sizes in omental, epididymal and inguinal fat depots. HFHC feeding induced an increase in adipocyte size in omental and epididymal fat depots. Early life ARA&DHA dietary intake reduced the adipocyte size within inguinal fat depots (N=15 per diet group). D) Representative image of histologically (H&E) stained fat tissue. * = $P \leq 0.05$; # = 0.08.

Cerebrovascular parameters

CBF analysis

We performed ASL-measurements to indicate CBF levels in the brains of ApoE3*Leiden mice (Fig. 3). In early life, mice fed the ARA&DHA-diet demonstrated an increased vasoactivity within the cortex, compared to the mice fed a chow-diet ($P=0.034$, $F(1,25)=5.04$) (Fig. 3C). These results illustrate an increased ability of the cerebrovasculature to react on vasoconstrictive stimuli.




Figure 3. CBF-levels and vasoactivity in early life. A) CBF-levels. B) CBF-levels under vasoconstrictive conditions. C) Vasoactivity. It represents the ability of the cerebrovasculature to react on vasoconstrictive stimuli. Mice fed ARA&DHA had an increased vasoactivity in the cortex. Chow: N=18; ARA&DHA: N=9; * = $P \leq 0.05$.

In 26 week-old mice the CBF level was significantly increased in the Chow→HFHC-diet group within the cortex ($P=0.044$, $F(1,14)=4.87$) and thalamus ($P=0.017$, $F(1,14)=7.33$) compared to the Chow→Chow-diet group (Fig. 4A). In addition, a trend for increased CBF-levels of mice on Chow→HFHC diets compared to mice on ARA&DHA→HFHC-diets was found in the thalamus ($P=0.074$, $F(1,17)=3.62$; Fig. 4). In vasoconstrictive conditions, the Chow→HFHC-group revealed again a significant increase in CBF levels in the cortex ($P=0.009$, $F(1,14)=7.33$), hippocampus ($P=0.054$, $F(1,14)=4.43$) and thalamus versus Chow→Chow ($P=0.014$, $F(1,14)=7.86$) (Fig. 4B). No significant diet effects were revealed on the vasoactivity at 26 weeks.

Immunohistochemistry GLUT-1

GLUT-1 is a transporter protein for glucose located in the blood vessel wall and can be used to elucidate vascular integrity in the brain. GLUT-1 fluorescent immunohistochemistry demonstrated significant effects on the number of GLUT-1 positive blood vessels per μm^2 (Fig. 4C). The ARA&DHA→HFHC fed mice showed an increased number of blood vessels compared to the Chow→HFHC fed mice in the cortex ($P=0.025$, $F(1,28)=5.64$), hippocampus ($P=0.019$, $F(1,28)=6.18$) and thalamus ($P=0.017$, $F(1,28)=6.45$) (Fig. 4C). Chow→HFHC fed mice also had less GLUT-1 positive blood vessels compared to the Chow→Chow fed mice within the hippocampus ($P=0.078$, $F(1,28)=3.35$) and thalamus ($P=0.072$, $F(1,28)=3.50$) (Fig. 4C). The amount of GLUT-1 protein within the cortex, hippocampus and thalamus was analyzed by intensity measurements, and mRNA levels of GLUT-1 were analyzed using qRT-PCRs. In both measurements no significant diet effects were found, indicating that the GLUT-1 protein and mRNA expression within the blood vessel wall was unchanged by diet while number of blood vessels was changed by diet. In addition, we performed biochemical experiments (qRT-PCRs) to analyze the blood brain barrier and vasculogenesis in the brain. Occludin and vascular endothelial growth factor A (VEGF-A) were examined, and yet no significant diet effects were found (data not shown).




Figure 4. Cerebrovascular parameters in later life.

A) CBF-levels. Chow→HFHC-diet group had an increased CBF-level within the cortex and thalamus. ARA&DHA→HFHC fed mice demonstrated decreased CBF levels in the thalamus. (Chow→Chow: N=7; Chow→HFHC: N=9; ARA&DHA→HFHC: N=10) B) CBF-levels under vasoconstrictive conditions. Chow→HFHC fed mice had increased CBF levels in the cortex, hippocampus and thalamus. (Chow→Chow: N=7; Chow→HFHC: N=9; ARA&DHA→HFHC: N=10) C) Quantification GLUT-1 immunohistochemistry in the cortex, thalamus and hippocampus. GLUT-1 is a transporter protein located in the blood vessel wall and is used as a marker for cerebrovascular integrity. Chow→HFHC-group showed a significant decreased number of GLUT-1 positive blood vessels in the cortex, hippocampus and thalamus. (N=15 per diet group) D) A representative immunofluorescent image of a GLUT-1 stained hippocampal section. *= $P \leq 0.05$; # ≤ 0.08 .

Cognitive parameters

Behavioral experiments

In early life, we examined locomotion, exploration, short term memory and sensory motor integration in the open field, ORT and rotarod tests. In none of these parameters a significant diet effect was revealed between mice fed the chow or ARA&DHA-diet.

In later life, mice were tested in the open field and rotarod, and again no significant diet effects were found in locomotion, exploration and sensory motor integration. A long term memory test, the MWM, was performed in mice aged 25 weeks. During acquisition, all mice learned and found the platform location. Analyses of the search strategies revealed a significant diet effect ($P=0.020$, $F(1,28)=6.30$); the ARA&DHA→HFHC fed mice used an increased level of spatial-hippocampus

dependent search strategies compared to the Chow→HFHC fed mice (Fig. 5A). In the probe trial, ARA&DHA→HFHC fed mice had a decreased swimming distance ($P=0.042$, $F(1,28)=4.55$) and velocity ($P=0.038$, $F(1,28)=4.73$) compared to Chow→HFHC fed mice.




Figure 5. MWM learning and memory, and DTI gray matter integrity in later life.

A) Classification of the search strategies in the MWM. ARA&DHA→HFHC-group had an increased hippocampal dependent spatial learning-ability. Y-axis displays percentage of used hippocampal dependent search strategy (↑ percentages= more use of a hippocampal dependent search strategy, $N=15$ per diet group) B) Mean diffusivity (MD) levels of mice aged 26 weeks. Chow→HFHC-group had increased MD levels in the hippocampus, motor cortex and somatosensory cortex, which indicates a decreased gray matter integrity. (Chow→Chow: $N=6$; Chow→HFHC: $N=8$; ARA&DHA→HFHC: $N=9$) * = $P \leq 0.05$; # = 0.08 AC: Auditory cortex; Hip: Hippocampus; MC: Motorcortex; SSC: Somatosensory cortex; VC: Visual cortex.

DTI and rsfMRI

DTI-experiments were assessed to determine white and gray matter integrity via fractional anisotropy (FA) and mean diffusivity (MD) levels; no significant diet effects were revealed in FA or MD levels in early life at 13 weeks. However, functional connectivity, assessed by rsfMRI, was increased in mice fed a ARA&DHA-diet in early life. These diet effects were mainly found in functional connections between the following brain regions: auditory, visual, motor and somatosensory cortex (Fig. 6A-B).

In the second set of MRI experiments at 26 weeks of life, DTI parameters, FA and MD levels, were again used to analyze white and gray matter integrity. No significant effects were found in FA-levels, while MD-levels of Chow→HFHC fed mice were affected compared to Chow→Chow fed mice. More specifically, the Chow→HFHC-diet group showed an increased MD-level within the hippocampus ($P=0.059$, $F(1,12)=4.35$), motor cortex ($P=0.026$, $F(1,12)=4.35$) and somatosensory cortex ($P=0.049$, $F(1,12)=4.82$) versus only Chow fed mice (Fig. 5B). In the motorcortex, the MD-levels of the Chow→HFHC fed mice were also significantly increased ($P=0.024$, $F(1,15)=5.80$) when compared to ARA&DHA→HFHC (Fig. 5B). Together, these results may indicate neuronal loss and decreased gray matter integrity in the Chow→HFHC fed mice in these areas (Fig. 5B). rsfMRI analysis revealed a decreased functional connectivity in Chow→HFHC relative to Chow→Chow. These significant diet effects were found in the connectivity between the ventral hippocampus, motor and somatosensory cortex (Fig. 6C-D). Compared to the Chow→HFHC fed mice, the ARA&DHA→HFHC fed mice showed an increased functional connectivity between the ventral hippocampus, motor and somatosensory cortex (Fig. 6E-F).




Figure 6. Early and later life connectivity between brain regions using rsfMRI.

A) Total correlation matrixes of the Chow and ARA&DHA diet group at 13 weeks of life. C-E) In later life at 26 weeks, the total ROI matrixes of the Chow→Chow, Chow→HFHC or ARA&DHA→HFHC fed mice. The selected brain regions, dorsal hippocampus (DH), ventral hippocampus (VH), auditory cortex (AC), motor cortex (MC), somatosensory cortex (SSC) and visual cortex (VC), are subdivided in left hemisphere region (first row) and right hemisphere region (second row). In general the DH and VH are strongly bilaterally connected, and AC, MC and SSC are highly interconnected. A higher Z-score (red) indicates a stronger functional connectivity. B-D-F) Statistical analyses of functional connectivity to reveal a significant diet effect are also represented as a matrix. Yellow= $P < 0.050$. A) In early life, ARA&DHA fed mice had an increased functional connectivity between the motor, visual and somatosensory cortex. B) Statistical matrix of Chow vs ARA&DHA in early life. (yellow= $P < 0.050$) C) The Chow→HFHC-group has a decreased functional connectivity between the ventral hippocampus, motor cortex and somatosensory cortex. D) Statistical matrix of Chow→Chow vs Chow→HFHC in later life. (yellow= $P < 0.050$) E) Functional connectivity was increased in ARA&DHA→HFHC fed mice compared to Chow→HFHC fed mice between the ventral hippocampus, auditory, motor and somatosensory cortex. F) Statistical matrix of Chow→HFHC vs ARA&DHA→HFHC in later life. (yellow= $P < 0.050$) Chow: $N=20$; ARA&DHA: $N=10$; Chow→Chow: $N=7$; Chow→HFHC: $N=7$; ARA&DHA→HFHC: $N=7$.

Discussion

In the present study, we are the first to show that a mild obesogenic HFHC diet influences brain structure and function by increasing CBF, decreasing gray matter integrity and functional connectivity. Secondly, brain structure and function can be preserved by an early life ARA&DHA dietary intake. Moreover, early life programming is important in the etiology of mild obesity.

Early life

Fetal and neonatal time periods are critical time windows in development. Nutrition in these critical periods can have long-term phenotypic consequences through permanent structural changes in organs, which is associated with the development of long term obesity (233-237). Furthermore, the amount and ratio of omega-6 and omega-3 LCPUFAs that are consumed during childhood may determine the adult metabolic profile and risk factors of disease (203). Failure to provide LCPUFAs in these critical developmental periods results in alterations in visual function, spatial learning and memory, nerve growth factor levels and dopamine production, as well as decreased glucose utilization in the brain (238-245). These effects could be related to, for instance, the amount of DHA incorporated in the neuronal membranes, because DHA can change the physicochemical properties of membranes, and deficiency of DHA can impair G-protein receptor-mediated signalling in neural membranes (238, 242, 244, 245). Furthermore, *Gomez-Pinilla et al.* showed that dietary DHA intake elevated the amount of leptin receptor b in the hypothalamus and hippocampus, but did not affect ghrelin receptors indicating that DHA specifically influences leptin signalling in the hypothalamic appetite system (246). Of note, the development of the hypothalamic appetite regulatory network occurs predominantly after birth (247). It is well established that ghrelin affects the orexigenic Neuropeptide Y (NPY) and Agouti-related peptide (AgRP) neurons in the arcuate nucleus of the hypothalamus resulting in an increased appetite (248, 249). In contrast, leptin inhibits the NPY and AgRP neurons and activates the anorexigenic neurons of the melanocortin (α -MSH)/cocaine and amphetamine-regulated transcript (CART) pathways in the arcuate nucleus, which results in decreased food intake and increased energy expenditure (248). ARA metabolites can modulate hypothalamic corticotropin-releasing hormone secretion, and induce a long-term activity-dependent enhancement of synaptic transmission in the hippocampus (16, 250). Thus, the hypothalamus works in concert with the hippocampus to regulate energy homeostasis, food intake and cognitive function (251-253). Treatment with ARA and DHA in early life may affect one or several of these key regulatory processes of metabolic homeostasis, and may involve effects in hypothalamus and hippocampus. The observed effects suggests that LCPUFAs like ARA and DHA exhibit promising effects in the prevention and treatment of obesity

and the metabolic profile, possibly via effects on the hypothalamic appetite system, already in early life.

Metabolism

In early life, ApoE3*Leiden mice fed LCPUFAs gained significantly less weight, and plasma cholesterol and triglycerides levels were decreased. These mice still had a lower body weight after 3 months of HFHC diet feeding, despite increased plasma cholesterol and leptin levels. In later life, we observed in all HFHC treated mice a decreased liver weight, while inguinal and epididymal fat weight increased. These results indicate a changed lipid storage in the HFHC-fed mice, with lipids possibly being stored into the epididymal and inguinal fat tissue instead of in the liver (254). Some studies reported an increased liver weight in mice after high fat and /or high carbohydrate feeding (210, 255), and most of them described an increased fat accumulation in the hepatic cells in the context of NAFLD (256-258). Although, *West et al.* found that an obesity inducing diet did not significantly increase the liver weight in nine inbred mice strains (259). In contrast, in our current study we found a decreased liver weight in later life. This effect appeared not to be related to ARA&DHA treatment. The observed reduction in liver weight upon switching from chow diet to a fat-containing obesogenic diet may be related to an effect on *de novo* lipogenesis in combination with a preferred storage of fat in white adipose tissue during obesogenic diet feeding. Under chow conditions energy comes mainly from carbohydrates and only in little amounts from dietary fat. Therefore, *de novo* lipogenesis is active to produce the fatty acids that are needed for organ function (e.g. heart). When animals are switched to obesogenic diets containing more dietary fat, *de novo* lipogenesis is reduced in the liver, and fat storage processes are activated in adipose tissue (206, 260). In parallel, under high fat diet conditions, fat storage in white adipose tissue is activated and, associated with this, liver fat and mass decline as storage pools in adipose tissue are filled. Indeed, we observe an increase in epididymal WAT tissue after the switch from chow to high fat diets. Of note, under more extreme conditions these initially physiological responses may develop into more pathophysiological responses in liver and adipose tissue, ultimately leading to adipose tissue hypertrophy and inflammation and associated hepatic steatosis with increased liver weights as shown recently in longitudinal studies of diet-induced obesity, adipose tissue inflammation and NAFLD (260, 261). ARA&DHA fed mice displayed reduced adipocyte size within the inguinal fat depot. In agreement, *Wielinga et al.* reported that inguinal adipocyte cells were smaller in ApoE3*Leiden mice with early life ARA&DHA intake after eight weeks of HFHC feeding (202). It is hypothesized that a reduced adipocyte size due to early ARA&DHA supplementation could be a result of variations in adipocyte differentiation, as expression profiling revealed significant differences in expression of hundreds of genes between different adipose depots in rodents (262). These

findings confirm that ARA&DHA dietary intake in early life strengthens the ability to cope with a mild obesogenic diet later in life.

Cerebrovasculature

LCPUFA intake improved the vasoactivity in the brains of ApoE3*Leiden mice in early life. In monkeys, a PET study revealed that DHA-supplementation for one week period resulted in a significantly increased regional CBF response to stimulation (263). *Ellis et al.* found in rabbits that ARA induced a dose-dependent blood vessels dilation response of which the maximum was 100%. DHA itself had no effect on vessel diameter, but reduced the dilation produced by ARA (19). It may be hypothesized that intake of combined ARA and DHA could neutralize the cerebrovascular reactivity.

It is well known that obese or overweight patients have a reduced CBF (18, 264-266). In the current study however, HFHC-diet feeding increased CBF levels, but not the vasoactivity in later life. Moreover, HFHC diet was found to decrease the number blood vessels in the cortex, thalamus and hippocampus, which indicates that a HFHC diet is able to affect cerebrovascular integrity. Other studies revealed that exposure to a high fat diet could induce cerebrovascular remodeling (267-269). It should be taken into account that the blood vessel walls in the relatively mild ApoE3*Leiden obese mouse model are not severely damaged yet and may still be able to adapt to the HFHC induced changes. To elucidate these findings, future research should focus on the role of LCPUFAs in endothelial function, atherosclerotic plaque development and blood pressure in mildly obese and more pronounced obese rodent models in critical periods of development. Overall, we demonstrated that a HFHC-diet provokes adaptations in the cerebrovascular integrity and CBF. Furthermore, dietary ARA&DHA intake in early life is able to counteract these adaptations.

Cognition, brain structure and function

We did not find ARA&DHA effects on cognition after eight weeks of dietary intake. In line with this, no significant differences were found by *Lim et al.* in the results of the maze-learning tests between Crj:CD-1 mice fed DHA diets for a period of one or two weeks (270). However, after a 12 week period of DHA-supplementation, mice required less time to reach the maze exit and strayed into blind alleys fewer times (270). These results suggest that it may take time before learning abilities improve after the incorporation of DHA into the neuronal membranes (270). Mice fed an ARA&DHA-diet showed an increased functional connectivity. In line with our findings, a study of *Grayson et al.* demonstrated that cortical modular organization in monkeys supplemented with DHA resembled a healthy human brain (271). Furthermore, individuals with low levels of dietary LCPUFAs had decreased functional connectivity within the early visual pathway and throughout

higher-order associative cortex and showed impairment of distributed cortical networks (271). These findings emphasize that LCPUFAs modulate brain structure, i.e. functional connectivity.

The ARA&DHA fed mice used a more specific, advanced, and hippocampal dependent cognitive search strategy later in life. This finding implies that dietary ARA&DHA intake can improve hippocampal function. Short exposure to a high fat and sucrose diet has been shown to impair hippocampal dependent place recognition, but not the object recognition (272, 273). HFD exposure in rats impaired long-term spatial reference memory in the Morris water maze, without affecting acquisition or short-term memory (274). Accordingly, LCPUFAs and high fat diets seem to specifically modulate spatial memory.

We revealed an increased structural loss of gray matter in later life due to HFHC exposure. Moreover, ARA&DHA dietary intake protected against neuronal loss. Human studies revealed that obese children had significant regional gray matter reduction in middle temporal gyrus, thalami, pre and postcentral gyrus, and cerebellum (92, 275). DHA plays a crucial role in maintaining cortical neuronal integrity (276). In healthy subjects supplementation with DHA, EPA and vitamin E significantly attenuated gray matter volume loss in the hippocampus and temporal brain regions (277). These findings emphasize the ability of a HFHC diet and LCPUFAs to affect gray matter structure and volume. *Kullman et al.* reported that obese individuals showed a decreased functional connectivity strength in the left insular cortex (278). The insular cortex is considered to be the primary gustatory cortex (279), as it represents an important relay of the neural circuitry, essential in food intake, connecting the hypothalamus, orbitofrontal cortex and limbic system. We showed that functional connectivity was reduced in mice fed HFHC-diet, especially within the hippocampus, motor, and somatosensory cortex. These findings indicate that a HFHC-diet induces impairments within the gray matter neuronal structure and functional connectivity in specific brain regions. Dietary ARA&DHA intake in early life is able to counteract these HFHC diet-induced detrimental effects on brain structure and function in later life.

The study further emphasizes the importance of LCPUFAs as dietary components to counteract detrimental effects related to overweight and obesity, and that early life LCPUFAs support cognitive flexibility during later life challenges. Sensitive imaging techniques are essential to investigate and demonstrate the subtle detrimental adaptations induced by a HFHC diet in a mildly obese rodent model. Future research should focus on interactions of dietary components, optimal supplementation period and the vulnerability within different periods of life.

3

BUTYRATE RESTORES HFD INDUCED ADAPTATIONS IN BRAIN FUNCTION AND METABOLISM IN MID-ADULT OBESE MICE

Arnoldussen I.A.C.

Wiesmann M.

Pelgrim C.E.

Wielemaker E.M.

van Duyvenvoorde W.

Amaral Santos P.

Verschuren L.

Keijser B.J.F.

Heerschap A.

Kleemann R.

Wielinga P.Y.

Kiliaan A.J.

Abstract

Midlife obesity affects cognition and increases risk of developing dementia. Recent data suggest that intake of the short chain fatty acid butyrate could improve memory function, and may protect against diet-induced obesity by reducing bodyweight and adiposity.

We examined the impact of a high fat diet (HFD) followed by intervention with 5% (w/w) dietary butyrate on metabolism, microbiota, brain function and structure in the low-density-lipoprotein receptor knockout (LDLR^{-/-}) mouse model in mid- and late life.

In mid-adult mice, 15 weeks of HFD induced adiposity, liver fibrosis and neuroinflammation, increased systolic blood pressure and decreased cerebral blood flow, and functional connectivity assessed with neuroimaging. The subsequent two months butyrate intervention restored these detrimental effects to chow fed control levels. Both HFD and butyrate intervention decreased variance in fecal microbiota composition. In late adult mice, HFD showed similar detrimental effects and decreased cerebral white and gray matter integrity, whereas butyrate intervention attenuated only metabolic parameters.

HFD induces detrimental effects in mid- and late adult mice, which can be attenuated by butyrate intervention. These findings are consistent with reported associations between midlife obesity and cognitive impairment and dementia in humans. We suggest that butyrate may have potential in prevention and treatment of midlife obesity.

Introduction

Obesity-related diseases are becoming one of the major health-care problems of the twenty-first century (1). Obesity is associated with structural brain changes, cognitive impairment and neurodegenerative diseases, besides the well-known metabolic complications (2, 3). Particularly in midlife, it has been reported that obesity can affect cognitive functioning and increase the risk of developing dementia (3). It has been hypothesized that obesity may exaggerate aging processes (280). Multiple rodent studies provided indications for adverse effects of obesity on cognitive functioning in aged mice (281, 282). Therefore, novel approaches related to weight management in aging are urgently required.

The potency of short chain fatty acids such as butyrate has been demonstrated in treatment of obesity and obesity related disease conditions in recent animal studies (20). Butyrate is a natural product of bacterial fermentation of mainly undigested plant polysaccharides and resistant starch in the colon. It has anti-inflammatory properties and is involved in the regulation of liver lipogenesis, adipocyte differentiation and energy intake (23). In addition, butyrate is associated with associative learning and memory functions in severe Alzheimer's disease mouse models via mediating histone acetylation (283). Obese humans have a decreased concentration of butyrate-producing bacteria (24).

In this study, we evaluated the effect of butyrate on microbiota, brain structure and function, and key metabolic organs in context of high fat diet (HFD)-induced obesity. LDLr^{-/-}-Leiden (LDLr^{-/-}) mice were assessed *in vivo* and *ex vivo* experiments in two age groups, mid- (3-6 months old (m.o.)) and late (6-12 m.o.) adult mice. The LDLr^{-/-} Leiden mouse model was chosen for this study because of its high sensitivity to develop obesity and obesity-associated metabolic and vascular complications in multiple organs when fed HFD with 45% fat content (190, 191, 284, 285), and subsequently we think these complications can detrimentally affect brain function. Our results provide evidence that butyrate can attenuate HFD induced impairment during mid- and late adulthood, whereas HFD induced alternations in brain function can only be restored in mid-adulthood. Overall, this study elucidates the effect of diet in the maintenance of metabolic and cognitive flexibility during life.

Materials and Methods

Animals and diets

Male LDLr^{-/-} Leiden mice originated from the SPF breeding stock at TNO (TNO Metabolic Health Research, Leiden, the Netherlands), and were housed in individually ventilated cages in conventional animal rooms in the preclinical imaging centre (PRIME) at the central animal laboratory, Radboudumc Nijmegen, the Netherlands (ad libitum access to acidified tap water and food; relative humidity 50–60%, temperature 21°C, light cycle 7 a.m.–7 p.m. and a maximum of 5 mice per cage). The animal experiments were approved by an independent institutional ethical committee on animal care and experimentation (Zeist, The Netherlands), approval number DEC3682 containing a statistical power analysis to minimize the number of animals. This study is performed and reported according to ARRIVE guidelines (286).

To examine age differences, we assessed a cohort of 30 LDLr^{-/-} mice of three m.o. and a second cohort of six m.o. mice representing mid- and late adulthood, respectively. Both cohorts consisted of three diet groups: Chow (33.0% kcal protein, 58.0% carbohydrates and 9.0% kcal fat, Sniff R/M-H diet V1530, Sniff Spezialdiäten GmbH, Soest, Germany), HFD (20.0% kcal protein, 35.0% carbohydrates and 45.0% kcal fat, D12451, Research Diets Inc, New Brunswick, USA) and HFD enriched with 5% (w/w) butyrate (HFDB). The dietary compositions are presented in supplementary Table 1. The first cohort, representing mid-adulthood, consisted of a group receiving a standardized chow diet from birth until the end of the study, a second group received a HFD at three months of age and a third group switched to a HFD after three months as well, and at seven m.o. the HFD was enriched with butyrate for two months (Fig. 1A). The second cohort representing late adulthood was subdivided in the same manner, only switched to a HFD at six m.o. and received at ten m.o. a butyrate intervention (Fig. 1B). Mice were examined for diet-induced changes in microbiome, metabolism, cognition, brain structure and function (Fig. 1).




Figure 1. Study design. A) Mid-adulthood, high fat diet (HFD) feeding started at 3 months of age and when 7 m.o. the butyrate intervention started (HFD enriched with butyrate (HFDB)). Cognitive and MRI experiments were performed at 9 months of age. B) Late adulthood, exposure of the HFD started at 6 m.o., and at 10 months of age the butyrate intervention started. Late adult mice were assessed in the cognitive and MRI experiments at 12 months of age. Individual body weight and food intake (at cage level) were monitored over time. Blood samples were taken after 5 hours of fasting (8 a.m.–1 p.m.). At the end of the experiment, all mice were anesthetized and sacrificed by transcardial perfusion with 0.1M phosphate-buffered saline (PBS). Thereafter organs were harvested and used for immunohisto- and biochemical experiments. MB= collect faeces for microbiome analyses; BL= blood sample collection. MWM= assessed in Morris water maze; BP=blood pressure measurements; MRI= examined with magnetic resonance imaging for resting state fMRI (rsfMRI), arterial spin labelling (ASL) and diffusion tensor imaging (DTI). HFD=high fat diet; HFDB=high fat diet enriched with butyrate; m.o.=months old; n= group size.

Plasma analyses

Total plasma cholesterol, insulin and triglyceride levels were measured using standardized ELISA kits (202). Glucose levels were assessed using the OneTouch Ultra glucometer (Lifescan Benelux, Beerse, Belgium).

Fecal sample collection and DNA extraction

Fecal samples were collected at the end of the study and frozen immediately at -80°C . Samples were mechanically homogenized and genomic DNA was isolated using the AGOWA mag Mini kit (DNA Isolation Kit, AGOWA, Berlin, Germany) according to the manufacturer's instructions.

Metagenomic sequencing

A fragment of the 16S rRNA gene (~ 270 bp), spanning the V4 hypervariable regions, was PCR amplified using F515/R806 primers within 30 cycles (287). Purified PCR products were paired-end sequenced on the Illumina MiSeq platform (Illumina, Eindhoven, The Netherlands). The sequence data was processed with Mothur v.1.33.2 (288) and grouped using Minimum Entropy Decomposition (MED) algorithm that clusters 16S rRNA gene amplicons in a sensitive manner (289). Sequences were taxonomically classified by the RDP-II Naive Bayesian Classifier using a 60% confidence threshold. Community profiles were compared by Bray–Curtis dissimilarity and Weighted Unifrac clustering of MED abundance (290). Sequences were normalized to 10000 sequences per samples. Population-level comparison of MED abundance between samples collected was performed using the Metastats tool (291).

Morris water maze

The Morris water maze (MWM) was assessed to examine long term memory and spatial learning abilities. We performed and analyzed the MWM according to standardized protocols previously described (292).

Blood pressure measurements

A computerized and warmed tail-cuff plethysmography device was used to measure systolic blood pressure (SBP; IITC Life Scientific Instruments, Woodland Hill, CA). The mice were habituated to the restrainer and acclimatized to the complete procedure. Six SBP-measurements per trained mouse were averaged to obtain the mean SBP.

MRI-experiments

MRI measurements were performed with a 11.7 T Biospec Avance III small animal MR system (Bruker BioSpin, Ettlingen, Germany) which operated with a Paravision 6.0 software platform. Isoflurane (Nicholas Primal (I) Ltd, London, United Kingdom) was used for anaesthesia (3.5% for induction for 2 minutes and approximately 1.8% for maintenance in a mixture of oxygen and medical air (1:2)). We used standardized protocols for these MRI measurements published by *Zerbi et al.* (226), and we assessed functional connectivity (FC) (resting state fMRI), regional cerebral blood flow (arterial spin labeling (ASL)) and gray and white matter

integrity (diffusion tensor imaging (DTI)). Imaging parameters are represented in supplementary table 2. Mice were excluded from further MRI analysis in case of positioning artefacts.

(Immuno)histochemistry

Mice were sacrificed via transcardial perfusion with 0.1 M phosphate-buffered saline (PBS), and organs harvested e.g. brain, liver and fat depots. Organs used in (immuno)histochemical experiments were post-fixed in 4% paraformaldehyde for 24 hours. 5 µm thick paraffin-embedded cross-sections of liver tissue were stained with hematoxylin and eosin. NAFLD was scored blinded using a general scoring system for rodent models according to an established protocol (293). Ionized calcium binding adapter molecule-1 (IBA-1) antibody was used to detect activated microglia (goat anti-IBA-1 (1:2500; Abcam, ab5076)) as indicator for neuroinflammation in 30-µm thick coronal brain sections. We used a previously described standardized protocol for immunohistochemistry and its quantification (213).

Quantitative real-time polymerase chain reaction

The snap-frozen dissected hippocampus of the right hemisphere was used for the analysis of mRNA levels using quantitative real-time polymerase chain reaction (qRT-PCR) in duplo according to a previously described protocol (213). Forward and reverse primer sequences are represented in supplementary table 3. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) is a reference gene serving as a control for general cell activity.

Statistical analysis

A random and double-blind selection procedure was used throughout the experiment. The results are expressed as means ± standard error of mean (SEM). Means were analyzed using repeated measures or multivariate analysis of variance (ANOVAs) with Bonferroni correction for multiple testing in, SPSS20 (IBM SPSS Statistics 20, IBM Corporation, Armonk, New York, USA). Microbiota correlation analyses were performed using Pearson correlations. Supplementary table 4 provides the P and F values of the found significant effects.

Results

Butyrate restored HFD induced metabolic adaptations in mid- and late adult mice

In the mid-adult cohort (9 m.o.), body weight of the HFD mice increased over time and was 30% higher ($P < 0.001$) compared to Chow mice at the end of the experiment. A two months butyrate intervention in mid-adult mice on HFD did not significantly reduce bodyweight of the HFD fed mice (16%; $P = 0.11$). However,

a two months butyrate intervention in late adult mice on HFD resulted in a body weight loss of 23% ($P<0.001$) compared to HFD fed mice. In this late adult cohort (12 m.o.), body weight of the HFD fed mice was 35% higher ($P<0.001$) compared to chow fed mice. (Fig. 2 and Supplementary table 4). Food intake was monitored per cage ($n=2$ or 3) in grams, and subsequent calculated into kilo calorie intake per day per mouse. No significant differences were found between the three dietary interventions in intake (kcal) over time indicating that the energy intake was the same for all diet groups (Fig. 2 and Supplementary table 4).

A HFD increased the cholesterol and triglycerides plasma levels in mid- and late adult mice compared to Chow. The subsequent butyrate intervention lowered the plasma levels of cholesterol and triglycerides in both mid- and late adult mice. No diet effects were found in glucose plasma levels in age cohorts. In mid-adulthood, plasma insulin levels were slightly increased by HFD. Butyrate intervention restored insulin levels compared to solely a HFD which were even below Chow insulin levels, which was more pronounced in late adulthood.

Liver, epididymal, inguinal and omental fat depots was weighed after sacrifice. Liver weight, omental and inguinal fat depot weight were increased in HFD fed mice both at age cohorts compared to chow fed mice. Subsequent butyrate intervention restored these metabolic adaptations (Supplementary table 4).

Butyrate recovered hepatic steatosis induced by HFD feeding in mid- and late adult mice

A HFD increased the NAFLD as hepatic steatosis and hypertrophy were increased in mid- and late adult mice. In late adult mice a HFD increased the percentage of fibrosis. A butyrate intervention restored these observed changes in the liver as it decreased hypertrophy, fibrosis and macrovesicular steatosis (Fig.2). In addition, inflammatory aggregates in the liver were increased in late adult mice fed a HFD (Supplementary Fig. 1).

Butyrate restored HFD induced spatial memory impairment in mid-adult obesogenic mice

Long term memory and spatial learning were assessed in the MWM. In mid-adulthood, HFDB mice had a shorter latency to reach the platform than HFD mice (Supplementary Fig. 2A). In the first 30 seconds of the probe trial, mid-adult HFD mice were in the former platform NE-quadrant for a shorter time period, compared to HFDB mice (Supplementary Fig. 2C), indicating a diminished spatial memory. The swim paths of all mice were classified to assess the degree of used hippocampus dependent search strategies. Mid-adult mice fed HFD showed less hippocampus-dependent search strategies than mid-adult mice fed chow (Supplementary Fig. 2D). In late adulthood, none of these parameters (acquisition, probe NE duration and search strategy) revealed a significant diet effect indicating that diets are no longer effective in late adulthood concerning spatial memory.




Figure 2. Body weight, calorie intake and liver tissue analysis. A) Body weight in mid adult (9 m.o.) mice at a HFD. A HFD enriched with butyrate (HFDB) started at 7 months of age. B) Body weight in late adult mice (12 m.o.). HFDB started at 9 months of age. C) Calorie intake in mid-adulthood. D) Calorie intake in late adulthood. E) Percentage of microvesicular steatosis. F) Percentage of macrovesicular steatosis. G) Percentage of hypertrophy of hepatic cells. H) Percentage of fibrosis within liver tissue. 9 m.o.= mid-adult. 12 m.o.= late adult. Data are presented as mean \pm SEM. * $P < 0.050$.

Butyrate restored HFD induced SBP and CBF in mid-adulthood

LDLr^{-/-} mice had an increased SBP between 120-150 mmHg whereas a normal SBP-range of an adult mouse is around 80-100 mmHg (294). Mid-adult mice fed a HFD had an increased SBP compared to chow fed mice (Fig. 3A). Mid-adult HFD mice had a decreased CBF within the hippocampus and thalamus, compared to both Chow and HFDB mice. Butyrate intervention restored the CBF to control levels in the hippocampus and thalamus (Fig. 3C-D).




Figure 3. Systolic blood pressure and cerebral blood flow in the cortex, hippocampus and thalamus (A-D). A) Mean systolic blood pressure (SBP) in mmHg in mid- (9 m.o.) and late (12 m.o.) adult LDLr^{-/-} mice. Six measurements were used to calculate the mean SBP per mouse. B, C, D) Mean cerebral blood flow (CBF) right and left cortex (B), hippocampus (C) and thalamus (D) in mL/100 g tissue/min. Data are presented in mean \pm SEM. * $P < 0.050$.

Butyrate restored functional connectivity in mid-adult HFD fed mice

Mid-adult HFD mice had a decreased FC within the somatosensory cortex and hippocampus compared to Chow mice (Fig. 4A-C). Butyrate intervention was able to restore impairments in FC induced by HFD feeding in mid-adult mice (Fig. 4D-F). FC results of late-adult mice are represented in supplementary figure 3. DTI was assessed to indicate diet effects on gray and white matter integrity. No diet effects were found in the mid-adult mice, whereas late adult mice showed a reduced white and gray matter integrity (Supplementary Fig. 4).




Figure 4. resting state fMRI (rsfMRI) of mid-adult mice (9 m.o.). A, B, D and E) Total correlation matrixes of the Chow (A), HFD (B & D) and HFDB (E) diet group of mid-adult mice (9 m.o.). The selected brain regions (dorsal hippocampus (DH), ventral hippocampus (VH), auditory cortex (AC), motor cortex (MC), somatosensory cortex (SSC) and visual cortex (VC)) are subdivided in left hemisphere region (first row) and right hemisphere region (second row). In general the DH and VH are strongly bilaterally connected, and AC, MC and SSC are highly interconnected. A higher Z-score (red) indicates a stronger functional connectivity (FC). C & F) Statistical analyses of FC to reveal a significant diet effect, which are represented as a matrix. Yellow: $P < 0.05$. More specifically, mid-adult mice on HFD showed reduced FC within the somatosensory cortex and hippocampus when compared to mice on Chow diet (A-C). In addition, butyrate intervention was able to restore the loss of FC mainly within the hippocampus and somatosensory cortex in mid-adult mice (D-F). Chow, $n=10$; HFD, $n=10$ and HFDB, $n=9$.

Butyrate decreased neuroinflammation in mid-adult HFD fed mice

IBA-1 immunohistochemistry indicated the level of activated microglia as measure of neuroinflammation. In the hippocampus and thalamus, the number of activated microglia was increased in mid-adult HFD mice compared to the chow fed mice (Fig. 5E-G). A butyrate intervention was able to decrease the number of activated microglia in the total hippocampus, cornu ammonis 1 (CA1) and thalamus (Fig. 5E-G). qRT-PCRs for tumor necrosis factor- α , interleukin-6 and β to analyze inflammatory levels in the hippocampus revealed no significant effects. We analyzed mRNA levels of synaptic markers: synaptophysin and postsynaptic density

protein-95 (PSD-95). In the hippocampus of mid-adult mice synaptophysin mRNA levels were higher in HFD than in chow and HFDB fed mice. No diet effects were found for PSD95 mRNA levels (Supplementary table 4 and Fig. 5).




Figure 5. Quantification of activated microglia as visualized by IBA-1 immunohistochemistry (E-H). E-G) Number of activated microglia within the hippocampus, cornu ammonis 1 (CA1) and thalamus in mid- (9 m.o.) and late (12 m.o.) adult mice. H) Visualization (late adult HFDB) of IBA-1 immunohistochemistry, positive for activated microglia, within the hippocampus at 5x and 40x magnification. Data are presented as mean \pm SEM. * $P < 0.050$.

HFD caused shifts in gut microbiota composition

We observed dramatic differences in microbiota at genus level between chow and HFD fed mice based on Bray-Curtis dissimilarity index regardless of age (Supplementary Fig. 6A). Butyrate intervention in both mid- as well as late adult mice resulted in distinct changes in microbiota composition (Supplementary Fig. 6B-C).

Butyrate-related changes in microbiota are associated with liver fibrosis and neuroinflammation

Pearson correlation analyses were used to identify a possible correlation of key pathological features (i.e. liver fibrosis in late adult mice and neuroinflammation in mid-adult mice) with the observed changes in fecal microbiota. Twelve bacterial genera showed significant correlation with liver fibrosis (Corr. c coefficient cut-off 0.4; $P < 0.05$; Fig. 6A-C). The abundance of *Clostridium Sensu Stricto* ($R = 0.59$; $P = 0.003$), *Bifidobacterium* ($R = 0.57$; $P = 0.005$) and *Enterorhabdus* ($R = 0.56$; $P = 0.006$) showed a positive correlation while *Clostridium XIVb* ($R = -0.50$; $P = 0.01$) showed a

negative correlation with liver fibrosis. Thirteen bacterial genera showed significant correlation with neuroinflammation (Corr. coefficient cut-off 0.4; $P < 0.05$; Fig. 6D-F). The abundance of *Dorea* ($R = 0.70$; $P < 0.001$), *Clostridium IV* ($R = 0.55$; $P = 0.007$) showed a positive correlation while *Clostridium Sensu Stricto* ($R = -0.58$; $P = 0.003$) showed a negative correlation with neuroinflammation.




Figure 6. Fecal microbiota changes associated with liver fibrosis and neuroinflammation.

A) Pearson correlation analysis was performed between fecal microbiota at the genus level and pathological liver fibrosis in late adult mice. B-C) examples of two bacterial genera that are correlated with liver fibrosis. D) Pearson correlation analysis was performed between fecal microbiota at the genus level and level of neuroinflammation in mid-adult mice. E-F) examples of two bacterial genera that are correlated with neuroinflammation. Data are presented in mean \pm SEM. * $P < 0.05$.

Discussion

In this study, we demonstrated that butyrate intervention is able to restore HFD induced changes in body weight, adiposity, liver pathology, cerebral blood flow, functional connectivity and neuroinflammation in mid-adulthood. In late adulthood, both HFD and butyrate did not affect brain function, which could be caused by irreversible aging processes in late adulthood e.g. in the vasculature.

HFD feeding increased adiposity and body weight in LDLr^{-/-} mice in this study, and these findings are repeatedly confirmed in rodent research (21, 295, 296). A butyrate intervention of two months restored these changes, and likewise others found that butyrate is able to reduce adiposity and body weight in obese mice (21, 22). Others found an increased activity in butyrate fed mice at night (21). We did not observe an increased activity in butyrate fed mice, like distanced swum or walked in respectively the MWM or open field test (results not shown) during day time, and unfortunately we did not examine night-time activity. Butyrate is thought to be involved in lipid metabolism by regulating and slowing down intestinal fat transport (297). Understanding of the molecular mechanism is still incomplete, but short chain fatty acids (SCFAs), like butyrate, can contribute to fatty acid oxidation in multiple tissues and decrease fat storage in white adipose tissue as butyrate can induce a switch from lipid synthesis to utilization (298). Typically, this metabolic adjustment involves a downregulation of processes orchestrated by the peroxisome proliferator-activated receptor- γ (PPAR γ) (299). An intake of butyrate affects the metabolic adaptations to obesogenic diet feeding via its stimulating effects on fatty acid oxidation and utilization. In future research physical activity levels and energy expenditure of these mice should be measured during day and night.

Obesity and non-alcoholic fatty liver disease (NAFLD) are marked by a state of inflammation initiated by HFD feeding (17). Moreover, average liver weight increased after HFD feeding in rats, and fat and cholesterol content in the liver was four to five times higher as the liver serves an alternative depot for free fatty acids (24). In mid- and late adult mice the liver weight, hepatic steatosis and hypertrophy was increased after six months of HFD feeding, and a butyrate intervention of two months was able to reverse these processes. A HFD induces a metabolic overload and thereby the metabolic pressure on the liver, resulting in alternative fat storage and NAFLD, whereas a butyrate intervention can restore metabolic pressure and adaptations.

An obesity-inducing diet diminishes hippocampal dependent search strategies and can generate cognitive deficits in rats (296). Reduced expression of synaptic proteins are suggested to be an underlying mechanism for this cognitive dysfunction (296). We found an increase in mRNA levels of the presynaptic marker synaptophysin

in mice on HFD, suggesting a possible mechanism to compensate a shortage of synaptophysin protein, and maybe the formation of presynapses. Butyrate intervention improved the spatial learning and memory performance in the Morris water maze, and preserved presynaptic synaptophysin mRNA levels. It has been reported that butyrate administration improved associative memory and learning function (283, 300). Butyrate most likely affects cognitive function via enhancing histone acetylation by inhibiting histone deacetylase (HDAC) (283). Dynamic changes in histone acetylation have been linked to the genetic programming required for memory formation (283). For instance, sodium butyrate is a HDAC inhibitor and adaptations in histone acetylation lead to neuroprotective and neuroregenerative changes in animal models for neurodegenerative diseases (283, 300).

Obesity and in particular central obesity, has been consistently associated with hypertension in humans (301). In rodents, the SBP increased after 6 months on high fat and high sucrose diet in an obese mouse model (302). We found that a HFD increased the SBP in mid-adult LDLr^{-/-} mice. An LDLr^{-/-} mouse model is prone to develop narrowed blood vessels, arterial stiffness, atherosclerosis and hypertension especially when fed a HFD, because oxidized LDL particles accumulate in the intima layer of arteries (303). Late adult mice had a high SBP, independent of diet, suggesting that aging processes result in structural changes within the vasculature and reaching a maximum in SBP.

To this date, little research has been performed using neuroimaging on brain regions of the LDLr^{-/-} mouse model. We gained novel insights using translational MRI techniques. Firstly, we revealed that cerebral blood flow (CBF) is decreased by a HFD in mid-adult LDLr^{-/-} mice. In humans it is well established that overweight and obese individuals often have a decreased CBF, gray and white matter volumes (18, 89, 90). Secondly, we have demonstrated that a HFD decreases functional connectivity (FC) between the somatosensory cortex and hippocampus in mid-adult mice. We assume that the adaptations in CBF and FC could be provoked by arterial wall stiffening, increased blood pressure and the development of hypertension induced by HFD. It can affect the cerebrovasculature and modify CBF and via blood oxygen level-dependent (BOLD) signaling it may affect the FC. Butyrate neutralized the changes in CBF and improved the impaired FC induced by a HFD. Butyrate is defined as a classic anti-angiogenic agent, however in low concentrations it can stimulate angiogenesis (304) and thereby affect BOLD signaling. In late adult mice, we found no HFD effects on SBP, CBF, FC and spatial memory. An exact mechanism for this phenomenon is unknown, although it is possible that the vascular vessels and capillaries may already have reached a pronounced dysfunctional state due to aging that cannot be further aggravated by HFD. It is known that old LDLr^{-/-}

mice on chow diet develop aortic plaques and atherosclerosis only due to aging (305). Moreover, the rate of development of atherosclerosis is more rapid between 10 and 14 months old LDLr^{-/-} mice compared to older and younger LDLr^{-/-} mice (306). Plasma cholesterol levels and vascular cell adhesion molecule 1 (VCAM-1) are possible contributors as their expression increases with age (306). We suggest that these aging processes induce a maximum, stationary, detrimental state within the vasculature. Furthermore this detrimental state cannot be further deteriorated by HFD nor improved by the dietary butyrate intervention tested herein. Mid-adult mice which have not reached this detrimental, potentially irreversible, state yet and may therefore be responsive to HFD as well as dietary intervention.

During mid-adulthood, HFD induced neuroinflammation within the thalamus and hippocampus. In line with this, a HFD increased the activation of microglia and inflammatory processes (295, 307). A butyrate intervention attenuated the inflammatory processes by decreasing the number of activated microglia within the thalamus, cortex and hippocampus, which confirmed earlier findings in literature (308). Butyrate has anti-inflammatory properties, and in more detail, butyrate activates free fatty acid receptor 3 (FFAR3) and 2 (FFAR2) (309-311) and therefore provide a link between butyrate and the central nervous system (CNS).

Analysis of the gut microbiome revealed pronounced changes in composition upon HFD feeding. These HFD induced changes were very consistent with observations made by others (312). The pronounced difference between chow and HFD composition made it difficult to discriminate the specific effects of butyrate from those of HFD on the microbiome. In line with observations made in obese humans which exhibit decreased levels of butyrate-producing bacteria compared to lean individuals (24), we found HFD induced in *Clostridium XIVa* abundancy. In an attempt to link the changes in microbiota with relevant pathophysiological endpoints, we tested whether specific genera were correlated to liver fibrosis and neuroinflammation. We observed a positive correlation between neuroinflammation and bacteria abundance of the genus *clostridia* (e.g. *Dorea*, *Clostridium IV*, *Lachnospiraceae*, *Lachnospiraceae incertae sedis*, and *Clostridium XIVa*) in fecal microbiota of mid-adult mice. A decreased abundance of these bacteria in HFDB fed mice correlated with a reduction in neuroinflammation as compared to HFD fed mice. Bacteria from the *clostridia* genus are known to be butyrate producing bacteria and are very abundant in the human colon (313). Our data suggest that dietary supplementation of butyrate in mid-adult mice could affect neuroinflammation despite a reduction of butyrate-producing bacteria in the fecal microbiota. Interestingly, this effect was less clear in the late adult mice in relation to hepatic fibrosis development where we observed both positive (*Clostridium sensu stricto*, *Clostridium XI*) as well as a negative correlation (*Clostridium XIVb*,

Clostridiales, *Clostridium XIVa*) of the butyrate-producing bacteria of the *Clostridia* genus. To our knowledge, these findings are novel however the data should be interpreted with caution because microbiome analysis and pathology have been performed at the same time point and microbiome changes may merely reflect metabolic adaptation but might not be causal to development of the end points. It is intriguing that the microbiota changes upon butyrate treatment differ in mid-adult and late adult mice and that the two age groups also develop different pathologic endpoints.

Conclusion

We hypothesize that midlife obesity increases the risk of cognitive impairment via mainly inflammatory responses and vascular damage. These findings emphasize the association between midlife obesity and cognitive impairment in humans (3). We report novel insights in the impact of diet-induced obesity and butyrate in a genetic mouse model on brain function and structure during mid- and late adulthood. In mid-adult humans, butyrate supplementation may serve as potential preventative for obesity and obesity-related diseases, and in the long-run as preventative for dementia.

Supplemental data:

<https://www.nature.com/articles/ijo201752#supplementary-information>

4

ADIPOSITY IS ASSOCIATED WITH VASCULAR AND VOLUMETRIC BRAIN OUTCOMES IN CEREBRAL SMALL VESSEL DISEASE.

THE RUN DMC STUDY

Arnoldussen I.A.C.¹

Gustafson D.R.¹

M.C. Leijssen E.M.C.

de Leeuw F.E.

Kiliaan A.J.

(2018) Submitted to Neurology.

¹ Authors contributed equally to the present work.

Abstract

Adiposity and changes in adiposity are associated with cerebrovascular diseases, such as cerebral small vessel disease (CSVD) and its sequelae, vascular cognitive impairments (VCI) and dementias. Body mass index (BMI), waist circumference (WC), and blood leptin and total adiponectin levels were associated with CSVD and brain volumetry in 503 adults age ≥ 50 years enrolled in the Radboud University Nijmegen Diffusion tensor and Magnetic resonance imaging Cohort (RUN DMC).

RUN DMC participants were followed for 9 years, from 2006 to 2015. BMI, WC, brain imaging and dementia diagnoses were evaluated at baseline and follow-up. Adipokines were measured at baseline. Brain imaging outcomes included CSVD components, white matter hyperintensities, lacunes and microbleeds; and gray and white matter, hippocampal, total brain, and intracranial volumes.

Cross-sectionally among men at baseline, higher BMI, WC and leptin were associated with lower gray matter and total brain volumes, and higher BMI and WC were associated with lower hippocampal volume. At follow-up 9 years later, cross-sectional analyses showed associations between higher BMI and lower gray matter volume in men, and that an obese WC ($>102\text{cm}$) was protective for presence of ≥ 1 lacune or ≥ 1 microbleed in men. In women, increasing BMI and overweight or obesity ($\text{BMI} \geq 25 \text{ kg/m}^2$ or $\text{WC} > 88 \text{ cm}$) were associated with ≥ 1 lacune.

Anthropometric and metabolic adiposity indicators were differentially associated with CSVD and brain volumes by sex. Adiposity is associated with a vascular-neurodegenerative spectrum that may influence and be influenced by brain pathologies associated with VCI and dementia.

Introduction

Approximately 50% of the global adult population is overweight or obese according to 2016 estimates, creating an obesity pandemic (1). Obesity during middle-age is associated with higher risk of several adult life cardiovascular risk factors and events including hypertension, hyperlipidemias, type 2 diabetes mellitus (T2DM), atherosclerosis, and myocardial infarction. These cardiovascular risk factors are subsequently associated with cerebrovascular events such as stroke and late-onset, sporadic dementias (315), in particular vascular cognitive impairments (VCI) and vascular forms of dementia (316).

The biological mechanisms linking obesity to VCI and dementias and underlying cerebrovascular and neuro-pathologies are not well understood. However, one potential mechanism is that obesity represents higher amounts of adipose tissue, which contribute to alterations in the peripheral and cerebral circulation, which may be more pronounced with aging (317). In the brain, this translates to decreased cerebral blood flow and cerebral small vessel disease (CSVD), which is considered the major vascular contributor to dementia (318, 319). While CSVD cannot be diagnosed *in vivo*, it is operationalized by the presence of brain MRI markers including white matter hyperintensities (WMH), lacunes and microbleeds.

Adipose tissue is the largest endocrine organ in the human body (77). Anthropometric measurements such as body mass index (BMI) and waist circumference (WC) are commonly used to estimate total and central adiposity, respectively. BMI grossly reflects total adult adiposity (5), whereas WC reflects the amount of highly bioactive visceral adipose tissue that surrounds internal organs (320). Secretory products of adipose tissue exhibit vascular and metabolic effects, and influence brain structure and function (77, 112, 321).

We examined the associations of anthropometric and metabolic measures of adiposity with MRI markers of CSVD (WMH, lacunes and microbleeds) and brain volumetry (gray and white matter, hippocampus and total brain). Given known sex differences in total and central adiposity, we stratified all analyses by sex. Identifying and understanding potential associations between potentially modifiable adiposity and CSVD, both of which increase risk for cognitive impairments and dementias, is important as this could facilitate interventions to reduce adverse effects of excess adiposity on public and personal health.

Methods

Study population

The RUN DMC study prospectively investigates vascular and other factors associated with CSVD and CSVD progression among a baseline sample of 503 adults, aged 50 to 85 years, with CSVD. Participants were recruited from consecutive patients referred to the department of Neurology of the Radboud university medical centre between October 2002 and November 2006, because they presented with symptoms of CSVD. These symptoms were acute, such as transient ischemic attack (TIA) or lacunar syndromes, but also subacute manifestations such as cognitive and motor (gait) disturbances (322). Patients were included based on age and the presence of CSVD on neuroimaging (WMH and/or lacunar infarcts) (322). Patients who were eligible because of a lacunar syndrome were included >6 months after the event to avoid acute effects on outcomes.

Exclusion criteria were: (a) dementia; (b) parkinsonism; (c) intracranial hemorrhage; (d) life expectancy <6 months; (e) intracranial space-occupying lesion; (f) psychiatric or other disease interfering with cognitive testing or ability to be followed up; (g) recent or current use of acetylcholinesterase inhibitors, neuroleptic agents, L-dopa or dopamine agonists or antagonists; (h) non-CSVD WMH (e.g. multiple sclerosis); (i) prominent visual or hearing impairment; (j) language barrier; or (k) MRI contraindications, e.g., claustrophobia.

Follow-up of participants after 9 years occurred among those who were alive and agreed to participate. Baseline participants were invited to follow-up via mail, with a follow-up telephone call to schedule their visit at our research center. During their follow-up visit, a subset of baseline assessments were re-administered, including a dementia assessment. All tests at baseline and follow-up were performed by two trained Neurology residents physicians.

All participants provided written informed consent. The study was approved by the Medical Review Ethics Committee region Arnhem-Nijmegen.

Structured interview

A structured interview was done, which included on demographics, lifestyle, vascular risk factors, and current medication use. *Demographics and lifestyle* included education (classified using 7 categories, ranging from primary school to an academic degree), marital status, living conditions, and lifestyle habits (alcohol consumption, smoking). *Vascular risk factors and cardiovascular disease* included history of hypertension, T2DM, atrial fibrillation, TIA, stroke, myocardial infarction, coronary artery bypass graft, per-cutaneous transluminal coronary angiography, aortic prosthesis, vascular prosthesis, carotid endarterectomy, and migraine. Family history of myocardial infarction, cerebrovascular disease and

diabetes mellitus was recorded. *Current medication* use was classified according to the anatomical therapeutic chemical (ATC) classification system (323).

Physical examination

Anthropometric measurements included body mass index (BMI), calculated as clinically measured body weight divided by body height squared (kg/m^2). Maximal waist circumference (WC) in centimeters (cm) was measured without a shirt, in a standing position, between the lowest rib and the iliac crest at the end of usual expiration.

Blood pressure measurements included systolic and diastolic pressures obtained in the sitting position. The average of three repeated measurements was used.

Blood measures at baseline were collected after a 12-hour, overnight fast, and serum aliquots were stored at -80°C . A lipid panel comprised of total cholesterol, HDL, LDL and triglyceride, was measured. Total serum leptin and adiponectin levels were determined using human leptin and adiponectin enzyme-linked immunosorbent assays (ELISA) (Laboratory Medicine, Radboud University Medical Center, Nijmegen, The Netherlands).

Magnetic Resonance Imaging protocol

MRI images were acquired on a 1.5T MRI scanner (2006: Siemens (Munich, Germany), Magnetom Sonata, and 2015: Siemens, Magnetom Avanto) using the same head coil at both time points. The scanning protocol included 3D T1 magnetization-prepared rapid gradient echo (MPRAGE) imaging (voxel size $1.0 \times 1.0 \times 1.0$ mm); and fluid-attenuated inversion recovery (FLAIR) pulse sequences (2006: voxel size $0.5 \times 0.5 \times 5.0$ mm, interslice gap 1.0 mm, and 2015: voxel size $0.5 \times 0.5 \times 2.5$ mm; interslice gap 0.5 mm). To minimize effects of changes in FLAIR sequence, we re-sliced follow-up FLAIR images to match slice thickness of baseline images using linear interpolation (324).

Brain volumetry outcomes

MRI outcomes analyses have been described in detail (324). In brief, gray matter, white matter, hippocampal and cerebrospinal fluid (CSF) volumes were calculated using SPM12 (fil.ion.ucl.ac.uk/spm/) unified segmentation routines on the T1 MPRAGE images (325). All images were visually checked for co-registration and segmentation artefacts. Gray matter volumes (GMV), white matter volumes (WMV), and CSF volumes were computed by summing all voxels belonging to that tissue class, multiplied by voxel volume in milliliters. Intracranial volume (ICV) was determined by summing GMV, WMV, and CSF volume; and total brain volume (TBV) by summing GMV and WMV. To account for interscan effects, we corrected for differences in ICV between baseline and follow-up. All volumes were normalized to baseline ICV to account for head size.

CSVD outcomes

CSVD was rated using STRIVE criteria for vascular changes on neuroimaging (326). WMH volumes were calculated using a semiautomatic WMH segmentation method (327). Segmentations were visually checked for segmentation errors by one trained rater, blinded to clinical data. The Fazekas scale was used to categorize baseline WMH severity (328).

Number and location of lacunes and microbleeds were rated manually on FLAIR/T1-weighted MRI scans according to the STRIVE criteria by 2 trained raters blinded to clinical data. Lacunes were defined as hypo-intense areas >2 mm and ≤ 15 mm on FLAIR and T1, ruling out enlarged perivascular spaces (≤ 2 mm, except around the anterior commissure, where perivascular spaces can be large) and infraputaminar pseudolacunes (329). Microbleeds were defined as small (<10 mm diameter), homogeneous, round foci of low signal intensity on T2* weighted images (330). Microbleed number per hemisphere was counted. Lesions were not considered microbleeds if they were symmetric hypointensities in the globus pallidus, likely calcifications or iron deposits (330). Inter-rater and intra-rater reliabilities were excellent (324). Incident lacunes and microbleeds during the follow-up period were identified (324).

Dementia assessment

Dementia screening was performed at 9-year follow-up and has been described in detail (325). The mini-mental state examination (MMSE)(331) was the initial screening instrument of global cognition. A follow-up MMSE score <26 or a decline of ≥ 3 points from baseline was considered screening positive. Those who screened positive based on these criteria were subsequently examined for the presence of dementia at the Radboud Alzheimer Center Nijmegen, The Netherlands. If a participant refused additional examinations at the Alzheimer Center, a consensus panel consisting of a neurologist, clinical neuropsychologist, and a geriatrician made the dementia diagnosis. This panel reviewed all available neuropsychological (322) and brain imaging information, which included a) difference in neuropsychological performance between baseline and follow-up; b) outcome of the Mini International Neuropsychiatric Interview MINI (332); c) the follow-up MRI scan and/or baseline MRI-scan to determine dementia subtype; and d) medical record review. In addition, participants' general practitioners and medical specialists were contacted regarding cognitive status. Cognitive tests were interpreted with consideration for age, level of education, and interference with daily living confirmed by family or caregiver (333). The dementia diagnosis was based on the diagnostic and statistical manual of mental disorders (DSM) IV criteria (334). Dementia onset was established as the date when clinical symptoms were consistent with the diagnosis (335).

Statistical analyses

Baseline and follow-up cross-sectional analyses included descriptive analyses of all primary adiposity exposures and brain outcomes for the entire sample and by sex. Means and standard deviations (SD) were computed for continuous variables. Due to skewed distributions, leptin, adiponectin and WMH were natural log transformed. As leptin resistance is often observed in obesity, we estimated leptin resistance by calculating the leptin to BMI (leptin:BMI) ratio (336). Pearson correlation analyses were conducted among adiposity measures. T-tests were used for means comparisons. Chi-square analyses were conducted for categorical variables.

Multivariable linear and logistic regression models were used to examine cross-sectional associations between adiposity measures and CSVD and volumetric brain outcomes at both time points, 2006 and 2015. Regression models were run separately for each adiposity measure as an independent predictor. Adiposity measures included BMI and WC in 2006 and 2015; and leptin and total adiponectin in 2006 only. Brain imaging outcomes included CSVD components and brain volumes. Logistic regression analyses were used to predict the presence *versus* absence of CSVD components. Odds ratios (OR) and 95% Confidence Intervals (CI) were calculated. WMH, GMV, WMV, TBV and hippocampal volume (HV) were considered as continuous outcome variables in linear regression analyses. Beta-coefficients (b) and 95% CI were calculated.

Given that CSVD and more severe change in body weight and BMI increases risk for dementia (337), we also examined change in anthropometric measures (BMI and WC) from 2006-2015 in association with MRI markers of CSVD and brain volumes measured in 2015. In analyses of anthropometric change, since no participant presents with the same body weight or BMI at two visits and 'any change' may not be informative, we categorized the sample by an *a priori* defined 'substantial' degree of change, either $\geq 5\%$ and $\geq 10\%$ increase or decrease from baseline body weight and BMI.

Selection of covariates originated from a pool of variables that were potentially biologically relevant and/or reported in the literature as CSVD risk and protective factors. Our pool consisted of age; sex; educational attainment (\leq primary education *versus* $>$ primary education); cigarette smoking in pack-years (calculated as number of packs of cigarettes smoked per day multiplied by number of years smoked); alcohol intake (units per day); clinically relevant depressive symptom burden (≥ 16 on the center of epidemiologic studies on depression scale (CES-D) or current depression medication use); MMSE score; use of glucose lowering medications; T2DM (self-reported or T2DM medication use); hypertension (blood pressure

$\geq 140/90$ mmHg or antihypertensive medication use); blood lipid levels; and hypercholesterolemia (cholesterol ≥ 6.2 mM or lipid-lowering drug use). From this pool, we systematically determined which ones to include in multivariable-adjusted regression models by evaluating each variable in age-adjusted regression models in association with brain MRI outcomes. If the variable was associated with an outcome at $P \leq 0.05$, the variable was included as a covariate in the final regression model. As a result, age, sex, education (\leq primary school versus $>$ primary school), cigarette smoking (ever/never), and T2DM (yes/no) were included. In analyses involving 2015 brain outcomes, presence of dementia and baseline WMH severity based on the Fazekas scale, were also included as covariates. Results for linear and logistic multivariable analyses were considered significant using two-tailed tests, $P < 0.05$. SPSS, version 22.0 (IBM Corporation, Armonk, New York, USA), was used to conduct data analyses.

Results

Baseline characteristics of the RUN DMC baseline study population ($N=503$) are reported in Table 1. Mean age of participants at baseline was 65.6 ± 8.8 years, and at follow-up, 71.3 ± 7.9 years. Differences in baseline characteristics between participants and those who died during follow-up are also presented in Table 1. Compared to surviving participants, those who died during follow-up were older; had a lower MMSE score; had higher average WC; had higher prevalence of central obesity using sex-specific cut points; were more likely to use lipid- glucose- and blood pressure-lowering medications; and were more likely to have T2DM. Those who died also evidenced lower baseline TBV, GMV, WMV and HV; a higher median volume of WMH; and were more likely to have a Fazekas score of moderate or severe (*versus* none or mild) or possess ≥ 1 lacune at baseline. Thus, ‘healthier’ surviving participants were observed in 2015.

Adiposity indicators, CSVD markers, and brain volumes

Increasing BMI and WC, as well as total and central obesity were associated with lower GMV and TBV in all male participants cross-sectionally at average baseline age, 66 years (Table 2). Higher serum leptin levels were associated with lower GMV and TBV in the total sample of men and women. Increasing BMI and WC were also associated with lower HV in men (Table 2). Interestingly, a higher estimate of leptin resistance (leptin: BMI), was also associated with lower GMV and TBV, particularly in men. At follow-up 9 years later, average age 71 years, cross-sectional analyses showed increasing BMI associated with lower GMV; and higher WC with lower HV, particularly in men (Table 3).




Figure 1. RUN DMC study design

In relation to CSVD outcomes, cross-sectionally at average age 66 years, leptin was protective in men. In particular, the highest quartile of leptin, compared to the lowest quartile, was associated with lower WMH volumes and lower odds of ≥ 1 lacune. Cross-sectional analyses at 9-year follow-up, showed that an obese WC lowered the odds of microbleeds and lacunes in men (Table 4). However, in women, increasing BMI, being overweight or obese based on BMI, or having an obese WC were associated with the presence of ≥ 1 lacune (Table 4).

Longitudinal analyses of baseline adiposity measures in association with CSVD and brain volumetry outcomes at follow-up 9 years later showed no associations (data not shown). Similarly, evaluation of 5% and 10% change in BMI and body weight over 9 years, showed no association with follow-up CSVD or brain volumetry outcomes (data not shown).

	Baseline 2006			Deaths during follow-up			Follow-up 2015		
	All N=503	Men N=284	Women N=219	All N=85	Men N=47	Women N=38	All N=287	Men N=164	Women N=123
Demographics									
Age (years), mean \pm SD	65.6 \pm 8.8	65.7 \pm 8.8	65.5 \pm 8.9	72.4 \pm 8.1 ^c	72.8 \pm 6.6 ^c	71.9 \pm 9.7 ^c	71.3 \pm 7.9	71.3 \pm 7.9	71.0 \pm 7.4
Education <8 years, N (%)	49 (9.7)	26 (9.2)	23 (10.5)	13 (15.3) ^c	6 (12.8)	7 (18.4) ^c			
Depressive symptoms \geq 16, N (%)	167 (33.2)	81 (28.5)	86 (39.3) ^b	50 (58.8)	20 (52.6)	19 (28.4) ^{b,c}			
MMSE score, mean \pm SD	28.1 \pm 1.6	28.1 \pm 1.7	28.2 \pm 1.6	27.3 \pm 1.9 ^c	27.1 \pm 1.9 ^c	27.6 \pm 1.9 ^c	27.5 \pm 3.5	28.1 \pm 2.0	28.3 \pm 2.0
Dementia incidence, N (%)				23 (27.1)	9 (19.1)	14 (36.8)	14 (4.9)	10 (6.1)	4 (3.3)
Vascular risk factors^a									
Alcohol consumption (units per week), mean \pm SD	7.9 \pm 9.3	10.8 \pm 10.3	4.1 \pm 6.2 ^b	8.4 \pm 10.8	12.9 \pm 12.5	2.8 \pm 3.8 ^a			
Smoking (ever), N (%)	353 (70.2)	236 (83.1)	117 (53.4) ^b	66 (77.6)	45 (95.7)	21 (55.3) ^a			
Lipid lowering drugs, N (%)	237 (47.1)	143 (50.4)	94 (42.9)	51 (60.0) ^c	30 (63.8)	21 (55.3)			
Glucose lowering drugs, N (%)	66 (13.1)	45 (15.8)	21 (9.6) ^b	22 (25.9) ^c	14 (29.8) ^c	8 (21.1) ^{b,c}			
Hypertension, N (%)	369 (73.4)	208 (73.2)	161 (73.5)	73 (85.9) ^c	41 (87.2)	32 (84.2)			
Diabetes mellitus, N (%)	75 (14.9)	50 (17.6)	25 (11.4) ^b	23 (25.9) ^c	15 (31.9) ^c	8 (21.1) ^{b,c}			
Adiposity indicators									
BMI (kg/m ²), mean \pm SD	27.2 \pm 4.1	27.3 \pm 3.8	27.0 \pm 4.5	26.9 \pm 4.3	27.0 \pm 4.5	26.8 \pm 4.5	27.2 \pm 4.0	27.2 \pm 3.6	26.9 \pm 4.3
Healthy (<25.0 kg/m ²), N (%)	152 (30.2)	78 (27.5)	74 (33.8)	29 (34.1)	78 (27.5)	74 (33.8)	90 (31.3)	48 (29.3)	42 (34.1)
Overweight (25.0-29.9 kg/m ²), N (%)	226 (44.9)	139 (48.9)	87 (39.7)	35 (41.2)	139 (48.9)	87 (39.7)	132 (45.9)	80 (48.8)	52 (42.3)
Obese (>30.0 kg/m ²), N (%)	125 (24.9)	67 (23.6)	58 (26.5)	21 (24.7)	67 (23.6)	58 (26.5)	65 (22.6)	36 (21.9)	29 (23.6)
Change in BMI, 2015-2006 (%), mean \pm SD							0.44 \pm 8.3	0.11 \pm 6.8	0.88 \pm 9.5
Waist circumference (WC, cm), mean \pm SD	96.0 \pm 12.2	99.9 \pm 10.6	91.0 \pm 12.4 ^b	97.0 \pm 12.4	101.4 \pm 10.2	91.6 \pm 12.7 ^b	99.1 \pm 12.4	102.4 \pm 11.2	93.5 \pm 12.0 ^b
Healthy WC (men < 102 cm, women < 88cm), N (%)	245 (48.7)	154 (54.2)	91 (41.6) ^b	36 (57.6)	22 (46.8)	14 (36.8) ^b	122 (42.7)	86 (52.8)	36 (30.1)
Obese WC (men \geq 102 cm, women \geq 88cm), N (%)	258 (52.3)	130 (45.8)	128 (58.4) ^b	68 (54.8)	25 (53.2)	24 (63.2) ^b	164 (57.3)	78 (47.9)	86 (69.9)
Change in WC, 2015-2006 (%), mean \pm SD							4.3 \pm 8.7	3.8 \pm 8.0	4.9 \pm 9.3
Leptin (ng/mL), mean \pm SD	23.4 \pm 16.2	13.9 \pm 10.5	35.7 \pm 22.5 ^b	26.4 \pm 23.6	15.3 \pm 11.4	39.8 \pm 27.6 ^b			
Leptin resistance (leptin:BMI)	0.8 \pm 0.6	0.5 \pm 0.3	1.2 \pm 0.7 ^b	0.9 \pm 0.8	0.5 \pm 0.4	1.4 \pm 0.9 ^b			
Adiponectin (ng/mL), mean \pm SD	4.9 \pm 3.3	3.6 \pm 2.2	6.5 \pm 3.8 ^b	6.2 \pm 4.5	3.9 \pm 2.2	9.0 \pm 5.2 ^b			
Brain volumes									
Total brain volume (mL), mean \pm SD	1060.9 \pm 80.1	1046.0 \pm 81.0	1080.0 \pm 74.9 ^b	573.0 \pm 52.7 ^c	559.9 \pm 47.9 ^c	589.2 \pm 54.4 ^c	1043.7 \pm 78.7	1025.5 \pm 80.8	1068.0 \pm 69.0 ^b
Gray matter volume (mL), mean \pm SD	606.2 \pm 52.6	593.2 \pm 51.4	623.1 \pm 49.2 ^b	427.3 \pm 45.9 ^c	420.8 \pm 53.3 ^c	435.4 \pm 33.9 ^c	597.6 \pm 50.6	584.67 \pm 47.7	617.4 \pm 47.4 ^b
White matter volume (mL), mean \pm SD	454.7 \pm 46.0	452.9 \pm 48.5	457.1 \pm 42.5	7.1 \pm 1.0 ^c	6.9 \pm 0.9 ^c	7.3 \pm 1.0 ^c	443.8 \pm 46.1	439.1 \pm 50.6	449.8 \pm 39.5
Hippocampal volume (mL), mean \pm SD	6.7 \pm 1.0	7.3 \pm 1.0	7.9 \pm 1.0 ^b				7.4 \pm 1.1	7.1 \pm 1.0	7.3 \pm 1.0 ^b
CSVD markers									
WMH (mL), median (IQR)	3.6 (10.0)	3.2 (9.6)	4.1 (10.8) ^b	11.5 (15.0) ^c	11.2 (14.4) ^c	12.6 (16.4) ^c	4.7 (10.3)	3.8 (7.6)	5.67 (13.7)
Microbleeds, N (%)	83 (16.5)	50 (17.6)	33 (15.1)	18 (21.2)	13 (27.7) ^c	5 (13.2)	70 (24.4)	38 (23.2)	32 (26.0)
Lacunae, N (%)	132 (26.2)	86 (30.3)	46 (21.0) ^b	43 (50.6) ^c	27 (57.4) ^c	16 (42.1) ^{b,c}	79 (27.5)	50 (30.4)	29 (23.6)

Table 1. Baseline and follow-up characteristics of participants, and baseline characteristics of those died during follow-up in the RUN DMC. ^aVascular risk factors assessed at baseline only. ^bSignificant at $P < 0.05$ men versus women. ^cSignificant $P < 0.05$ participants who died during follow up versus baseline. WMH: White Matter Hyperintensities. CSVD: cerebral small vessel disease. IQR: interquartile range.

Baseline 2006		Total Brain Volume		Gray Matter Volume		White Matter Volume		Hippocampal Volume	
	N	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
Continuous adiposity measures									
BMI (kg/m ²)	503	-0.05 (-0.13- 0.03)	0.21	-0.16(-0.24- -0.09)	0.00	0.07(-0.01-0.16)	0.11	-0.08(-0.16- -0.01)	0.04
BMI (kg/m ²), men	284	-0.12(-0.23-0.04)	0.00	-0.28(-0.41- -0.18)	0.00	0.08(-0.04-0.22)	0.19	-0.12(-0.26- 0.01)	0.04
BMI (kg/m ²), women	219	0.01 (-0.08- 0.09)	0.90	-0.04 (-0.14- -0.06)	0.48	0.06(-0.06-0.17)	0.35	-0.06(-0.17- 0.06)	0.38
WC (cm)	503	-0.07(-0.16-0.02)	0.11	-0.18(-0.26- -0.10)	0.00	0.05(-0.04-0.14)	0.31	-0.11(-0.20- -0.02)	0.01
WC (cm), men	284	-0.13(-0.25- -0.06)	0.00	-0.25(-0.40- -0.16)	0.00	0.03(-0.11-0.18)	0.31	-0.10(-0.26- 0.02)	0.08
WC (cm), women	219	-0.02 (-0.11- 0.08)	0.76	-0.09 (-0.19- 0.03)	0.16	0.07(-0.06-0.19)	0.32	-0.11(-0.23- 0.02)	0.10
Leptin (ng/mL)	503	-0.18(-0.26- -0.07)	0.00	-0.13(-0.22- -0.04)	0.00	0.04(-0.06-0.15)	0.42	-0.05(-0.14-0.05)	0.36
Leptin (ng/mL), men	284	-0.07(-0.19- 0.02)	0.11	-0.16(-0.32- -0.06)	0.00	0.05(-0.09-0.21)	0.43	-0.07(-0.22-0.06)	0.27
Leptin (ng/mL), women	219	-0.02 (-0.13- 0.10)	0.76	-0.05 (-0.19- -0.08)	0.44	0.02(-0.13-0.18)	0.74	-0.00(-0.15-0.16)	0.97
Adiponectin (ng/mL) ^b	503	-0.07(-0.16-0.02)	0.11	0.02(-0.06-0.10)	0.65	0.04(-0.06-0.13)	0.44	0.01(-0.08-0.10)	0.78
Adiponectin (ng/mL), men	284	0.04(-0.04-0.14)	0.28	0.06(-0.05-0.17)	0.30	0.03(-0.09-0.16)	0.62	0.05(-0.07-0.17)	0.44
Adiponectin (ng/mL), women	219	0.01(-0.10-0.12)	0.89	-0.03(-0.17-0.10)	0.60	0.04(-0.10-0.20)	0.53	-0.02(-0.19-0.13)	0.73
Leptin resistance	503	-0.16(-0.24- -0.08)	0.00	-0.11(-0.18- -0.03)	0.01	0.01(-0.08-0.09)	0.92	-0.03(-0.12-0.06)	0.51
Leptin:BMI, men	284	-0.05(-0.14-0.03)	0.20	-0.13(-0.24- -0.02)	0.02	0.04(-0.08-0.16)	0.48	-0.07(-0.19-0.05)	0.27
Leptin:BMI, women	219	-0.07(-0.16-0.04)	0.22	-0.08(-0.19-0.05)	0.24	-0.04(-0.18-0.09)	0.54	0.04(-0.10-0.17)	0.61
Categorical adiposity measures									
BMI (kg/m²)									
Total sample									
Healthy: $\leq 24.9^a$	152	1.0		1.0		1.0		1.0	
Overweight & Obese: ≥ 25.0	351	-0.03(-0.11- 0.05)	0.53	-0.09(-0.17- -0.02)	0.02	0.06(-0.03-0.15)	0.17	-0.03(-0.12-0.05)	0.44
Men vs women									
Normal weight: $\leq 24.9^a$	78/74	1.0		1.0		1.0		1.0	
Overweight & Obese: men ≥ 25.0	206	-0.07(-0.16- 0.01)	0.08	-0.16(-0.27- -0.05)	0.00	0.04(-0.08-0.17)	0.46	-0.12(-0.27-0.05)	0.17
Overweight & Obese: women ≥ 25.0	145	0.03 (-0.06- 0.12)	0.56	-0.03 (-0.11- -0.11)	0.96	0.08(-0.05-0.19)	0.25	-0.06(-0.22-0.12)	0.54
WC (cm)									
Total sample									
Healthy WC: men ≤ 102 & women $\leq 88^a$	245	1.0		1.0		1.0		1.0	
Obese WC: men > 102 , women > 88	258	-0.10(-0.18- -0.02)	0.01	-0.15(-0.23- -0.07)	0.00	0.03(-0.06-0.12)	0.53	-0.10(-0.19- -0.01)	0.02
Men vs women									
Small WC: men ≤ 102 & women $\leq 88^a$	154/91	1.0		1.0		1.0		1.0	
Large WC: men > 102	130	-0.12(-0.20- -0.03)	0.01	-0.15(-0.23- -0.07)	0.00	0.02(-0.10-0.14)	0.53	-0.13(-0.25- -0.02)	0.03
Large WC: women > 88	128	-0.03 (-0.12- 0.07)	0.60	-0.06 (-0.17- 0.06)	0.32	0.03(-0.10-0.16)	0.62	-0.07(-0.20-0.07)	0.32
Leptin (ng/mL)									
Total sample									
Q1: men ≤ 7.4 , women $\leq 19.6^a$	127	1.0		1.0		1.0		1.0	
Q4: men > 21.7 , women > 52.6	126	-0.21(-0.25- -0.07)	0.00	-0.17(-0.20- -0.04)	0.00	0.02(-0.08- 0.11)	0.74	-0.11(-0.18- -0.02)	0.10
Men vs women									
Q1: men ≤ 7.4 , women $\leq 19.6^a$	72/55	1.0		1.0		1.0		1.0	
Q4: men > 21.7	71	-0.09(-0.16- 0.03)	0.17	-0.19(-0.24- -0.02)	0.02	0.05(-0.10- 0.17)	0.74	-0.16(-0.25-0.01)	0.07
Q4: women > 52.6	55	-0.12(-0.20- 0.04)	0.18	-0.16(-0.23- 0.02)	0.10	-0.04(-0.20- 0.13)	0.68	-0.02(-0.19-0.15)	0.82

Table 2. Linear regression models estimating baseline cross-sectional associations between adiposity exposures and brain volumes in RUN DMC. Linear regression models for gray matter, white matter and hippocampal volumes were adjusted for age, education, sex, smoking, and type 2 diabetes mellitus (T2DM). N: number of participants. Seven participants were excluded in the hippocampal analyses due to imaging artifacts (motion). β : beta, standardized regression coefficient. Continuous measures of total brain volume, gray matter volume, white matter volume and hippocampal volume (mL). Continuous levels of leptin and adiponectin are categorized in quartiles, and the lowest quartile (Q1) was compared to the highest quartile (Q4). ^aReferent category. ^bNo significant associations were observed for adiponectin quartiles (data not shown).

Follow-up 2015		Total brain volume		Gray matter volume		White matter volume		Hippocampal volume	
	N	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
Continuous adiposity measures									
BMI (kg/m ²)	287	-0.04(-0.12- 0.05)	0.38	-0.12(-0.23- -0.02)	0.02	0.06(-0.05-0.18)	0.29	-0.02(-0.11- -0.08)	0.71
BMI (kg/m ²), men	164	-0.05(-0.18- 0.07)	0.41	-0.14(-0.29- -0.03)	0.04	0.05(-0.11-0.24)	0.45	0.05(-0.09-0.19)	0.48
BMI (kg/m ²), women	123	-0.03(-0.14-0.09)	0.69	-0.09(-0.22-0.07)	0.30	0.05(-0.10-0.19)	0.56	-0.08(-0.21-0.07)	0.32
WC (cm)	286	-0.05(-0.14- 0.04)	0.26	-0.13(-0.24- -0.01)	0.03	0.05(-0.08-0.18)	0.46	-0.03(-0.14- -0.07)	0.56
WC (cm), men	163	-0.04(-0.18- 0.08)	0.44	-0.11(-0.26- 0.04)	0.16	0.02(-0.16-0.21)	0.80	0.06(-0.09-0.21)	0.42
WC (cm), women	123	-0.05(-0.18- 0.08)	0.44	-0.16(-0.31- 0.01)	0.07	-0.12(-0.28-0.03)	0.11	-0.12(-0.28-0.03)	0.11
Categorical adiposity measures									
BMI (kg/m²)									
Total sample									
Normal weight: $\leq 24.9^a$	88	1.0		1.0		1.0		1.0	
Overweight & Obese: ≥ 25.0	199	-0.03(-0.08- 0.08)	0.94	-0.09(-0.19- -0.01)	0.08	0.08(-0.04-0.19)	0.19	0.00(-0.09-0.10)	0.93
Men vs women									
Normal weight: men/women $\leq 24.9^a$	47/41	1.0		1.0		1.0		1.0	
Overweight & Obese: men ≥ 25.0	117	0.07(-0.11- 0.12)	0.89	-0.09(-0.19- -0.01)	0.08	0.08(-0.05-0.26)	0.17	0.03(-0.10-0.15)	0.70
Overweight & Obese: women ≥ 25.0	82	-0.01(-0.13-0.11)	0.86			0.05(-0.11-0.19)	0.95	-0.01(-0.15-0.14)	0.95
WC (cm)									
Total sample									
Small WC: men ≤ 102 , women $\leq 88^a$	122	1.0		1.0		1.0		1.0	
Large WC: men >102 , women >88	164	0.03(-0.16- 0.18)	0.94	-0.08(-0.37- -0.06)	0.18	0.07(-0.10-0.38)	0.26	-0.05(-0.10- 0.29)	0.34
Men vs women									
Small WC: men ≤ 102 , women $\leq 88^a$	85/37	1.0		1.0		1.0		1.0	
Large WC: men >102	78	0.01(-0.11- 0.12)	0.89	-0.08(-0.37- -0.06)	0.18	0.08(-0.14-0.50)	0.28	0.14(-0.01-0.52)	0.04
Large WC: women >88	86	-0.01(-0.13-0.11)	0.86	-0.06(-0.20-0.09)	0.46	0.07(-0.22-0.48)	0.47	-0.06(-0.47-0.19)	0.41

Table 3. Linear regression models estimating cross-sectional associations between anthropometric measures and brain volumes at 9-year follow-up in RUN DMC. Linear regression models for gray matter volume, white matter volume and hippocampal volume were adjusted for: age, education, sex, smoking, type 2 diabetes mellitus (T2DM), WMH severity, and dementia incidence. N: number of participants. Five participants were excluded in the hippocampal analyses due to imaging artifacts (motion). β : beta, standardized regression coefficient. Continuous measures of total brain volume, gray matter volume, white matter volume and hippocampal volume (mL). ^aReferent category.

Follow-up 2015	WMH ^b			≥ 1 Microbleeds ^c			≥ 1 Lacunes ^c	
	N	β (95% CI)	p	N	OR (95% CI)	p	OR (95% CI)	p
Continuous adiposity measures								
BMI (kg/m²)	287	0.05(-0.05-0.15)	0.31	276	0.95(0.88-1.03)	0.20	1.03(0.95-1.11)	0.51
BMI (kg/m ²), men	164	0.05(-0.10-0.19)	0.50	161	0.90(0.80-1.01)	0.09	0.93(0.83-1.04)	0.19
BMI (kg/m ²), women	123	0.05(-0.10-0.20)	0.53	116	0.99(0.90-1.11)	0.97	1.17(1.03-1.32)	0.01
WC (cm)	286	-0.05(-0.06-0.16)	0.39	274	0.98(0.95-1.01)	0.15	1.01(0.97-1.03)	0.69
WC (cm), men	163	0.03(-0.13-0.19)	0.69	157	0.96(0.92-1.0)	0.05	0.97(0.94-1.01)	0.19
WC (cm), women	123	0.04(-0.13-0.21)	0.61	116	0.99(0.95-1.03)	0.60	1.04(0.99-1.09)	0.07
Categorical adiposity measures								
BMI (kg/m²)								
Men vs women								
Normal weight: men/women ≤24.9 ^a	47/41	1.0		47/40	1.0		1.0	
Overweight & Obese: men ≥25.0	117	0.04(-0.11-0.18)	0.62	113	0.51(0.23-1.15)	0.11	0.60(0.26-1.40)	0.23
Overweight & Obese: women ≥25.0	82	0.10(-0.06-0.25)	0.21	76	1.73(0.64-4.62)	0.28	6.97(1.68-28.97)	0.01
WC (cm)								
Men vs women								
Small WC: men ≤102, women ≤88 ^a	85/37	1.0		85/36	1.0		1.0	
Large WC: men >102	78	0.03(-0.21- 0.35)	0.63	73	0.42(0.18-0.97)	0.04	0.32(0.14-0.75)	0.01
Large WC: women >88	86	0.07(-0.23-0.51)	0.45	80	0.79(0.30-2.08)	0.64	4.66(1.11-19.42)	0.03

Table 4. Linear and logistic regression models estimating cross-sectional associations between adiposity exposures and cerebral Small Vessel Disease outcomes at 9-year follow-up in RUN DMC. Linear regression models for WMH were adjusted for: age, education, sex, smoking, type 2 diabetes mellitus (T2DM), CSVD severity, and dementia incidence. Logistic regression models for incident microbleeds were adjusted for: age, education, sex, WMH severity, and dementia incidence; and incident lacunes for: age, education, sex, smoking, hypertension, T2DM, CSVD severity, and dementia incidence. N: number of participants. Nine participants were excluded from microbleeds analyses due to imaging artifacts (motion). WMH: White Matter Hyperintensities. ^aReferent category. ^bContinuous measures of WMH (mL), ^cpresence of any microbleeds or lacunes. CSVD: cerebral Small Vessel Disease. β: beta, standardized regression coefficient. OR: Odds Ratio. (95% CI): 95% Confidence Interval.

Discussion

To our knowledge, these data comprise a first report suggesting sex differences underlying associations of anthropometric and metabolic adiposity measures with cerebrovascular and neurodegenerative events in a clinical cohort of CSVD patients. Overweight and obesity, as well as higher leptin levels, in men with CSVD are associated with neurodegeneration, particularly in their 60s. However, among men with CSVD surviving into their 70s, higher adiposity is associated with fewer cerebrovascular markers. In contrast, women with CSVD and higher levels of adiposity in their 70s are more likely to experience lacunes and evidence no associations with neurodegeneration. These data also illustrate the nuances of following a healthier group of surviving participants with a clinically diagnosed cerebrovascular phenotype. Most observed associations occurred among men at two cross-sectional time points, at average age 66 years, and 9 years later at average age 71 years, with adverse associations with adiposity diminishing with survivorship.

Associations of higher BMI and WC with underlying cerebrovascular pathologies such as white matter hyperintensities (WMH) (338), cerebral microbleeds (339) and lacunes (340) are sparsely reported in the literature. Most data suggest higher levels of vascular risk factors, including higher BMI and central obesity or visceral adipose, to be associated with higher cerebrovascular risk (318, 341). In addition, a J-shaped BMI curve, i.e. both low and high BMI, is associated with higher risk of intracerebral haemorrhage (342). Higher WC has been associated with CSVD components, notably characterized by microbleeds, lacunes, and WMH in Korean samples (318, 339).

In addition to considering cerebrovascular components of CSVD, we also evaluated neurodegeneration. Cross-sectional associations between higher anthropometric measurements and lower TBV, GMV and HV is comparable to others reporting that higher levels of anthropometric measures are associated with a lower TBV, GMV and HV (57, 343, 344). Higher BMI has been associated with lower brain volumes, particularly GMV, in adults (345, 346), including those with metabolic syndrome (347). Higher BMI and lower GMV have also been associated with lower cognitive function (80); as has central obesity (89). Visceral adipose tissue (highly correlated with WC) measured via abdominal MRI, has been associated with lower HV in cognitively healthy elderly age 55-90 years (348). Despite the majority of studies reporting associations between higher adiposity and lower brain volumes, there are conflicting reports. For example, higher BMI was associated with higher HV in older adults with or at risk for cardiovascular disease (349).

Our findings suggest that first, higher levels of adiposity, classified as anthropometric overweight and obesity, may be related to earlier neurodegenerative outcomes, such as lower TBV and GMV, and later protection for lacunes and microbleeds in men. In contrast, among women, higher levels of adiposity are related to later occurrence of at least one marker of vascular damage, lacunes. This suggests a cerebrovascular-neurodegenerative spectrum and distinct temporality of brain outcomes associated with adipose tissue exposures that differ between men and women with CSVD. Second, these findings emphasize that adipose tissue may play endocrine, neuronal and vascular roles in brain health (77, 104, 350). These roles may be both cause and consequence of cerebrovascular and neurodegenerative pathologies associated with CSVD and/or its clinical symptoms. Third, these data underscore the profound influence of selective survival in longitudinal analyses of brain outcomes, and imply a role of brain reserve in survival, since those who died during the follow-up period had lower baseline brain volumes.

From the standpoint of metabolic adipose tissue, higher blood leptin levels and calculated leptin resistance are associated with lower TBV and GMV in older adults, particularly men with CSVD. This is somewhat novel, especially for older adults. Higher blood leptin levels have been associated with lower GMV in young adults (mean age 32.0 ± 1.0 years) (351). Among adolescents (mean age 15.3 ± 1.1 years) who gained body fat, there was observed greater GMV reduction in the putamen (352). Another study found higher leptin associated specifically with lower insula GMV among adults age 25-40 years who had a family history of obesity (353). Higher leptin levels are found in ischemic stroke patients (354). In contrast, among Japanese elderly, higher blood leptin levels were associated with higher region-specific GMV, including in the hippocampus (355). Higher leptin levels have also been shown to protect against age-related cognitive decline (355) and dementia (111) in older adults. Given that leptin signaling is dysregulated in degenerating hippocampal neurons, potentially indicating neuronal leptin resistance in Alzheimer's disease patients (356), inconsistent cross-sectional leptin associations with brain outcomes, cognition and dementias in older men may be obscured. Leptin may play a pivotal role in an aging fat-brain axis, as well as be a critical marker of aging-related changes in satiety, glucose homeostasis, inflammatory processes, vascular function, and memory formation (112, 117, 321, 357-360).

Sex differences in total and central obesity is a known phenomenon (361, 362). Differential storage of white adipose tissue is influenced by sex hormones, as well as fat depot-specific release of hormones and cytokines (363). Larger WC, a marker of visceral adipose tissue and denoting a more androgenic hormonal environment, is more common among adult men and postmenopausal women (364, 365), and

associated with higher cardiovascular disease risk (366). Sex-specific associations between anthropometric measures and brain volumes are inconsistent. As aforementioned, higher BMI is associated with lower GMV in Japanese men (367); and with temporal lobe atrophy in women aged ≥ 70 years (368). Given that the average age of menopause is 52 years in northern European populations (369), our sample most likely includes women who were going through the menopausal transition. Menopause influences the sex hormone milieu (370), as well as body composition (365), including adipose tissue and consequently, blood adipokine levels.

Differential associations between adiposity measures and brain outcomes may also be dependent on age of measurement and temporality. A changing role of adipose tissue throughout aging and in later life, by age decade, may be a real phenomenon. Our sample was examined on average, during their 7th and 8th decades of life. During this aging period, an inflection point in vascular risk factor trajectories has been observed. These trajectories are described for BMI, blood pressure and blood lipids in association with late-onset sporadic dementias (371-373). While we cannot evaluate life course trajectories in RUN DMC given a 9-year follow-up of adults age 50 years and older, their potential contribution to a changing risk profile, particularly among women, cannot be ignored. Midlife obesity has been associated with reduced GMV and greater dementia risk, whereas later life overweight and obesity has been associated with lower dementia risk (3, 343, 351, 368, 374). Given cross-sectional findings in RUN DMC, we cannot elucidate causality, nor gain insights regarding temporality of these associations.

Lack of associations between higher levels of adiposity measures and CSVD components, may be due to an attenuation of the role of vascular risk factors among those presenting with cerebrovascular disease, similar to observations in dementia epidemiology among those with prodromal dementias such as VCI (371, 375, 376). Given that aging- and dementia-related neurodegenerative events affect homeostatic regulation of energy balance, blood pressure and other aspects of metabolism, temporal paradoxical associations are often observed between vascular risk factors, such as obesity, and symptomatic, clinical outcomes (3, 77, 337, 363).

CSVD is a diagnosis that is evolving over time due to emerging phenotypic characterization and accumulating mechanistic evidence in association with clinical outcomes. CSVD is a group of pathological processes with a heterogeneous etiology and pathogenesis that involve the small arteries, arterioles, venules, and capillaries of the brain. In 2006, the baseline year of the RUN DMC, CSVD was diagnosed on the basis of the presence of any WMH and the presence of any lacunes (377), with accompanying acute symptoms (transient ischemic attacks or lacunar

syndromes), or subacute manifestations (cognitive, motor (gait) and/or mood disturbances) (378). Later recognized markers of CSVD, such as microbleeds, were also rated in RUN DMC. Therefore, our approach of evaluating adiposity measures in association with individual components of CSVD is appropriate, in addition to being more informative for intervention development.

Longitudinal analyses of risk factors for CSVD and its components are challenged since WMH, lacunes and microbleeds change over time in ways that are difficult to characterize. In RUN DMC, it was reported (324) that CSVD progression is nonlinear, accelerating over time, and is a dynamic process, with progression interrupted by reduction in some adults who, on average, show progression. RUN DMC has also reported the disappearance of lacunes and microbleeds over this 9-year follow-up period. In RUN DMC, these unexpected changes tended to occur among those presenting with a worse CSVD phenotype at baseline. Since RUN DMC participants who died during the 9-year follow-up had worse baseline CSVD and volumetric phenotypes compared to those who survived and participated in 2015, and there is worsening of CSVD on average among surviving participants, our longitudinal results are not as likely influenced by these unexpected changes.

Major strengths of our analyses are the following. First, this report is based on 9-year follow-up of men and women presenting with a distinct cerebrovascular phenotype, CSVD, with quantified neurodegenerative outcomes, notably brain volumetry, and thorough clinical and risk factor evaluation. Second, brain imaging data were analyzed by raters who were blind to clinical information, with good inter- and intra-rater agreement. Third, statistical analyses were adjusted for multiple potential confounders. Fourth, while our primary exposure was adiposity, we measured this phenotype in several ways. Using anthropometric (total and central obesity) and metabolic (leptin and adiponectin) markers of adipose tissue, differential associations were observed. This may point to different mechanistic pathways linking adipose tissue biology to neurodegeneration and cerebrovascular events. Fifth, multiple brain imaging outcomes were investigated. This is important for more precise determination of etiologic mechanisms associated with adiposity.

Despite the aforementioned strengths, several limitations should be addressed. First, serum leptin, adiponectin and lipid levels, were not measured at the follow-up examination; blood was only collected at baseline. Second, sociodemographic and health status information was not updated at follow-up in the RUN DMC. Third, nutritional intake and physical activity, known influencers of overweight and obesity, were not measured. Fourth, RUN DMC participants were all diagnosed with CSVD, which makes it difficult to generalize these results to an otherwise 'healthy' older population. It is also known that CSVD brain pathologies exist

without symptomatic burden ('silent' or preclinical CSVD) and are relatively common among elderly. Fifth, the RUN DMC sample is comprised of Dutch adults of northern European ancestry. These results require replication in heterogeneous samples due to global variations in anthropometric and brain characteristics (315, 379). Sixth, due to the small number of adults developing dementia, it could not be evaluated as a clinical endpoint. Finally, no information was available on menopausal status of women.

In conclusion, our analyses demonstrate sex differences in a potential manifestation of the fat-brain axis, and different cross-sectional and longitudinal adiposity-brain associations with aging over a 9-year period in men and women with CSVD. Future research is necessary to elucidate biological mechanisms and fat-brain associations over the life course. Elucidation of both independent and interdependent neuropathological versus vascular mechanisms underpinning the evolving aging brain is imperative.

5

A 10-YEAR FOLLOW-UP OF ADIPOSITY AND DEMENTIA IN SWEDISH ADULTS AGED 70-YEARS AND OLDER

Arnoldussen I.A.C.

Sundh V.

Bäckman K.

Kern S.

Ostling S.

Blennow K.

Zetterberg H.

Skoog I.

Kiliaan A.J.

Gustafson D.R.

Abstract

Adiposity measured in mid- or late-life, and estimated using anthropometric measures such as body mass index (BMI) and waist-to-hip ratio (WHR), or metabolic markers such as blood leptin and adiponectin levels, is associated with late-onset dementia risk. However, during later life this association may reverse and aging- and dementia-related processes may differentially affect adiposity measures. We explored associations of concurrent BMI, WHR, and blood leptin and high molecular weight adiponectin levels with dementia occurrence.

924 Swedish community-dwelling elderly without dementia, aged 70 years and older, systematically-sampled by birth day and birth year population-based in the Gothenburg city region of Sweden. The Gothenburg Birth Cohort Studies are designed for evaluating risk and protective factors for dementia. All dementias diagnosed after age 70 for 10 years were identified. Multivariable logistic regression models were used to predict dementia occurrence between 2000-2005, 2005-2010, and 2000-2010 after excluding prevalent baseline (year 2000) dementias. Baseline levels of BMI, WHR, leptin and adiponectin were used.

Within 5 years of baseline, low BMI ($<20 \text{ kg/m}^2$) was associated with higher odds of dementia compared to those in the healthy BMI category ($\geq 20\text{-}24.9 \text{ kg/m}^2$). Compared to the lowest quartile, leptin levels in the second quartile were associated with lower odds of dementia in women ($P<0.05$).

In late-life, anthropometric and metabolic adiposity measures appear to be differentially associated with dementia risk. While BMI and leptin levels are highly positively correlated, our results show that their association with dementia at age ≥ 70 years, is asynchronous. These data suggest that with aging, the complexity of the adiposity exposure may increase and suggests metabolic dysregulation. Additional studies are needed to better understand this complexity.

Introduction

In later life, lower body mass index (BMI), being underweight by definition, and declining BMI (despite one's BMI level at onset of decline), are associated with greater risk of late-onset dementias (3, 380). In mid- or adult life, higher BMI is prospectively associated with dementia risk (3). This difference in the mid-life *versus* late-life BMI-dementia risk association is often referred to as the 'obesity paradox' and evidence of reverse causality (315).

Anthropometric characteristics are sculpted by amount and distribution of adipose tissue. Changes in this important tissue result from lifetime variations in energy balance, sociocultural background, and epigenetic phenomena, and develop secondarily to central pathophysiological changes in preclinical and clinical dementia (381). Therefore, adiposity indicators like BMI, waist-hip ratio (WHR), and circulating blood levels of adipokines, such as leptin and adiponectin, are potential risk markers for dementia (88, 382-384). Since adipose tissue is the primary source of both of these adipokines, leptin and adiponectin are good biomarkers of this tissue's metabolic activity.

There is limited and ambiguous published research regarding the link between adipokine levels and sporadic, late-onset dementias (360, 385). However, leptin is associated with insulin resistance and brain health (77, 88), and is the adipose tissue hormone (111) most studied in association with brain structure and function. Leptin has numerous effects on brain development (110), and potentially on brain health in cognition and aging. Peripheral leptin enters the central nervous system and interacts with specific areas of the brain such as the hypothalamus and hippocampus (386). In addition to crossing the blood brain barrier (BBB) (387), several studies indicate that leptin is also produced in the brain, for example in the hypothalamus, cortex and cerebellum (388, 389). Leptin has been prospectively associated with dementia over 8 years (111), but not over 24 years, from mid- to late-life (368). Leptin is positively correlated with total and central anthropometric adiposity measures (390, 391).

Adiponectin is a visceral adipose tissue marker and exists as complex multimeric isoforms comprised of high molecular weight (HMW) hexamers and trimers (392). It is an effective insulin sensitizer, and circulating levels are inversely correlated with insulin resistance, metabolic syndrome, adiposity, type 2 diabetes mellitus, and cardiovascular diseases. Adiponectin modulates inflammatory responses, energy expenditure (CNS and periphery), food intake (CNS), and a number of metabolic processes, including glucose regulation and fatty acid catabolism in the periphery. Since BBB transport mechanisms for adiponectin are unclear, blood levels may not indicate potential interactions between adiponectin and the brain (393).

We investigated the association between dementia occurrence and both anthropometric and adipokine indicators of adiposity - BMI, WHR, and blood leptin and adiponectin levels - in a 10-year prospective population-based study of Swedish adults, aged 70 years and older. Moreover, we examined whether these adiposity markers predict dementia occurrence over the short- and/or long-term. This study will further reveal the complex interplay between surrogate markers of adiposity and dementia onset.

Methods

Participants

Participants comprising the Gothenburg Birth Cohort Studies, originate from two epidemiological studies in Gothenburg, Sweden: 1) the Prospective Population Study of Women (PPSW) and 2) the Gerontological and Geriatric Population Studies (H70). Both have been described in detail previously (394-396). Eligible participants were sampled from the Swedish Population Register based on birth date. PPSW participants were born in 1908, 1914, 1918, 1922 and 1930. H70 participants were born in 1930. Adults living in private households and residential care were included. There were 1725 eligible individuals in 2000-2001, and 1018 agreed to participate (response rate 59.0%). Among these, 857 (84.2%) consented to genetic analyses. Due to the nature of PPSW and H70, women (n=700) were 70-92 years and men (n=224) 70 years. Participants included 339 H70 females at age 70, 224 H70 males at age 70, and 361 PPSW females age 70 years and older (Fig. 1). Follow-up examinations were conducted in 2005-06 and 2009-10, and 689 participated in 2005-06 (response rate among survivors, 82.2%), and 518 in 2009-10 (response rate among survivors, 76.7%).

Those who died, or declined participation during follow-up, were traced in records from hospitals and homes for the aged, inpatient and outpatient departments in psychiatric hospitals and clinics, municipal psychiatric outpatient departments in Gothenburg, the hospital-linkage system, and death certificates (397). All participants (or their closest relatives) gave informed consent, which was conducted in accordance to the provisions of the Helsinki Declaration. The study was approved by the Ethics Committee for Medical Research at the University of Gothenburg.




Figure 1. Flowchart of participant characteristics and number at baseline and in follow-up examinations.

Dementia assessment

Dementia diagnoses were made according to the diagnostic and statistical manual of mental disorders 3rd edition revised (DSM-III-R) (334, 398) at examinations in 2000-2001, 2005-2006 and 2009-2010. This dementia diagnosis was made according to components of the Alzheimer's disease assessment scale-cognitive subscale (ADAS-Cog). A close informant interview was conducted using IQCode (399). Symptom domains relevant for making a dementia diagnosis include: memory (short-term, long-term), aphasia (language disturbance), apraxia (inability to carry out motor activities despite intact function), agnosia (failure to identify or recognize objects despite intact sensory function), executive function (planning, organizing, abstract thinking) and personality changes (used in DSM-III-R criteria for dementia, not in DSM-IV or DSM-V). Participants must have evidenced significant impairment in social or occupational functioning representing significant decline from a previous level of functioning. There was also a global rating of symptom status made by the interviewer (psychiatrist or psychiatric

nurse) and information from close informants related to dementia symptoms. The comprehensive psychiatric rating scale (CPRS) (400), an extensive neuropsychiatric inventory, was used to rate psychiatric symptoms and signs at face-to-face examinations. The mini-mental examination (MMSE) was used as an additional evaluation of global cognitive impairment, but is not considered diagnostic by itself. In addition, the Swedish Hospital Discharge Registry provided diagnostic information for individuals discharged from hospitals since 1978. Medical records were collected from hospitals and outpatient departments in Sweden's public health care system. Incident cases of dementia up to December 31, 2010 were also based on information from the Swedish Hospital Discharge Register (ICD-10: F00.1, F01.8, F01.9, F03.9, G30.9). Thus, dementia diagnoses were obtained for all study participants, since almost all people in Sweden receive health care from the community and have an equal chance of having a medical record. Dementia was diagnosed by neuropsychiatrists at consensus meetings based on all information sources available, e.g., neuropsychiatric examinations, medical records, and close informant interviews.

Anthropometric assessments

Anthropometric measurements were conducted in the morning when participants wore light clothing at each examination (401). Body weight was recorded to the nearest 0.1 kg, and body height was measured to the nearest 0.5 cm. BMI was calculated as kg/m^2 . Waist and hip circumferences were measured to the nearest 0.5 cm. Measures of each were conducted until there was agreement within 0.5 cm.

Blood measures

Blood samples were collected after a 12-hour fast, and plasma aliquots stored at -70°C . Regarding baseline adipokine measures, control standards and participant samples were tested in duplicate using high molecular weight (HMW) adiponectin and leptin ELISA assays (Linco Research, Inc, St. Louis, MO 63304) in the Clinical Chemistry Department at the University of Gothenburg. Blood lipids including cholesterol, high density lipoprotein (HDL), low density lipoprotein (LDL), and triglycerides, were measured at the certified Sahlgrenska Hospital Laboratory.

DNA was extracted from blood samples according to standard procedures. *APOE* (gene map locus 19q13.2) genotyping was performed by mini-sequencing as previously described in detail (402). Genotypes were obtained for the two SNPs (rs7412 and rs429358), which are used to unambiguously define $\epsilon 2$, $\epsilon 3$, and $\epsilon 4$ alleles.

Lifestyle assessments and medical history

Level of education (completing six years compulsory education or less *vs* at least

compulsory education); socioeconomic status (SES, working vs middle/upper class); alcohol consumption and cigarette smoking (ever vs never use); medication use; and medical history were queried at each examination. Diagnoses of myocardial infarction, stroke, cancer, and diabetes were self-reported as well as determined by clinical examinations (ECG and blood samples), case records, hospital discharge registry, national cancer registry, and national stroke registry. Systolic (SBP) and diastolic blood pressures (DBP) were measured while participants were sitting down and at rest for at least 5 minutes. Depressive symptoms were measured using the Montgomery–Åsberg depression rating scale (MADRS) (403).

Statistical analyses

Means and standard deviations were calculated for all continuous variables, and frequencies and percentages for categorical variables. BMI, WHR, leptin and adiponectin were used in analyses as continuous, and as traditional quartiled categorical variables. We considered baseline adiposity exposures at 2000 and 2005, and percent BMI change from 2000-2005 in association with dementia occurring between 2005 and 2010. Clinically relevant BMI change was categorized as $\pm > 5\%$, and $\pm > 10\%$. We created combined anthropometric and adipokine categories as potentially better surrogates of adiposity-associated risk (404, 405). We cross-tabulated leptin quartiles by the four traditional BMI (underweight, $< 20.00 \text{ kg/m}^2$; healthy, $20.00 - 24.99 \text{ kg/m}^2$; overweight, $25.00 - 29.99 \text{ kg/m}^2$; and obese $\geq 30.00 \text{ kg/m}^2$) categories. In other words, those with a low leptin and low BMI were compared to those with a high leptin and high BMI, versus either alone. Similarly, leptin quartiles were cross-tabulated by WHR (central obesity: men $> 0.90 \text{ cm}$, women > 0.85).

Multivariable logistic regression analyses were used to calculate odds ratios (OR) for developing dementia over different time periods. First, regression models estimated the odds of dementia over ten years (2000-2010) given adiposity exposures measured at baseline in 2000. Second, we estimated the association between baseline adiposity indicators measured in 2000 among those without dementia and subsequent dementia occurring during two five year periods, 2000-2005 and 2005-2010. To reiterate, prevalent dementia cases at baseline were excluded from all models. In models predicting dementia by adipokines, we adjusted for BMI if the correlation between adipokine and BMI did not exceed $r \geq 0.70$.

Selection of covariates originated from a pool of potentially biologically relevant variables: age; sex; education; socioeconomic status (SES); smoking; alcohol intake; depressive symptoms; diabetes; cancer; heart attack; stroke/transient ischemic attack; hypertension; blood triglyceride, cholesterol, HDL, LDL, and glucose levels; DBP; SBP; and APOE ϵ 4 genotype. All potential covariates were included in age-adjusted

logistic regression models predicting dementia over the entire follow-up period. Age, sex, SES, smoking status (ever/never), diabetes (yes/no), alcohol intake (ever/never), depressive symptoms (MADRS score > 12), and presence of at least one APOE ϵ 4 allele were significant at $P \leq 0.05$ and included in multivariable models. Two-tailed tests were used with a significance level of $P \leq 0.05$. SPSS, version 20.0 (IBM Corporation, Armonk, New York, USA), was used to perform data analyses.

Results

Overall, 924 adults (224 men and 700 women) without dementia, participated in the baseline (year 2000) examination. PPSW and H70 were representative of the population base with regard to sex; marital status; income; community rent allowance for those who could not afford housing; rate of inpatient and outpatient care in psychiatric hospitals, clinics, and municipal outpatient departments; and rates of registration with the Temperance Board (national registry for alcohol abuse). Over ten years, 134 participants developed dementia.

The average age of the participants at baseline ($n=924$) was 74.0 years, and when stratified by sex, the average age of men was 70.0 years and for women, 75.3 years (Table 1). Compared to men, women were more likely to be underweight and had higher plasma levels of leptin and adiponectin. Approximately 25% of the study population had at least one APOE ϵ 4 allele (27.7% in men and 24.4% in women).

Multivariable logistic regression models were used to examine the relationship between continuous baseline adiposity indicators and subsequent dementias. No associations were found between BMI, WHR, leptin or adiponectin and dementia occurrence in the entire sample over the ten year follow up (Table 2; data not shown for adipokines).

To better understand the temporality of association between adiposity markers and dementia, we divided the ten year follow-up period into two five year dementia occurrence intervals - 2000-2005 and 2005-2010 (Table 2), as has been done previously (406). When analyzed continuously, a higher baseline BMI lowered the odds of dementia within five years. However, using traditional BMI categories, participants with an 'underweight' BMI ($< 20 \text{ kg/m}^2$) had higher odds of developing dementia compared to participants with a 'healthy' BMI (reference category BMI 20.00-24.9 kg/m^2) (Table 2). This finding among underweight may have driven the protective association observed when BMI was modelled as a continuous variable. This association was driven by female participants (Table 2).

Variable	All (n=924)	Men (n=224)	Women(n=700)
Age, years, mean \pm SD	74.0 \pm 5.3	70.0 \pm 0.0	75.3 \pm 5.5
Education <8 years, N (%)	544 (58.9)	125 (55.8)	419 (59.9)
Socioeconomic status (SES)			
working class, N (%)	351 (38.0)	57 (25.4)	294 (42.0)
middle or upper class, N (%)	300 (32.5)	100 (44.6)	200 (28.6)
Waist-hip ratio (WHR), mean \pm SD	0.87 \pm 0.083	0.95 \pm 0.06	0.84 \pm 0.07
Low WHR: men \leq 0.90, women \leq 0.85, N (%)	375 (40.6)	33 (14.7)	342 (48.9)
High WHR: men > 0.90, women > 0.85, N (%)	509 (55.1)	186 (83.0)	323 (46.1)
BMI, kg/m ² , mean \pm SD	26.7 \pm 4.2	29.9 \pm 3.9	26.6 \pm 4.3
\leq 20.00, N (%)	28 (3.0)	4 (1.8)	24 (3.4)
20.01 - 24.99, N (%)	290 (31.4)	66 (29.5)	224 (32.0)
25.00 - 29.99, N (%)	375 (40.6)	108 (48.2)	267 (38.1)
\geq 30.00, N (%)	180 (19.5)	45 (20.1)	135 (19.3)
Leptin (ng/mL), mean \pm SD	31.6 \pm 24.0	15.07 \pm 10.5	37.28 \pm 24.7
Q1 (ng/mL)	\leq 13.36	\leq 8.06	\leq 18.65
Q2 (ng/mL)	13.37 - 25.24	8.07 - 12.04	18.66 - 30.52
Q3 (ng/mL)	25.25 - 44.00	12.05 - 21.07	30.53 - 50.83
Q4 (ng/mL)	>44.00	>21.07	>50.83
Adiponectin (ng/mL), mean \pm SD	6.10 \pm 4.54	3.52 \pm 2.34	6.97 \pm 4.77
Q1 (ng/L)	\leq 3.11	\leq 1.97	\leq 3.76
Q2 (ng/L)	3.12 - 4.89	1.98 - 3.01	3.77 - 5.71
Q3 (ng/L)	4.90 - 7.84	3.02 - 4.57	5.72 - 8.90
Q4 (ng/L)	>7.84	>4.57	>8.90
Alcohol consumption			
g per week, mean \pm SD	37.5 \pm 66.9	72.2 \pm 101.9	25.63 \pm 44.4
Smoking, ever, N (%)	353 (38.2)	151 (67.4)	202 (28.9)
Serum levels (mmol/L)			
Triglycerides, mean \pm SD	1.4 \pm 0.7	1.41 \pm 0.6	1.46 \pm 0.8
Cholesterol, mean \pm SD	5.9 \pm 1.1	5.5 \pm 0.9	6.1 \pm 1.0
High density lipoprotein (HDL), mean \pm SD	1.6 \pm 0.4	1.3 \pm 0.4	1.7 \pm 0.4
Low density lipoprotein (LDL), mean \pm SD	3.7 \pm 0.9	3.6 \pm 0.9	3.8 \pm 0.9
Systolic blood pressure, mean \pm SD	157.7 \pm 22.5	155.1 \pm 23.1	158.2 \pm 22.3
Diabetes mellitus, N (%)	111 (12.0)	41 (18.2)	70 (9.9)
Depression,* N (%)	99 (10.7)	15 (6.7)	84 (12.0)
Presence of any Apoe ϵ 4 allele,† N (%)	233 (25.2)	62 (27.7)	171 (24.4)

Table 1. Baseline characteristics of participants without dementia in the Gothenburg Birth Cohort Studies. *Clinically relevant depressive symptom burden, MADRS score > 12. †APOE ϵ 4 allele: apolipoprotein epsilon 4 allele.

In women, we also found an association between leptin levels and odds of dementia within the first five years of follow up (2000-2005, Table 3). No associations were observed in men. Baseline leptin levels in the second quartile (18.66 - 30.52 ng/mL) were associated with lower odds of developing dementia compared to leptin levels in the first quartile (≤ 18.65 ng/mL, reference). No associations were found between serum levels of HMW adiponectin and dementia occurrence.

Considering only dementias occurring during 2005-2010, there were no associations with baseline BMI, WHR, leptin or adiponectin (Table 2). In addition, BMI change from 2000-2005 was not associated with dementias occurring from 2005-2010 (data not shown). Combinations of anthropometric BMI or WHR risk categories by leptin quartiles did not shed any additional light on these associations (data not shown).

	2000-2010			2000-2005			2005-2010		
	Total	Men	Women	Total	Men	Women	Total	Men	Women
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Baseline BMI (kg/m²)									
Continuous	0.94 (0.86-1.02)	0.99 (0.85-1.10)	0.92 (0.83-1.01)	0.88 (0.79-0.97)*	0.89 (0.67-1.18)	0.87 (0.78-0.97)**	0.99 (0.89-1.11)	0.99 (0.83-1.21)	0.98 (0.86-1.12)
<20.0	2.94 (0.73-11.88)	NE†	4.59 (0.95-22.22)	4.21 (1.03-17.21)*	NE	5.34 (1.19-23.87)	1.12 (0.13-9.41)	NE	1.32 (0.14-11.45)
20.00 - 24.99 †	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
25.00 - 29.99	0.79 (0.39-1.56)	1.34 (0.34-5.24)	0.67 (0.29-1.50)	0.13 (0.25-1.19)	0.39 (0.04-3.37)	0.54 (0.24-1.27)	0.69 (0.28-1.74)	1.91 (0.33-11.26)	0.45 (0.14-1.43)
≥30.00	0.93 (0.39-2.22)	0.80 (0.13-5.10)	0.92 (0.34-2.51)	0.25 (0.19-1.54)	0.51 (0.04-6.94)	0.51 (0.16-1.66)	1.21 (0.41-3.57)	0.76 (0.06-9.29)	1.29 (0.38-4.40)
Baseline WHR (cm)									
Continuous	1.07 (0.69-1.65)	2.95 (0.92-9.42)	0.87 (0.54-1.39)	0.95 (0.59-1.51)	0.80 (0.17-3.80)	0.96 (0.59-1.57)	1.20 (0.65-2.24)	4.75 (1.03-21.78)	0.81 (0.42-1.55)
Low WHR†	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
High WHR	0.80 (0.42-1.52)	3.13 (0.37-26.52)	0.62 (0.30-1.28)	0.63 (0.32-1.27)	0.54 (0.05-5.45)	0.63 (0.31-1.32)	1.11 (0.47-2.65)	NE	0.77 (0.29-2.06)

Table 2. Odds of developing dementia from 2000-2010, 2000-2005, and 2005-2010 by baseline BMI and WHR among participants without dementia. The Gothenburg Birth Cohort Studies. Multivariable models included the following covariates: in 2000-2010, age, sex, socio-economic status (SES), ApoEε4 allele, and smoking; in 2000-2005, age, sex, depressive symptoms and SES; in 2005-2010, age, sex and smoking. †Reference category. Multivariable models were assessed for the complete cohort (total), and subsequent for both men and women in the cohort. Low WHR: men ≤ 0.90 cm, women ≤ 0.85 ; High WHR: men > 0.90 cm, women > 0.85 . Number of incident dementias in 2000-2010, N=134 (men N=18 and women N=116); in 2000-2005, N=68 (men N=8 and women N=60); and in 2005-2010, N=66 (men N=10 and women N=56). * NE: not estimable due to small number. *P<0.05; ** P<0.001.

2000-2010			2000-2005			2005-2010					
	Total	Men	Women	Total	Men	Women	Total	Men	Women		
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)		
Baseline leptin (ng/mL)											
Continuous	0.55 (0.23-1.30)	0.96 (0.18-5.16)	0.48 (0.17-1.31)	0.33 (0.11-0.99)*	0.42 (0.04-4.33)	0.36 (0.1-1.0)*	0.82 (0.23-2.89)	1.76 (0.17-18.70)	0.56 (0.12-2.56)		
Quartiles											
	Men	Women									
Q1†	≤8.06	≤18.64	1.0	1.0	1.0	1.0	1.0	1.0	1.0		
Q2	8.07-12.04	18.65-30.52	1.15 (0.51-2.63)	0.30 (0.06-1.55)	1.87 (0.69-5.10)	2.62 (0.73-9.42)	1.91 (0.21-23.52)	0.32 (0.10-0.97)*	0.65 (0.22-1.89)	0.35 (0.05-2.65)	0.92 (0.25-3.36)
Q3	12.05-21.07	30.53-50.83	1.17(0.51-2.71)	0.56 (0.11-2.67)	0.63 (0.59-4.52)	3.47 (1.0-12.02)	1.65 (0.12-22.41)	0.82 (0.32-2.04)	0.37 (0.10-1.31)	0.29 (0.03-3.13)	0.35 (0.07-1.73)
Q4	>21.07	>50.83	0.33 (0.07-1.59)	0.44 (0.08-2.57)	0.79 (0.26-2.47)	0.66 (0.11-3.83)	NE	0.45 (0.13-1.13)	0.39 (0.08-1.99)	0.69 (0.10-4.58)	0.62 (0.25-2.18)
Baseline adiponectin (ng/mL)											
Continuous	1.14 (0.38-3.40)	0.49 (0.07-3.45)	1.63 (0.42-6.33)	1.63 (0.32-8.31)	1.18 (0.06-25.13)	1.77 (0.26-12.22)	0.89 (0.22-3.54)	0.30 (0.03-3.29)	1.53 (0.25-9.30)		
Quartiles											
	Men	Women									
Q1†	≤1.97	≤3.76	1.0	1.0	1.0	1.0	1.0	1.0	1.0		
Q2	1.98-3.01	3.78-5.71	0.39 (0.15-1.51)	0.19 (0.02-1.74)	0.49 (0.16-1.48)	0.43 (0.10-1.82)	1.22 (0.07-22.11)	0.29 (0.05-1.65)	0.36 (0.10-1.24)	NE	0.67 (0.16-2.73)
Q3	3.02-4.57	5.73-8.90	0.74 (0.31-1.07)	0.89 (0.21-3.83)	0.62 (0.22-1.77)	1.06 (0.33-3.43)	2.52 (0.19-34.24)	0.81 (0.21-3.13)	0.53 (0.18-1.61)	0.53 (0.09-2.91)	0.44 (0.09-2.03)
Q4	>4.57	>8.90	0.93 (0.42-2.08)	0.38 (0.06-2.27)	1.19 (0.47-3.03)	1.19 (0.38-3.77)	1.19 (0.07-21.63)	1.08 (0.31-3.76)	0.79 (0.28-2.21)	0.22 (0.02-2.23)	1.29 (0.36-4.56)

Table 3. Odds of developing dementia from 2000-2010, 2000-2005, and 2005-2010 by baseline leptin and adiponectin among participants without dementia. The Gothenburg Birth Cohort Studies. Multivariable models include in 2000-2010, age, sex, social economic status (SES), ApoEε4 allele, and smoking; in 2000-2005, age, sex, depressive symptoms and SES; in 2005-2010, age, sex and smoking. †Reference category. Multivariable models were assessed for the complete cohort (total), and subsequent for men and women. In adiponectin analyses BMI was included as covariate. NE: not estimable due to small number. Number of incident dementias in: 2000-2010, N=134 (men N=18 and women N=116), 2000-2005 N=68 (men N=8 and women N=60).

Discussion

In an elderly Swedish sample of women and men, we observed that a low BMI (<20 kg/m²) was associated with over a 5-fold higher odds of developing dementia within five years in women, aged 70 years and older. Given published literature on mid-life obesity and later onset dementia, this may be evidence of an obesity paradox and/or reverse causality. However, without mid-life measures, we cannot be sure. In addition, the odds of developing dementia were 70% lower in women aged 70 years and older who had intermediate serum leptin levels based on analysis of leptin quartiles. Thus, despite a robust correlation of $r = 0.71$ between BMI and leptin in this sample, there is a suggestion that these adiposity indicators were differentially associated with development of dementia within 5 years of measurement in elderly women.

Other studies in middle-aged and elderly women have suggested contradictory associations between anthropometric versus leptin levels and cognition or dementia (360, 407, 408).

Our analyses support a first-ever published report from the original Gothenburg Birth Cohort Study (H70) among those born 1901/02 (406). Similar to the 2003 report, there was no association between higher BMI and dementia observed during the first 10 years of follow-up. In contrast to the 2003 report, the current analysis includes primarily those born 1930. Thus, with further follow-up, we will be able to observe a 30 year cohort comparison of the BMI-dementia association. In addition, we also chose to focus on all dementias, due to controversies and changes in the field regarding diagnoses of dementia subtypes without the use of neuroimaging or cerebrospinal fluid biomarkers (409).

Our study has many advantages. First, the population sample is part of a long-standing series of rich longitudinal birth cohort studies in Sweden beginning in 1968 (PPSW) and 1971 (the first H70 cohort) (394, 395), with consistent measures over time among unique birth cohorts. Second, there is long follow-up of a large sample of rigorously studied elderly, age 70 years and older, and adjustment for multiple potential confounders. Third, we measured blood levels of adipokines reflecting the metabolic activity of adipose tissue. Finally, we used the study design to our advantage, to estimate temporal (shorter versus longer term) associations of adiposity indicators with dementia. Given the important role of temporality in understanding risk and protective factor-dementia associations, we see this as a definite strength of these studies.

As with any study design, there are limitations. First, while data from the Gothenburg Birth Cohort Studies have been published using this merged sample examined with identical methods by the same research group at the same time (410, 411), the study is unbalanced regarding sex and age. All men were aged 70 years at baseline, whereas women were aged 70 years and older. Thus, our observations regarding sex differences should be interpreted cautiously. Second, a small number of male participants reduced the power for analyzing associations in men given that dementia onset was monitored during a period of relatively low risk, notably the 8th decade of life (406). Third, leptin and adiponectin measures were only measured at baseline, thus repeated measures and change in these important hormones over time cannot be considered. Fourth, other nutritional influencers of BMI and WHR, including estimates of energy intake and expenditure, are unavailable. Fifth, our leptin findings are rather conservative. While leptin has strong mechanistic actions in the brain, the use of this marker in the periphery as a reflection of the fat-brain axis remains unclear, especially at a life stage when there are ageing-related body

composition changes. In addition, replication of our observations regarding leptin is essential. Finally, some may deem four and five year examination intervals to characterize date of dementia onset as unacceptable. However, given the insidious and often slowly progressive course of dementia onset and a population-based sample, we deem it highly acceptable for these types of analyses. Calculating odds ratios (OR), given this nuance, closely approximates the risk. We have shown this in a previous publication on adiposity measures and dementia (406).

While we found that the odds of developing dementia are higher in elderly women with low BMI within five years, this is not a new finding. The association between BMI and dementia appears to change over the life course. Some studies show that adults with a high midlife BMI or central obesity have a higher risk of late-life dementia (3). Others have reported that weight loss or BMI decline is associated with dementia risk in elderly (71, 73). Our data did not suggest associations of baseline BMI and dementias occurring between 2000-2010 nor dementias occurring between 2005 and 2010. Nor was BMI change from 2000-2005 related to dementias occurring between 2005-2010. This suggests temporal issues regarding the BMI-dementia association in this sample, however we cannot evaluate this historically since we do not have measures of BMI prior to 70 years. Thus, a lower BMI may be a preclinical marker for underlying dementia development within five years of a clinical diagnosis in women aged 70 years and older. These results should be interpreted with caution due to a small number of women in the lowest BMI category. The association of lower baseline BMI with dementia may result from a lengthier period of weight loss accompanying prodromal dementia prior to age 70 years, supporting BMI as a possible preclinical marker of dementia.

That low BMI may be a preclinical marker for dementia, is suggested by neuropathophysiological changes accompanying dementia. These changes can induce changes in body weight and adipose storage. For example, brain areas that control weight (i.e. mesial temporal cortex) are affected during the preclinical dementia phase; and adipose tissue loss may result from preclinical apathy, reduced olfactory function, difficulty with eating (e.g., aphasia), inadequate nutrition or prescribed medicines targeting dementia related symptoms e.g. depression or cognitive impairments (337).

Furthermore, we found that the odds of developing dementia in elderly women were lower among those with intermediate blood leptin levels. Postmenopausal women may have a greater risk of developing dementia than men, perhaps due to changes in leptin and brain function due to a fall in endogenous estrogen levels following menopause (412). Estrogen and leptin have neuroprotective effects on cognition by regulating neurogenesis, hippocampal synaptic plasticity, and axonal

growth (77). However, the roles of adipokines are diverse, and often enigmatic. The adiposity-dementia association, remains to be elucidated, as this is by no means clear.

As higher levels of adiposity, commonly measured as anthropometric overweight and obesity, are associated with higher risk for cardiovascular disease (CVD) (85, 87), and CVD is a risk factor for dementia (67), we explored CVD variables as potential covariates. None were significantly associated with dementia in age-adjusted models. This null observation associating dementia with CVD risk in late-life concurs with other published reports (65). In addition, adiponectin, known to be associated with CVD, was not associated with dementia.

The underlying neurodegenerative and vascular pathologies observed in dementia, and subsequent impairment of key fat-brain and gut-brain feedback loops (413), may be at the root of conflicting observations; and seemingly intertwined adiposity factors may be differentially associated with dementia when measured within five years. Our findings are supported by studies reporting that higher levels of leptin are prospectively associated with lower odds of dementia or mild-cognitive impairment in elderly (360, 414, 415). Metabolic dysregulation may accompany or promote late-life alternations in leptin levels and dementias (416). Atrophy of the posterior hypothalamus affects appetite and feeding behavior via disintegration of network connections and effects on hormone synthesis (416, 417). In addition, higher levels of leptin are observed to be neuroprotective (418).

Conclusion

Two highly correlated adiposity variables measured in late life do not associate similarly with late-onset dementia in women. Our analyses show a higher odds of dementia with a low BMI, and a lower odds of dementia with intermediate leptin levels when dementia occurs within 5 years of the adiposity measurement among women aged 70 years and older. Lower BMI and higher leptin may potentially characterize aspects of metabolic dysregulation associated with presence versus absence of prodromal dementia. However, both associations suggest a similar adiposity phenomenon, whether cast as lower BMI being risky or higher leptin levels being protective.

6

GENERAL DISCUSSION AND CONCLUDING REMARKS

General discussion

In 2015, nearly 40% of the adult population in the United States was obese, and the obesity epidemic is still rising (419). Obesity is characterized by accumulation of white adipose tissue (WAT; adiposity), which eventually may have adverse effects on human health and cognition. Accumulation of WAT is potentially modifiable, therefore it is important to identify and better understand affected mechanisms in the adiposity-brain-axis. Eventually, knowledge on the adiposity-brain-axis may facilitate interventions to reduce the impact on public health. Therefore, with the research performed in this thesis we aimed to reveal mechanisms underlying the adiposity-brain-axis, in particular the involvement of cerebrovasculature, adipokines and aging, and the effects of fatty acids in these mechanisms in both mice and men. In this General discussion, it will be considered to what extent the results of the experiments described in the previous chapters may elucidate the adiposity-brain-axis.

Mouse models

Transgenic mice models are commonly used in preclinical obesity research, because in a relatively short period of time diet-induced-adiposity can be provoked, and/or the susceptibility to develop co-morbidities like cardiovascular disorders. In our preclinical studies (**Chapter 2 and 3**), we examined the ApoE3*Leiden and low-density lipoprotein receptor knock-out Leiden (LDLr^{-/-}) mouse models. Both mouse models are susceptible for accumulation of WAT, and the development of cardiovascular disorders like atherosclerosis and hypertension. We elucidated cerebral blood flow (CBF), structural brain changes and functional connectivity in these mouse models using high resolution (11.7 Tesla) magnetic resonance imaging (MRI)-scanner. Thereby, we were, to our knowledge, one of the first examining these parameters at this high resolution in context of rodent obesity. Nevertheless, there are some limitations concerning mouse models. For instance, anesthesia is a limitation within MRI research in rodent models. Anesthesia is stressful, induces hyperglycemia, and some types of anesthesia are stored within WAT (420).

Secondly, metabolic phenotypes vary between mouse strains, and even vary between subpopulations of the same strain. For instance, C57BL/6J (The Jackson Laboratory) mice are predisposed to the obese and diabetic phenotypes, and show impaired insulin secretion compared to C57BL/6N (NIH) mice (421). Moreover, transgenic mouse models are characterized by pathological features. For instance, the LDLr^{-/-} mouse model is susceptible to develop adiposity, hyperlipidemia, and atherosclerosis (**Chapter 3**). In detail, the loss of hepatic low-density lipoprotein receptor (LDLr) expression can increase circulating lipoproteins providing

increased triglyceride substrate for uptake by adipocytes (422). Alternatively, loss of the LDLr may have indirect effects on reducing lipolysis in adipocytes or altering thermogenesis, which would also increase lipid storage in WAT (422). The exact biological mechanisms involving lipid intake and accumulation in specific adipose depots is still under debate in the LDLr^{-/-} mouse model, which might complicate the comparison of findings in this mouse model compared to other mouse models. Therefore, it is important to recognize the details of genetic background when interpreting experimental results, and the affected biological mechanisms involved in transgenic mouse models.

Thirdly, it remains challenging to compare mice to humans, particularly aging effects as for instance mice exert a maturation phase of 12 weeks which includes three weeks of weaning. Thus compared with humans, mice have an accelerated maturation rate during childhood because they develop rapidly during infancy (150 times faster than humans), and generally, at 3–7 months are considered mature adults, equivalent to 20–30 years in human (421, 423). These limitations of mouse models must be taken into account when designing an experiment. Some possible directions to overcome these limitations could be elucidating the exact biological processes affected in transgenic mouse models. And last, the use of a standardized method such as proposed by the ARRIVE guidelines (286). Nevertheless, animal models still have the potential and are indispensable for examining biological mechanisms in anatomical, metabolic, and neurochemical detail.

Cerebral circulation

The influence of adiposity on the vasculature may play a key role in the development of cognitive impairment, and dementia (424, 425). In detail, adiposity can induce atherosclerosis in large cerebral arteries, and more importantly it can impair the cerebral microcirculation (425). Alterations in cerebral circulation can induce a decreased cerebral blood flow (CBF; hypoperfusion) or cerebral small vessel disease (CSVD) (318, 319, 339). CSVD can be diagnosed by markers like white matter hyperintensities (WMH), lacunes and microbleeds using MRI imaging. Biological mechanisms involved are inflammation, loss of pericyte coverage on cerebral microvessels, blood brain barrier disruption, altered hippocampal angiogenic gene expression (282, 317). Alterations within the cerebral circulation can contribute to cognitive impairment as CBF is essential for normal neuronal function (319). Obese individuals often have an increased systolic blood pressure, and reduced CBF (18, 426). In **chapter 3** we reinforce these finding in obese mice. Nevertheless, in a model for mild obesity we found reverse effects in CBF (**Chapter 2**). Plausibly, the blood vessel walls in the relatively mild obese ApoE3*Leiden mouse model were not severely affected or atherosclerotic, and may still be able to compensate the

HFD induced changes. Moreover, we examined adiposity related to brain volumes in **chapter 4**, where we reported that body mass index (BMI), waist circumference (WC), and leptin were negatively associated with total gray matter volume (GMV). Moreover, we found that leptin was negatively associated with severe white matter hyperintensities (WMH) and presence of lacunes, which indicated a protective effect of leptin in relation to CSVD outcomes. Whereas others reported that adiposity indicated by the amount of visceral adipose tissue, is positively associated with CSVD indicated by microbleeds, lacunes, and WMH (318, 339). Possibly, anthropometric measurements like BMI and WC may not be accurate enough to indicate specifically visceral adipose tissue. In addition, CSVD progression seems to be nonlinear, accelerating over time, and seems to be a highly dynamic process (324), which might complicate the over time analyses of the association between adiposity and CSVD markers. Moreover, in humans we used 1.5 Tesla MRI images to indicate CSVD markers. High resolution images, used in our preclinical research (11.7 Tesla), might provide the opportunity to precisely examine microstructural changes within the cerebral circulation. Thus, in rodent models we reinforce the adverse effects of obesity within cerebrovascular pathology. However, in humans more research is necessary to elucidate the effect of obesity on cerebrovascular pathology, in specific brain regions and its progression.

White adipose tissue (WAT)

Adiposity seems to be a key inducer of adverse effects on health, and cause of dysfunction of numerous types of tissues including the brain (112, 117, 321). WAT is a large endocrine organ, secreting hundreds of adipokines, and particularly the endocrine activity of WAT can be deregulated in adiposity (77). Adipokines contribute to the regulation of appetite and satiety, fat distribution, insulin secretion and sensitivity, energy expenditure, endothelial function, inflammation, and blood pressure (427). Therefore, alterations in adipokine secretion may link adiposity to its inflammatory, metabolic, and vascular comorbidities (427). Functional characterization, in particular of the more recently identified adipokines like vaspin, apelin and resistin, is a major task in future adipokine research.

In mouse models, we found that leptin is not increased in mildly obese mice, whereas leptin levels were significantly increased in the obese mice (428)(**Chapter 2 and 3**). In **Chapter 4 and 5**, we reported that obese individuals have increased leptin levels, and women exert a higher level than men. These findings are in coherence with findings in literature (77, 389). Moreover, we reported asynchronous results for leptin in **chapter 4 and 5**, for instance leptin was negatively associated with GMV in men, whereas it was suggested to be protective in relation to CSVD outcomes in men, and dementia in women. These findings might indicate that leptin's effects in

the adiposity-brain-axis are diverse, change with aging and are influenced by sex. Notably, leptin levels are mostly determined in blood plasma, which provides an accurate indication of metabolic activity. However, leptin signaling within the brain depends on transport of leptin across the blood brain barrier, and availability of its receptors in specific brain regions (77). Therefore, further research is necessary to elucidate the exact function of leptin in specific brain regions, aging and sex.

Adiponectin is a protein hormone, most well-described for modulating inflammatory responses, energy expenditure, energy intake and a number of metabolic processes, including glucose regulation and fatty acid catabolism in the periphery (136-138, 140, 429). In the periphery, adiponectin is released from adipose tissue into the blood circulation as full-length trimers, hexamers, high molecular weight, multimers, and a globular fraction called globular adiponectin (430, 431). In humans, plasma adiponectin levels are, in contrast to most adipokines, inversely correlated to an increase in WAT, as well as to surrogate measures such as percentage body fat, waist-to-hip ratio and BMI (432, 433). In **chapter 4** we confirm these findings, nevertheless we found no association between adiponectin and brain volumes, CSVD markers or dementia. In addition, leptin and adiponectin exert insulin-sensitizing effects, they are considered as important regulators of β -cell mass and survival (77, 427). Therefore, both adipokines might influence the development insulin resistance, and eventually type 2 diabetes mellitus. The mouse studies described in **chapter 2 and 3** provide blood glucose and plasma insulin levels, however both studies are limited in examining glucose homeostasis. Following studies should investigate the functionality of α - and β -cells, glucagon, or assess a glucose tolerance test.

Aging

During aging the associations clarifying the adiposity-brain-axis can be rather ambiguous. In detail, midlife obesity is associated with reduced GMV and greater dementia risk, possibly via impairment of cerebral circulation and/or neuroinflammation (3, 343, 351, 368, 374). High adiposity might contribute to neurodegenerative and cerebrovascular changes through vascular and metabolic pathways (434). However, others reported that in humans that midlife overweight and obesity were not associated with infarct-like brain lesions, cerebral microbleeds, or dementia compared with normal weight after a mean follow-up of 26.2 years (435). In **chapter 4**, we found that BMI, waist circumference (WC) and leptin are negatively associated with GMV in adult men (> 50.0 years). Nevertheless, if obesity is included in forecast models prevalence of dementia is estimated to be 9% for the United States, 14 % in Australia, and 19% higher in China, in 2050 (436, 437). In mid-adult obese rodents, a butyrate supplementation served as preventative for

obesity and obesity related co-morbidities (**Chapter 3**). We indicated in this rodent study (**Chapter 3**) that particularly in midlife inflammatory and cerebrovascular mechanisms can be influenced by for instance a butyrate intervention. These findings might indicate that midlife obesity can be a susceptible life period in which a dietary intervention can beneficially effect obesity and its co-morbidities.

Additionally, examining aging in the adiposity-brain-axis is complicated by sex hormones, which induce sex-specific features in WAT storage. For instance, importantly, women differ with respect to distribution of WAT, men tend to accrue more WAT within the visceral adipose depots, which has been highly correlated to increased cardiovascular risk (438). Women on the other hand, accrue more WAT in the subcutaneous depots prior to menopause, a feature associated with protection from the negative consequences associated with obesity and the metabolic syndrome (438). In detail, after menopause, deposition of adipose tissue shifts to favor the visceral adipose depot. This shift is accompanied by a parallel increase in metabolic risk reminiscent to that seen in men (438). In this thesis we found sex differences in associations between adiposity and brain volumes mainly in adult men (> 50.0 years), and between adiposity and dementia particularly in older women (>70.0 years) (**Chapter 4 and 5**). Postmenopausal women may have a greater risk of developing dementia than men, perhaps due to changes in leptin and brain function due to a fall in endogenous estrogen levels following menopause (412). Estrogen seems to protect against adiposity, and it can influence the function of adipokines like leptin and adiponectin (439). Moreover, estrogen and leptin can have neuroprotective effects on cognition by regulating neurogenesis, hippocampal synaptic plasticity, and axonal growth (77). Future studies should elucidate the role of sex-hormones in an aging adiposity-brain-axis.

Weight loss in late life is strongly associated with dementia risk. Underweight in elderly is strongly associated with increased dementia risk within a 5 year follow up (315, 440) (**Chapter 5**), which might be associated with the presence versus absence of prodromal dementia. The biological mechanisms involved in late life weight changes and the risk of dementia are not fully understood but might be associated with pathophysiological changes in brain areas regulating energy intake, behavioral disturbances, and medication (71, 337, 381). Moreover, we found that leptin can be differently associated with dementia odds than BMI in elderly (**Chapter 5**). This might indicate that the role of leptin changes during aging. Our findings are supported by studies reporting that higher levels of leptin are prospectively associated with lower odds of dementia or mild-cognitive impairment in elderly (360, 414, 415). Moreover, plasma leptin levels have been positively correlated with GMV in various brain regions including the hippocampus, and high leptin levels might protect against age-related cognitive decline in an elderly population study

(355). A study of *Bonda et al.* revealed that leptin signaling is deregulated in the degenerating hippocampal neurons in patients with Alzheimer's disease (AD), which may indicate neuronal leptin resistance in AD (356). Metabolic dysregulation may accompany or promote late-life alternations in leptin levels and dementia (416). Thereby, leptin might play a pivot role in an aging adiposity-brain-axis, and possibly by affecting satiety, glucose homeostasis, inflammatory processes, vascular function, and memory formation (112, 117, 321, 357-360). A limitation in human studies is that the interpretation and comparing epidemiological findings is rather complicated as a study population of people born in 1930 exert different anthropometric features as individuals born in 1990, plausibly due to changes dietary habits, socioeconomic environment, and scientific and technological progress.

Thus, obesity in midlife most likely forecasts a higher dementia risk in late life, although in elderly weight loss might moreover indicate presence of prodromal dementia. Future studies could elucidate the progression of structural brain changes affected by adiposity in aging, by plotting for example GMV against an obese and lean population over a life period.

Diets

High fat diets (HFDs) are regularly used to elicit changes in body composition like provoking adiposity, inflammation, non-alcoholic fatty liver disease (NALFD), high level of circulating triglycerides and cholesterol, and cognitive deficits (441, 442). We confirm these effects of a HFD in **chapter 2 and 3**. The duration of HFD feeding is an important factor to consider concerning aim of the study and mouse model. As for instance, in a LDLr^{-/-} mouse model, overweight develops after 6-10 weeks on a 24% w/w HFD, and the first structural brain changes are observed after about 12 weeks on HFD and it requires about 24-32 weeks to develop cognitive impairment. Male mice are generally more sensitive to develop obesity on HFD than female mice. In addition, genetic variation and modification in mouse models can affect metabolic function, and therefore the effects of HFD exposure may differ between mouse models. In most studies on diet-induced obesity, mice are fed on a HFD for 8–15 weeks (421).

In severe obese individuals the standard lifestyle recommendations fail, or is not compatible with modern life routines. Therefore, alternative strategies for intervention are urgently required. Preventive nutritional strategies during critical periods in life seem promising to reduce the risk of obesity development. For instance, omega-3 fatty acids are able to prevent body weight gain, hepatic lipid accumulation by inducing hepatic fatty acid oxidation (421, 443, 444). We revealed that the intake of LCPUFAs, ARA and DHA, in early life can protect against

HFD-induced-obesity in late life (**Chapter 2**). In children daily consumption of DHA and eicosapentaenoic acid (EPA) improved anthropometric and lipid parameters (27). These findings confirm that ARA and DHA dietary intake in early life strengthens the ability to cope with an obesogenic diet. In **chapter 3** we found that butyrate, a short chain fatty acid (SCFA), restores HFD-induced obesity on metabolic, cerebrovasculature and cognitive function. Possible mechanisms by which ARA, DHA and butyrate can protect or restore HFD-induced obesity and co-morbidities is still incomplete. Although, it has been reported that ARA and DHA intake results in variations in adipocyte differentiation, whereby storage of WAT can be affected (262). Moreover, DHA and ARA are the two most abundant LCPUFAs in the brain, and are essential to support optimal brain and visual system development (204, 205). Butyrate has anti-inflammatory properties, contributes to fatty acid oxidation, and can induce a switch from lipid synthesis to lipid utilization (298, 308). In addition, butyrate most likely affects cognitive function via enhancing histone acetylation by inhibiting histone deacetylase (283). Dynamic changes in histone acetylation have been linked to the genetic programming required for memory formation (283). Additionally, adiposity can be affected by the gut microbiome through energy harvesting and lipid storage by the bacteria. Moreover, butyrate is a product of bacterial fermentation of mainly indigestible plant polysaccharides and resistant starch in the colon, and obese individuals have a decreased level of butyrate producing microbiota (23). Furthermore, analysis of the gut microbiome revealed pronounced changes in composition upon HFD feeding (**Chapter 3**). These HFD induced changes were very consistent with observations made by others (312). In line with observations made in obese humans which exhibit decreased levels of butyrate-producing bacteria compared to lean individuals (24), we found HFD induced in *Clostridium XIVa* abundancy (**Chapter 3**). Moreover, the pronounced difference between the microbiome of mice fed a control diet or fed a HFD made it difficult to discriminate the specific effects of butyrate (**Chapter 3**). However, in literature it was reported that both butyrate and omega-3 fatty acids modify microbiome composition, and the expression of anti-inflammatory compounds (445-447). Thereby, butyrate and DHA are able to affect adiposity by modifying microbiome composition. Thus, fatty acids like DHA and butyrate, exert a moreover protective effect in adiposity and its co-morbidities. Nevertheless, it has to be taken into account that dietary interventions might provoke slightly altered effects in other mouse models as their metabolic function may differ due to genetic modification.

In humans, LCPUFAs like ARA and DHA are added in milk formula's and they are beneficially affecting the development of infants (448). In addition, in adults LCPUFAs improve vascular endothelial function and can reduce blood pressure,

circulating lipids, and inflammatory processes (449, 450). We indicated that SCFAs, like butyrate, may serve as potential preventative for obesity and obesity related diseases (**Chapter 3**). However, up till now it is rather difficult for obese individuals to obtain a relatively high concentration of butyrate within their blood plasma. Some adversities to overcome are for example the distinctive odor of butyrate, and relatively low uptake of butyrate in the blood circulation when consuming fiber rich diets (451). Nevertheless, *Van der Beek et al.* implied that rectal butyrate administration was safe at a dose of 100 mmol sodium butyrate/L, and that it led to higher portal butyrate concentrations compared with placebo (452). Future research could address the effectiveness of a lower concentration of butyrate and/or capsules that release butyrate within the intestine.

Concluding remarks of this thesis research

Overall, in this thesis we clearly demonstrate a link between adiposity and brain function. Animal studies have the potential to examine biological mechanisms and dietary interventions within the adiposity-brain-axis in great detail. Moreover, with this thesis we have furthered the understanding of biological mechanisms by which adiposity can affect brain function, e.g. via adipokines like leptin or affecting cerebral circulation. However, it seems to be a highly interactive and multi-factorial process, which is clearly altered by sex and during aging. Therefore, future research should account for age and sex-hormones while examining the adiposity-brain-axis. Moreover, we underline the role of diets in health. Firstly, intake of LCPUFAs (ARA and DHA) in early life prevents WAT accumulation, improves gut microbiome and preserves brain function. Secondly, a butyrate intervention, and plausible the intake of indigestible fibers (butyrate is a product of bacterial fermentation of indigestible fibers), can counteract HFD-induced adverse health effect in midlife in adiposity, gut microbiome, cerebrovasculature, liver and brain function. Thereby, balanced diets containing undigested fibers and LCPUFAs have a crucial role in attenuating the genesis of adiposity and its co-morbidities throughout lifespan.

7

APPENDICES

Summary

Worldwide, obesity has reached epidemic proportions, and each year obesity or obesity-related-diseases lead to the death of 2.8 million adults. Obesity has an undefined aetiology, but is generally caused by imbalance of energy intake versus energy output. This imbalance results in excessive, mainly white adipose tissue (WAT) accumulation, adiposity, which eventually may have an adverse effect on human health, such as cardiovascular disease, type 2 diabetes mellitus and cognitive impairment. The aim of the research presented in this thesis was to elucidate the adiposity-brain-axis involving dietary fatty acids (arachidonic acid (ARA), docosahexaenoic acid (DHA) and butyrate), (cerebro)vasculature, adipokines, and the effect of aging on these processes.

Chapter 1 provides the background information for this thesis research, including the role of diets in the progression, and contrarily the prevention, of adiposity. Moreover, we discuss two pathways via which adiposity might affect brain structure, function and cognition with emphasis on the (cerebro)vascular pathway and adipokines. In short, diets with a high content of saturated fat (HFD) or high fat and high carbohydrates (HFHC), are considered to be important contributing factors of obesity, whereas dietary interventions containing adequate levels of LCPUFAs, butyrate or even fiber rich diets, could contribute to the decline of adiposity. Moreover, we hypothesize that adiposity can induce (cerebro)vascular changes such as arterial stiffness, inflammation and hypoperfusion, which can subsequently disrupt brain structure and function, and increase the risk for cognitive impairment and dementia. Lastly, we speculate that adipokine release may be dysregulated in adiposity, possibly because the adipose tissue may be 'diseased'. Since neurodegenerative and vascular processes observed in dementia affect several brain regions and nuclei, the action of adipokines might be altered during neurodegenerative and vascular events, and may even feedback to contribute to neurodegeneration.

In the first experimental work, described in **chapter 2**, we examined the effect of early life dietary long chain polyunsaturated fatty acids (LCPUFAs, ARA and DHA) intake in mild-obese transgenic ApoE3*Leiden mice. A broad and multidisciplinary experimental approach was used to examine energy metabolism, brain structure, cerebral circulation, and cognition. We found that a HFD diet causes adverse structural brain and metabolic adaptations in mildly obese mice, like a decrease in functional connectivity and gray matter integrity, and an increase in cholesterol and triglycerides levels. Importantly, most of these alterations due to HFD can be averted by dietary intake of ARA and DHA for 8 weeks early in life, which supports metabolic flexibility and cerebral integrity later in life.

In **chapter 3** a severe mouse model of obesity, low-density lipoprotein receptor knock-out Leiden (LDLr^{-/-}) mouse, have been examined for the development of HFD-induced-obesity, inflammation within WAT, increased fat storage within the liver, increased systolic blood pressure, and cerebral hypoperfusion (**Chapter 3**). Subsequently, we aimed to examine the preventive abilities of short chain fatty acid, butyrate in obese LDLr^{-/-} male mice. We found that butyrate restored HFD-induced-obesity parameters towards normal levels in two months in mid-adult obese mice, for instance it restored body weight, neuroinflammation, liver steatosis, cerebral blood flow and functional connectivity. Additionally, we found that a HFD can lower the variance the gut microbiome, and thereby a HFD can affect adiposity through energy harvesting and lipid storage by the bacteria. Moreover, butyrate could modify microbiome composition, nevertheless the pronounced effects of HFD made it difficult to discriminate the specific effects of butyrate. Furthermore, we suggest that butyrate may have potential in prevention and treatment of midlife obesity.

In **chapter 4 and 5** we aimed to analyze associations concerning the adiposity-brain-axis in humans. In the RUN DMC study (**Chapter 4**), we found that adiposity indicators, body mass index (BMI), waist circumference (WC) and leptin are negatively associated with gray matter volume (GMV) in men diagnosed with cerebral small vessel disease (CSVD) at baseline. BMI and WC were also inversely associated with hippocampal volume in men. At follow-up, 9 years later, cross-sectional analyses showed inverse associations between BMI and GMV; and WC and hippocampal volume, particularly in men. In relation to CSVD outcomes, the metabolic adiposity indicator, leptin, was protective in men. In particular, the highest quartile of leptin (compared to the first quartile) was inversely associated with severe white matter hyperintensities, the presence of any lacunes and the presence of at least one component of CSVD. These findings indicate that adiposity measures, both anthropometric and metabolic were differentially associated with brain volumes and CSVD, and by sex. These findings suggest an Alzheimer-vascular spectrum associated with adiposity, a known marker of both neurodegeneration and vascular risk, that both influences and is influenced by brain pathologies associated with dementia.

In **chapter 5**, we examined in older Swedish adults individuals participating in the Gothenburg studies. These Gothenburg studies provided us a 10-year prospective population-based study of 70 years and older inhabitants of Gothenburg, Sweden. Subsequently, we analyzed if anthropometric parameters as BMI and waist-hip ratio, and serum adipokines levels like adiponectin and leptin could predict dementia risk in older Swedish adults over a 5 and 10 year period. In addition, we examined whether these associations are different between older men and women.

We found that in late-life, anthropometric and metabolic adiposity measures appear to be differentially associated with dementia risk in women aged 70 years and older (Gothenburg). In detail, while BMI and leptin levels are highly positively correlated, our results show that their association with dementia at age ≥ 70 years, is asynchronous. Leptin is associated with decreased dementia risk, whereas low BMI ($<20 \text{ kg/m}^2$) predicted an increased dementia risk over a 5 year period.

In **chapter 6**, we provide an extensive discussion of the research findings summarized above, and furthermore we propose follow-up studies are needed to address the questions that have arisen from our work. The most important conclusions of this thesis research are: 1) A clear link between adiposity and brain function. 2) The potential of animal studies to examine biological mechanisms and dietary interventions within the adiposity-brain-axis in great detail. 3) Intake of LCPUFAs in early life prevents WAT accumulation, and preserves brain function. 4) Butyrate intervention and plausible the intake of indigestible fibers (butyrate is a product of bacterial fermentation of indigestible fibers), can counteract HFD-induced adverse health effect in midlife in adiposity, gut microbiome, cerebrovasculature, liver and brain function. 5) The adiposity-brain-axis is a highly interactive and multi-factorial process, which is clearly altered by sex and during aging. Overall, we think that the research described in this thesis has furthered the understanding of biological mechanisms by which adiposity can affect brain function. Thereby, balanced diets containing undigested fibers and LCPUFAs have a crucial role in attenuating the genesis of adiposity and its co-morbidities throughout lifespan.

Nederlandse Samenvatting

Obesitas is uitgegroeid tot een wereldwijde epidemie, en elk jaar resulteren obesitas of obesitas-gerelateerde-ziekten in de dood van 2.8 miljoen volwassenen. Obesitas heeft onbepaalde etiologie, maar wordt over het algemeen veroorzaakt door een disbalans tussen energie inname en energie verbranding. Deze disbalans resulteert in extreme vetopslag van vooral het zogenaamde witte vet. Deze extreme vetopslag wordt ook wel adipositas of vetzucht genoemd. Adipositas heeft een negatieve invloed op de gezondheid en kan uiteindelijk bijdragen aan het ontwikkelen van ziekten, zoals bijvoorbeeld cardiovasculaire ziekten, type 2 diabetes mellitus of zelfs cognitieve achteruitgang en dementie. In het onderzoek beschreven in dit proefschrift hebben we onderzocht of de adipositas-brein-as beïnvloed kan worden door vetzuren in het dieet (arachidonzuur (ARA), docosahexaeenzuur (DHA) en butyraat) en daarnaast, of er in deze as een rol is weggelegd voor de hersendoorbloedingen adipokinen. De laatste genoemde, adipokinen zijn hormonen, eiwitten en cytokinen geproduceerd door het witte vetweefsel. Als laatste zullen we onderzoeken wat de invloed is van veroudering op de adipositas-brein-as.

Hoofdstuk 1 schetst de achtergrond van het onderzoek beschreven in dit proefschrift. We bespreken in dit hoofdstuk de rol van vetzuren in het ontstaan en tegengaan van adipositas. Zo is het bekend dat diëten met een hoge concentratie aan verzadigd vet en/of in combinatie met veel koolhydraten een belangrijke rol spelen in het veroorzaken van obesitas. Als tegengewicht, zijn er vetzuren die de ontwikkeling adipositas kunnen tegenwerken, voorbeelden zijn lange-keten-meervoudig-onverzadigde-vetzuren (LCPUFAs) en korte-keten-vetzuren (SCFAs, butyraat). De SCFAs kunnen worden gevormd door bacteriën in de darmwand wanneer deze bacteriën vezels verteren. Daaropvolgend bediscussiëren we via welke wegen adipositas de brein functie en cognitie zou kunnen beïnvloeden. Eén van de mogelijke wegen richting het brein zou kunnen zijn door het aantasten van de hersendoorbloeding, namelijk onder andere door verstijving en/of vernauwing van de vaten rondom het brein en ontstekingen in het brein. Deze veranderingen in de hersendoorbloeding kunnen daaropvolgend een negatieve invloed hebben op zenuwcellen in het brein, wat kan resulteren in cognitieve achteruitgang of zelfs dementie. Een andere weg richting het brein is vanuit het witte vetweefsel, zo zou de adipokinen productie verstoord kunnen zijn. Adipokinen zijn betrokken in vele biologische processen en kunnen ook op vele hun invloed uitoefenen. Als de productie van deze adipokinen verstoord is of uit balans doordat er extreem veel vetweefsel is, dan zouden zij neurodegeneratieve en vasculaire processen kunnen beïnvloeden wat uiteindelijk zou kunnen leiden tot cognitieve achteruitgang en dementie.

In het eerste experimentele onderzoek beschreven in **hoofdstuk 2**, onderzochten wij of de inname van de LCPUFAs, ARA en DHA in het vroege leven een beschermend effect kon hebben in muizen (ApoE3*Leiden). Deze muizen kregen drie weken na hun geboorte voor acht weken een dieet met de LCPUFAs, daaropvolgend kregen deze muizen voor 12 weken een hoog vet dieet (HFD). Om de effecten van deze diëten te onderzoeken hebben we een veelzijdige aanpak gebruikt om onder andere veranderingen in energie metabolisme, hersendoorbloeding, het brein en cognitie te kunnen analyseren. Uiteindelijk vonden wij dat een dieet met veel verzadigde vetzuren (hoog vet dieet; HFD) aanpassingen veroorzaakten op metabool vlak en in het brein, zoals bijvoorbeeld een toename van cholesterol en triglyceriden in het bloed, vermindering in verbindingen tussen hersengebieden en afname in grijze stof. Ook konden we concluderen dat de inname van ARA en DHA in het vroege leven bescherming biedt tegen de meeste veranderingen veroorzaakt door het HFD op latere leeftijd. Deze bevinding benadrukt de metabole flexibiliteit en cerebrale integriteit in het latere leven.

In **hoofdstuk 3** beschrijven we een onderzoek met als doel om het effect van korte-keten-vetzuur, butyraat te achterhalen in middelbare en oudere obese muizen. In dit onderzoek gebruikten we muizen die het gen voor de receptor van de lage-dichtheid-lipoproteïne missen, deze zogenaamde LDLr-/-.Leiden muizen. Deze muizen ontwikkelen op een HFD een zware vorm van obesitas. In dit onderzoek waren we met name geïnteresseerd in de ontwikkeling van adipositas, ontsteking van het witte vetweefsel, opslag van vet in de lever, bloeddruk en de cerebrale doorbloeding. We vonden dat een butyraat-interventie van twee maanden de negatieve effecten van een HFD voor een groot deel terug kon brengen naar normale controle waarden, en dit met name in de muizen van middelbare leeftijd. Daarbij ontdekten we dat een HFD de variatie verlaagt van bacteriën in de darm, en daarmee kan een HFD effect hebben op het verkrijgen van energie en vetopslag door bacteriën. Als laatste, vonden we dat butyraat de samenstelling van darmbacteriën kan veranderen, ondanks dat het lastig was om het effect van butyraat te onderscheiden van de effecten van een HFD op het darm bacteriën. We concludeerden uit dit onderzoek dat butyraat een belangrijke rol kan spelen in het tegenwerken van de ontwikkeling van adipositas op middelbare leeftijd.

Wij beschrijven in **hoofdstuk 4 en 5** onderzoeken uitgevoerd in mensen. In beide hoofdstukken hebben we met statische analysemodellen onderzocht of en hoe adipositas parameters zijn geassocieerd met het brein en dementie, om zo de adipositas-brein-as duidelijker te kunnen schetsen in mensen. De adipositas parameters die we analyseerden waren BMI, buikomvang, leptine en adiponectine. In **hoofdstuk 4** analyseerden we in de RUN DMC studie, body mass index (BMI),

buikomvang (WC), leptine en adiponectine (parameters voor adipositas), en in het brein bekeken we de afwijkingen aan de vaten, en grijze en witte stof volumes. De RUN DMC studie bestaat uit mannen en vrouwen die gediagnosticeerd zijn met ziekte aan de kleine vaten in het brein (CSVD). We vonden dat BMI, WC en leptine negatief geassocieerd zijn met volume van de grijze stof, met name in mannen (>50.0 jaar). Dit indiceert dat adipositas geassocieerd is met een lager grijze stof volume in mannen. Daarbij vonden we dat leptine een beschermend effect kan hebben in mannen, met betrekking tot de vasculaire parameters in het brein. In detail, leptine was negatief geassocieerd met verlies van integriteit van de witte stof (zogenaamde white matter hyperintensities) en de aanwezigheid van lacunes in het brein, welke beiden parameters zijn voor CSVD. Deze bevindingen indiceren dat parameters voor adipositas verschillend zijn geassocieerd met brein volume, vasculaire parameters, en geslacht. In **hoofdstuk 4** bediscussiëren we dat adipositas invloed kan hebben op hersenaandoeningen, zoals bijvoorbeeld verminderde grijze stof, en dat deze hersenaandoeningen zijn sterk geassocieerd met dementie.

In **hoofdstuk 5** wordt een Zweeds onderzoek, de Gothenburg cohort studies beschreven. Deze studies bestaan uit Zweedse ouderen, man en vrouw, van 70 jaar en ouder. In deze studie analyseerden we of parameters voor adipositas (BMI, buik-heup-ratio, leptine en adiponectine) geassocieerd zijn met dementie over een periode van tien jaar. Daarbij analyseerden we ook of er een verschil was tussen mannen en vrouwen. Wij vonden dat in oudere vrouwen (>70.0 jaar), leptine en BMI verschillend geassocieerd zijn met het risico op dementia binnen 5-jaar. Dit terwijl je verwacht dat leptine en BMI dezelfde associatie hebben omdat ze sterk met elkaar gecorreleerd zijn. Zo is leptine is geassocieerd met een verminderde kans op dementie, terwijl een lage BMI (<20 kg/m²) een verhoogd risico geeft om dementie te ontwikkelen binnen 5 jaar.

We bediscussiëren al onze resultaten in **hoofdstuk 6**, en daarnaast stellen we mogelijke andere toekomstige onderzoeken voor die vragen zou kunnen beantwoorden ontstaan uit het onderzoek beschreven in dit proefschrift. Ook bespreken we de belangrijkste conclusies van dit proefschrift, deze zijn: 1) er is een duidelijk verband tussen adipositas en brein functie. 2) Onderzoek in muismodellen biedt de kans om biologische processen en dieet interventies betrokken in de adipositas-brein-as in diepgaand detail te onderzoeken. 3) De inname van LCPUFAs in het vroege leven beschermt tegen adipositas, en behoudt van een gezonde brein functie. 4) Een butyraat interventie, en mogelijk de inname van vezels, kan de negatieve effecten van een HFD-dieet tegenwerken met name op middelbare leeftijd op het gebied van darm microbioom, hersendoorbloeding, lever en brein functie.

5) De adipositas-brein-as is een interactieve en multifactoriële as, die zeker wordt beïnvloed door geslacht en gedurende veroudering. Al met al, denken wij dat het onderzoek beschreven in dit proefschrift bijdraagt aan een verbeterd inzicht in de biologische mechanismen betrokken in de adipositas-brein-as. Uitgebalanceerde vezelrijke diëten met LCPUFAs kunnen een belangrijke rol spelen in het tegengaan van adipositas en de ontwikkeling van obesitas-gerelateerde-ziekten.

List of Abbreviations

AC	Auditory cortex
AD	Alzheimer's disease
ADAS-Cog	Alzheimer's disease assessment scale-cognitive subscale
AgRP	Agouti-related peptide
ANOVA	Analysis of variance
APOC1	Apolipoprotein C-I
APOE	Apolipoprotein E
APOE3	Apolipoprotein E3
ApoE3*Leiden	Apolipoprotein E3 Leiden
APOEε4	Apolipoprotein E ε4 allele
ARA	Arachidonic acid
ASL	arterial spin labeling
BBB	Blood brain barrier
BMI	Body mass index
BOLD	Blood oxygen level-dependent
CA1	Cornu Ammonis area 1,
CART	cocaine and amphetamine-regulated transcript
CBF	Cerebral blood flow
cDNA	Complementary deoxyribonucleic acid
CM	Custom made
CNS	Central nervous system
CPRS	Comprehensive psychiatric rating scale
CRP	C-reactive protein
CSF	Cerebrospinal fluid
CSVD	Cerebral small vessel disease
CT	Cycle threshold
CVD	Cardiovascular disease
DH	Dorsal hippocampus
DHA	Docosahexaenoic
DNA	Deoxyribonucleic acid
DSM-III-R	Diagnostic and statistical manual of mental disorders 3rd edition revised
DTI	Diffusion tensor imaging
ELISA	Enzyme-linked immunosorbent assay
EPA	Eicosapentaenoic acid
FA	Fractional anisotropy
FAIR	Flow-sensitive alternating inversion recovery
FC	Functional connectivity

FFAR2	Free fatty acid receptor 2
FFAR3	Free fatty acid receptor 3
fMRI	Functional magnetic resonance imaging
FW	Forward
GAPDH	Glyceraldehydes 3-phosphate dehydrogenase
GLUT-1	Glucose transporter 1
GMV	Gray matter volume
H70	Gerontological and geriatric population studies
HDAC	Histone deacetylase
HDL	High density lipoprotein
HFD	High fat diet
HFDB	High fat diet enriched with butyrate
HFHC	High fat and high carbohydrates diet
HMW	High molecular weight
HV	Hippocampal volume
IBA-1	Ionized calcium binding adapter molecule-1
ICV	Intracranial volume
IL-1 β	Interleukin-1 beta
IL-6	Interleukin-6
InsulinR	Insulin receptor
IVC	Individually ventilated cages
LCPUFAs	Long-chain polyunsaturated fatty acids
LDL	Low-density lipoprotein
LDL	Low density lipoprotein
LDLr	Low-density lipoprotein receptor
LDLr-/-	Low-density lipoprotein receptor knock-out Leiden
LTP	Long term potentiation
m.o.	Months old
MC	Motor cortex
MCP-1	Monocyte chemotactic protein-1
MD	Mean diffusivity
MED	Minimum entropy decomposition
MINI	Mini international neuropsychiatric interview
MMSE	Mini-Mental State Examination
MRI	Magnetic resonance imaging
mRNA	Messenger ribonucleic acid
MWM	Morris water maze
n	Group size
NALFD	Non-alcoholic fatty liver disease
NaN3	Sodium azide
NE	North-East

NE	Not estimable
NMDA	N-methyl-D-aspartate
NPY	Neuropeptide Y
ObRA	Leptin receptor A
OBRB	Leptin receptor B
OF	Open field
OR	Odds ratio
ORT	Object recognition test
PAI-1	Plasminogen activator inhibitor
PBS	Phosphate-buffered saline
pCVD	Premature cardiovascular disease
PET	Positron emission tomography
PPAR γ	Peroxisome proliferator-activated receptor- γ
PPSW	Prospective population study of women
PRIME	Preclinical imaging centre
PSD-95	Postsynaptic density protein-95
Q	Quartile
qRT-PCR	Quantitative real time polymerase chain reaction
REV	Reverse
RNA	Ribonucleic acid
ROI	Region of interest
rRNA	Ribosomal ribonucleic acid
rsfMRI	Resting state functional magnetic resonance imaging
RUN DMC	Radboud University Nijmegen Diffusion tensor and Magnetic resonance imaging Cohort
SAA	Serum amyloid A
SBP	Systolic blood pressure
SCFAs	Short chain fatty acids
SD	Standard deviations
SEM	Standard error of mean
SPF	Specific pathogen free
SSC	Somatosensory cortex
STRIVE	Strategies and therapies for reducing ischemic and vascular events
T2DM	type two diabetes mellitus
TBV	Total brain volume
TIA	Transient ischemic attack
TNF- α	Tumor necrosis factor alpha
TNO	Toegepast-natuurwetenschappelijk onderzoek
VC	Visual cortex
VCAM-1	Vascular cell adhesion molecule 1

VEGF	Vascular endothelial growth factor
VEGF-A	Vascular endothelial growth factor A
VH	Ventral hippocampus
VLDL	Very low density lipoprotein
w/w	Weight by weight
WAT	White adipose tissue
WC	Waist circumference
WHR	Waist hip ratio
WMH	White matter hyperintensities
WML	White matter lesion
WMV	White matter volume
α -MSH	Alpha-melanocyte-stimulating hormone
β (95% C.I.)	Beta-coefficients and 95% confidence Intervals

References

1. WHO WHO. Obesity and Overweight 2017 [updated May 2018. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>.
2. Anstey KJ, Cherbuin N, Budge M, Young J. Body mass index in midlife and late-life as a risk factor for dementia: a meta-analysis of prospective studies. *Obes Rev*. 2011;12(5):e426-37.
3. Emmerzaal TL, Kiliaan AJ, Gustafson DR. 2003-2013: a decade of body mass index, Alzheimer's disease, and dementia. *J Alzheimers Dis*. 2015;43(3):739-55.
4. Nevill AM, Stewart AD, Olds T, Holder R. Relationship between adiposity and body size reveals limitations of BMI. *Am J Phys Anthropol*. 2006;129(1):151-6.
5. Stevens J, Cai J, Juhaeri, Thun MJ, Williamson DF, Wood JL. Consequences of the use of different measures of effect to determine the impact of age on the association between obesity and mortality. *Am J Epidemiol*. 1999;150(4):399-407.
6. Ferland M, Despres JP, Tremblay A, Pinault S, Nadeau A, Moorjani S, et al. Assessment of adipose tissue distribution by computed axial tomography in obese women: association with body density and anthropometric measurements. *The British journal of nutrition*. 1989;61(2):139-48.
7. Pouliot MC, Despres JP, Lemieux S, Moorjani S, Bouchard C, Tremblay A, et al. Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *Am J Cardiol*. 1994;73(7):460-8.
8. Seidell JC, Oosterlee A, Deurenberg P, Hautvast JG, Ruijs JH. Abdominal fat depots measured with computed tomography: effects of degree of obesity, sex, and age. *European journal of clinical nutrition*. 1988;42(9):805-15.
9. Janssen I, Katzmarzyk PT, Ross R. Body mass index, waist circumference, and health risk: evidence in support of current National Institutes of Health guidelines. *Arch Intern Med*. 2002;162(18):2074-9.
10. Sanchez-Castillo CP, Velasquez-Monroy O, Lara-Esqueda A, Berber A, Sepulveda J, Tapia-Conyer R, et al. Diabetes and hypertension increases in a society with abdominal obesity: results of the Mexican National Health Survey 2000. *Public Health Nutr*. 2005;8(1):53-60.
11. Das UN. Is obesity an inflammatory condition? *Nutrition*. 2001;17(11-12):953-66.
12. Antuna-Puente B, Feve B, Fellahi S, Bastard JP. Adipokines: the missing link between insulin resistance and obesity. *Diabetes Metab*. 2008;34(1):2-11.
13. Garcia C, Feve B, Ferre P, Halimi S, Baizri H, Bordier L, et al. Diabetes and inflammation: fundamental aspects and clinical implications. *Diabetes Metab*. 2010;36(5):327-38.
14. Luchtman DW, Song C. Cognitive enhancement by omega-3 fatty acids from child-hood to old age: findings from animal and clinical studies. *Neuropharmacology*. 2013;64:550-65.

15. Arnoldussen IA, Kiliaan AJ. Impact of DHA on metabolic diseases from womb to tomb. *Mar Drugs*. 2014;12(12):6190-212.
16. Williams JH, Errington ML, Lynch MA, Bliss TV. Arachidonic acid induces a long-term activity-dependent enhancement of synaptic transmission in the hippocampus. *Nature*. 1989;341(6244):739-42.
17. Itariu BK, Zeyda M, Hochbrugger EE, Neuhofer A, Prager G, Schindler K, et al. Long-chain n-3 PUFAs reduce adipose tissue and systemic inflammation in severely obese nondiabetic patients: a randomized controlled trial. *Am J Clin Nutr*. 2012;96(5):1137-49.
18. Alosco ML, Spitznagel MB, Raz N, Cohen R, Sweet LH, Colbert LH, et al. Obesity interacts with cerebral hypoperfusion to exacerbate cognitive impairment in older adults with heart failure. *Cerebrovascular diseases extra*. 2012;2(1):88-98.
19. Ellis EF, Police RJ, Yancey LM, Grabeel MN, Heizer ML, Holt SA. Effect of fish oil n-3 fatty acids on cerebral microcirculation. *Am J Physiol*. 1990;258(6 Pt 2):H1780-5.
20. Jin CJ, Sellmann C, Engstler AJ, Ziegenhardt D, Bergheim I. Supplementation of sodium butyrate protects mice from the development of non-alcoholic steatohepatitis (NASH). *Br J Nutr*. 2015;114(11):1745-55.
21. Gao Z, Yin J, Zhang J, Ward RE, Martin RJ, Lefevre M, et al. Butyrate improves insulin sensitivity and increases energy expenditure in mice. *Diabetes*. 2009;58(7):1509-17.
22. Lin HV, Frassetto A, Kowalik EJ, Jr., Nawrocki AR, Lu MM, Kosinski JR, et al. Butyrate and propionate protect against diet-induced obesity and regulate gut hormones via free fatty acid receptor 3-independent mechanisms. *PLoS One*. 2012;7(4):e35240.
23. Canfora EE, Jocken JW, Blaak EE. Short-chain fatty acids in control of body weight and insulin sensitivity. *Nature reviews Endocrinology*. 2015;11(10):577-91.
24. Elli M, Colombo O, Tagliabue A. A common core microbiota between obese individuals and their lean relatives? Evaluation of the predisposition to obesity on the basis of the fecal microflora profile. *Medical hypotheses*. 2010;75(4):350-2.
25. Decsi T, Minda H, Hermann R, Kozari A, Erhardt E, Burus I, et al. Polyunsaturated fatty acids in plasma and erythrocyte membrane lipids of diabetic children. Prostaglandins, leukotrienes, and essential fatty acids. 2002;67(4):203-10.
26. Burrows T, Collins CE, Garg ML. Omega-3 index, obesity and insulin resistance in children. *Int J Pediatr Obes*. 2011;6(2-2):e532-9.
27. Vasickova L, Stavek P, Suchanek P. Possible effect of DHA intake on body weight reduction and lipid metabolism in obese children. *Neuro Endocrinol Lett*. 2011;32 Suppl 2:64-7.
28. Juarez-Lopez C, Klunder-Klunder M, Madrigal-Azcarate A, Flores-Huerta S. Omega-3 polyunsaturated fatty acids reduce insulin resistance and triglycerides in obese children and adolescents. *Pediatric diabetes*. 2013;14(5):377-83.
29. Damsgaard CT, Stark KD, Hjorth MF, Biloft-Jensen A, Astrup A, Michaelsen KF, et al. n-3 PUFA status in school children is associated with beneficial lipid profile, reduced physical activity and increased blood pressure in boys. *Br J Nutr*. 2013;110(7):1304-12.

30. Dangardt F, Chen Y, Gronowitz E, Dahlgren J, Friberg P, Strandvik B. High physiological omega-3 Fatty Acid supplementation affects muscle Fatty Acid composition and glucose and insulin homeostasis in obese adolescents. *Journal of nutrition and metabolism*. 2012;2012:395757.
31. Rivellese AA, Maffettone A, Iovine C, Di Marino L, Annuzzi G, Mancini M, et al. Long-term effects of fish oil on insulin resistance and plasma lipoproteins in NIDDM patients with hypertriglyceridemia. *Diabetes Care*. 1996;19(11):1207-13.
32. Mori TA, Bao DQ, Burke V, Puddey IB, Beilin LJ. Docosahexaenoic acid but not eicosapentaenoic acid lowers ambulatory blood pressure and heart rate in humans. *Hypertension*. 1999;34(2):253-60.
33. Mori TA, Watts GF, Burke V, Hilme E, Puddey IB, Beilin LJ. Differential effects of eicosapentaenoic acid and docosahexaenoic acid on vascular reactivity of the forearm microcirculation in hyperlipidemic, overweight men. *Circulation*. 2000;102(11):1264-9.
34. Woodman RJ, Mori TA, Burke V, Puddey IB, Watts GF, Best JD, et al. Docosahexaenoic acid but not eicosapentaenoic acid increases LDL particle size in treated hypertensive type 2 diabetic patients. *Diabetes Care*. 2003;26(1):253.
35. Kelley DS, Siegel D, Vemuri M, Mackey BE. Docosahexaenoic acid supplementation improves fasting and postprandial lipid profiles in hypertriglyceridemic men. *Am J Clin Nutr*. 2007;86(2):324-33.
36. Sneddon AA, Tsofliou F, Fyfe CL, Matheson I, Jackson DM, Horgan G, et al. Effect of a conjugated linoleic acid and omega-3 fatty acid mixture on body composition and adiponectin. *Obesity (Silver Spring)*. 2008;16(5):1019-24.
37. Micallef M, Munro I, Phang M, Garg M. Plasma n-3 Polyunsaturated Fatty Acids are negatively associated with obesity. *Br J Nutr*. 2009;102(9):1370-4.
38. Stirban A, Nandrea S, Gotting C, Tamler R, Pop A, Negrean M, et al. Effects of n-3 fatty acids on macro- and microvascular function in subjects with type 2 diabetes mellitus. *Am J Clin Nutr*. 2010;91(3):808-13.
39. Labonte ME, Couture P, Tremblay AJ, Hogue JC, Lemelin V, Lamarche B. Eicosapentaenoic and docosahexaenoic acid supplementation and inflammatory gene expression in the duodenum of obese patients with type 2 diabetes. *Nutrition journal*. 2013;12(1):98.
40. Singhal A, Lanigan J, Storry C, Low S, Birbara T, Lucas A, et al. Docosahexaenoic Acid supplementation, vascular function and risk factors for cardiovascular disease: a randomized controlled trial in young adults. *Journal of the American Heart Association*. 2013;2(4):e000283.
41. McDonald DM, O'Kane F, McConville M, Devine AB, McVeigh GE. Platelet redox balance in diabetic patients with hypertension improved by n-3 fatty acids. *Diabetes Care*. 2013;36(4):998-1005.
42. Virtanen JK, Mursu J, Voutilainen S, Uusitupa M, Tuomainen TP. Serum Omega-3 Polyunsaturated Fatty Acids and Risk of Incident Type 2 Diabetes in Men: The Kuopio Ischaemic Heart Disease Risk Factor Study. *Diabetes Care*. 2013.

43. Woodman RJ, Mori TA, Burke V, Puddey IB, Barden A, Watts GF, et al. Effects of purified eicosapentaenoic acid and docosahexaenoic acid on platelet, fibrinolytic and vascular function in hypertensive type 2 diabetic patients. *Atherosclerosis*. 2003;166(1):85-93.
44. Lemaitre RN, King IB, Mozaffarian D, Kuller LH, Tracy RP, Siscovick DS. n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr*. 2003;77(2):319-25.
45. Tsitouras PD, Gucciardo F, Salbe AD, Heward C, Harman SM. High omega-3 fat intake improves insulin sensitivity and reduces CRP and IL6, but does not affect other endocrine axes in healthy older adults. *Horm Metab Res*. 2008;40(3):199-205.
46. Heine-Broring RC, Brouwer IA, Proenca RV, van Rooij FJ, Hofman A, Oudkerk M, et al. Intake of fish and marine n-3 fatty acids in relation to coronary calcification: the Rotterdam Study. *Am J Clin Nutr*. 2010;91(5):1317-23.
47. Djousse L, Biggs ML, Lemaitre RN, King IB, Song X, Ix JH, et al. Plasma omega-3 fatty acids and incident diabetes in older adults. *Am J Clin Nutr*. 2011;94(2):527-33.
48. Mozaffarian D, Lemaitre RN, King IB, Song X, Spiegelman D, Sacks FM, et al. Circulating long-chain omega-3 fatty acids and incidence of congestive heart failure in older adults: the cardiovascular health study: a cohort study. *Ann Intern Med*. 2011;155(3):160-70.
49. Virtanen JK, Nyantika AN, Kauhanen J, Voutilainen S, Tuomainen TP. Serum long-chain n-3 polyunsaturated fatty acids, methylmercury and blood pressure in an older population. *Hypertension research : official journal of the Japanese Society of Hypertension*. 2012;35(10):1000-4.
50. Gustafson DR. A life course of adiposity and dementia. *Eur J Pharmacol*. 2008;585(1):163-75.
51. Gustafson DR. Adiposity hormones and dementia. *Journal of the neurological sciences*. 2010;299(1-2):30-4.
52. Gustafson DR, Steen B, Skoog I. Body mass index and white matter lesions in elderly women. An 18-year longitudinal study. *Int Psychogeriatr*. 2004;16(3):327-36.
53. Businaro R, Ippoliti F, Ricci S, Canitano N, Fuso A. Alzheimer's disease promotion by obesity: induced mechanisms-molecular links and perspectives. *Curr Gerontol Geriatr Res*. 2012;2012:986823.
54. Haltia LT, Viljanen A, Parkkola R, Kemppainen N, Rinne JO, Nuutila P, et al. Brain white matter expansion in human obesity and the recovering effect of dieting. *J Clin Endocrinol Metab*. 2007;92(8):3278-84.
55. Delgado TC, Violante IR, Nieto-Charques L, Cerdan S. Neuroglial metabolic compartmentation underlying leptin deficiency in the obese ob/ob mice as detected by magnetic resonance imaging and spectroscopy methods. *J Cereb Blood Flow Metab*. 2011;31(12):2257-66.

56. Brooks SJ, Benedict C, Burgos J, Kempton MJ, Kullberg J, Nordenskjold R, et al. Late-life obesity is associated with smaller global and regional gray matter volumes: a voxel-based morphometric study. *Int J Obes (Lond)*. 2013;37(2):230-6.
57. Pannacciulli N, Del Parigi A, Chen K, Le DS, Reiman EM, Tataranni PA. Brain abnormalities in human obesity: a voxel-based morphometric study. *Neuroimage*. 2006;31(4):1419-25.
58. Raji CA, Ho AJ, Parikshak NN, Becker JT, Lopez OL, Kuller LH, et al. Brain structure and obesity. *Hum Brain Mapp*. 2010;31(3):353-64.
59. Stanek KM, Grieve SM, Brickman AM, Korgaonkar MS, Paul RH, Cohen RA, et al. Obesity is associated with reduced white matter integrity in otherwise healthy adults. *Obesity (Silver Spring)*. 2011;19(3):500-4.
60. Hassenstab JJ, Sweet LH, Del Parigi A, McCaffery JM, Haley AP, Demos KE, et al. Cortical thickness of the cognitive control network in obesity and successful weight loss maintenance: a preliminary MRI study. *Psychiatry Res*. 2012;202(1):77-9.
61. Nummenmaa L, Hirvonen J, Hannukainen JC, Immonen H, Lindroos MM, Salminen P, et al. Dorsal striatum and its limbic connectivity mediate abnormal anticipatory reward processing in obesity. *PLoS One*. 2012;7(2):e31089.
62. Stoeckel LE, Kim J, Weller RE, Cox JE, Cook EW, 3rd, Horwitz B. Effective connectivity of a reward network in obese women. *Brain Res Bull*. 2009;79(6):388-95.
63. Cazettes F, Cohen JJ, Yau PL, Talbot H, Convit A. Obesity-mediated inflammation may damage the brain circuit that regulates food intake. *Brain Res*. 2011;1373:101-9.
64. Widya RL, de Roos A, Trompet S, de Craen AJ, Westendorp RG, Smit JW, et al. Increased amygdalar and hippocampal volumes in elderly obese individuals with or at risk of cardiovascular disease. *Am J Clin Nutr*. 2011;93(6):1190-5.
65. Alonso A, Jacobs DR, Jr., Menotti A, Nissinen A, Dontas A, Kafatos A, et al. Cardiovascular risk factors and dementia mortality: 40 years of follow-up in the Seven Countries Study. *J Neurol Sci*. 2009;280(1-2):79-83.
66. Gallagher D, Visser M, Sepulveda D, Pierson RN, Harris T, Heymsfield SB. How useful is body mass index for comparison of body fatness across age, sex, and ethnic groups? *Am J Epidemiol*. 1996;143(3):228-39.
67. Kivipelto M, Ngandu T, Fratiglioni L, Viitanen M, Kareholt I, Winblad B, et al. Obesity and vascular risk factors at midlife and the risk of dementia and Alzheimer disease. *Arch Neurol*. 2005;62(10):1556-60.
68. Rosengren A, Skoog I, Gustafson D, Wilhelmsen L. Body mass index, other cardiovascular risk factors, and hospitalization for dementia. *Arch Intern Med*. 2005;165(3):321-6.
69. Whitmer RA, Gunderson EP, Barrett-Connor E, Quesenberry CP, Jr., Yaffe K. Obesity in middle age and future risk of dementia: a 27 year longitudinal population based study. *BMJ*. 2005;330(7504):1360.

70. Whitmer RA, Sidney S, Selby J, Johnston SC, Yaffe K. Midlife cardiovascular risk factors and risk of dementia in late life. *Neurology*. 2005;64(2):277-81.
71. Atti AR, Palmer K, Volpato S, Winblad B, De Ronchi D, Fratiglioni L. Late-life body mass index and dementia incidence: nine-year follow-up data from the Kungsholmen Project. *J Am Geriatr Soc*. 2008;56(1):111-6.
72. Dahl AK, Loppinen M, Isoaho R, Berg S, Kivela SL. Overweight and obesity in old age are not associated with greater dementia risk. *J Am Geriatr Soc*. 2008;56(12):2261-6.
73. Hughes TF, Borenstein AR, Schofield E, Wu Y, Larson EB. Association between late-life body mass index and dementia: The Kame Project. *Neurology*. 2009;72(20):1741-6.
74. Mielke MM, Zandi PP, Sjogren M, Gustafson D, Ostling S, Steen B, et al. High total cholesterol levels in late life associated with a reduced risk of dementia. *Neurology*. 2005;64(10):1689-95.
75. Ronan L, Alexander-Bloch AF, Wagstyl K, Farooqi S, Brayne C, Tyler LK, et al. Obesity associated with increased brain age from midlife. *Neurobiol Aging*. 2016;47:63-70.
76. Sriram K, Benkovic SA, Miller DB, O'Callaghan JP. Obesity exacerbates chemically induced neurodegeneration. *Neuroscience*. 2002;115(4):1335-46.
77. Arnoldussen IA, Kiliaan AJ, Gustafson DR. Obesity and dementia: adipokines interact with the brain. *Eur Neuropsychopharmacol*. 2014;24(12):1982-99.
78. Castro MG, Venutolo C, Yau PL, Convit A. Fitness, insulin sensitivity, and frontal lobe integrity in adults with overweight and obesity. *Obesity (Silver Spring)*. 2016;24(6):1283-9.
79. Melka MG, Gillis J, Bernard M, Abrahamowicz M, Chakravarty MM, Leonard GT, et al. FTO, obesity and the adolescent brain. *Hum Mol Genet*. 2013;22(5):1050-8.
80. Walther K, Birdsill AC, Glisky EL, Ryan L. Structural brain differences and cognitive functioning related to body mass index in older females. *Hum Brain Mapp*. 2010;31(7):1052-64.
81. Zhang B, Tian X, Tian D, Wang J, Wang Q, Yu C, et al. Altered Regional Gray Matter Volume in Obese Men: A Structural MRI Study. *Front Psychol*. 2017;8.
82. van Greevenbroek MM, Schalkwijk CG, Stehouwer CD. Obesity-associated low-grade inflammation in type 2 diabetes mellitus: causes and consequences. *Neth J Med*. 2013;71(4):174-87.
83. Andersson CX, Gustafson B, Hammarstedt A, Hedjazifar S, Smith U. Inflamed adipose tissue, insulin resistance and vascular injury. *Diabetes Metab Res Rev*. 2008;24(8):595-603.
84. Heni M, Kullmann S, Preissl H, Fritsche A, Haring HU. Impaired insulin action in the human brain: causes and metabolic consequences. *Nature reviews Endocrinology*. 2015;11(12):701-11.
85. Stillman CM, Weinstein AM, Marsland AL, Gianaros PJ, Erickson KI. Body-Brain Connections: The Effects of Obesity and Behavioral Interventions on Neurocognitive Aging. *Frontiers in aging neuroscience*. 2017;9:115.

86. Bender SB, McGraw AP, Jaffe IZ, Sowers JR. Mineralocorticoid receptor-mediated vascular insulin resistance: an early contributor to diabetes-related vascular disease? *Diabetes*. 2013;62(2):313-9.
87. van Emmerik NM, Renders CM, van de Veer M, van Buuren S, van der Baan-Slootweg OH, Kist-van Holthe JE, et al. High cardiovascular risk in severely obese young children and adolescents. *Arch Dis Child*. 2012.
88. Kiliaan AJ, Arnoldussen IA, Gustafson DR. Adipokines: a link between obesity and dementia? *Lancet Neurol*. 2014;13(9):913-23.
89. Debetto S, Wolf C, Lambert JC, Crivello F, Soumare A, Zhu YC, et al. Abdominal obesity and lower gray matter volume: a Mendelian randomization study. *Neurobiol Aging*. 2014;35(2):378-86.
90. Karlsson HK, Tuulari JJ, Hirvonen J, Lepomäki V, Parkkola R, Hiltunen J, et al. Obesity is associated with white matter atrophy: A combined diffusion tensor imaging and voxel-based morphometric study. *Obesity*. 2013;21(12):2530-7.
91. Mueller K, Horstmann A, Moller HE, Anwander A, Lepsien J, Schroeter ML, et al. Obesity Associated Cerebral Gray and White Matter Alterations Are Interrelated in the Female Brain. *PLoS One*. 2014;9(12):e114206.
92. Ou X, Andres A, Pivik RT, Cleves MA, Badger TM. Brain gray and white matter differences in healthy normal weight and obese children. *J Magn Reson Imaging*. 2015.
93. Gunstad J, Paul RH, Cohen RA, Tate DF, Gordon E. Obesity is associated with memory deficits in young and middle-aged adults. *Eating and weight disorders : EWD*. 2006;11(1):e15-9.
94. Gunstad J, Paul RH, Cohen RA, Tate DF, Spitznagel MB, Gordon E. Elevated body mass index is associated with executive dysfunction in otherwise healthy adults. *Comprehensive psychiatry*. 2007;48(1):57-61.
95. Li Y, Dai Q, Jackson JC, Zhang J. Overweight is associated with decreased cognitive functioning among school-age children and adolescents. *Obesity (Silver Spring)*. 2008;16(8):1809-15.
96. Prunet-Marcassus B, Cousin B, Caton D, Andre M, Penicaud L, Casteilla L. From heterogeneity to plasticity in adipose tissues: site-specific differences. *Experimental cell research*. 2006;312(6):727-36.
97. Campbell DR, Kurzer MS. Flavonoid inhibition of aromatase enzyme activity in human preadipocytes. *The Journal of steroid biochemistry and molecular biology*. 1993;46(3):381-8.
98. Avram AS, Avram MM, James WD. Subcutaneous fat in normal and diseased states: 2. Anatomy and physiology of white and brown adipose tissue. *J Amer Acad Dermatology*. 2005;53(4):671-83.
99. Sbarbati A, Accorsi D, Benati D, Marchetti L, Orsini G, Rigotti G, et al. Subcutaneous adipose tissue classification. *Eur J Histochem*. 2010;54(4):e48.

100. Vindigni V, Tonello C, Lancerotto L, Abatangelo G, Cortivo R, Zavan B, et al. Preliminary report of in vitro reconstruction of a vascularized tendonlike structure: a novel application for adipose-derived stem cells. *Annals of plastic surgery*. 2013;71(6):664-70.
101. Stedman's Medical Dictionary: Lippincott Williams & Wilkins; 2006.
102. Villarroya J, Cereijo R, Villarroya F. An endocrine role for brown adipose tissue? *American journal of physiology Endocrinology and metabolism*. 2013;305(5):E567-72.
103. Esteve Rafols M. Adipose tissue: Cell heterogeneity and functional diversity. *Endocrinología y nutrición : organo de la Sociedad Espanola de Endocrinología y Nutricion*. 2013.
104. Feng B, Zhang T, Xu H. Human adipose dynamics and metabolic health. *Annals of the New York Academy of Sciences*. 2013;1281:160-77.
105. Wenk GL. Neuropathologic changes in Alzheimer's disease. *The Journal of clinical psychiatry*. 2003;64 Suppl 9:7-10.
106. Lehr S HS, Lamers D, et al. . *Mol Cell Proteomics* 2012; 11(1): M111 010504. Identification and validation of novel adipokines released from primary human adipocytes.
107. Duvnjak L, Duvnjak M. The metabolic syndrome - an ongoing story. *J Physiol Pharmacol*. 2009;60 Suppl 7:19-24.
108. Bluher M. Adipose tissue dysfunction in obesity. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association*. 2009;117(6):241-50.
109. Abizaid A, Horvath TL. Brain circuits regulating energy homeostasis. *Regulatory peptides*. 2008;149(1-3):3-10.
110. Harvey J, Shanley LJ, O'Malley D, Irving AJ. Leptin: a potential cognitive enhancer? *Biochem Soc Trans*. 2005;33(Pt 5):1029-32.
111. Lieb W, Beiser AS, Vasan RS, Tan ZS, Au R, Harris TB, et al. Association of plasma leptin levels with incident Alzheimer disease and MRI measures of brain aging. *JAMA*. 2009;302(23):2565-72.
112. Harvey J. Leptin regulation of neuronal excitability and cognitive function. *Current opinion in pharmacology*. 2007;7(6):643-7.
113. Oomura Y, Hori N, Shiraishi T, Fukunaga K, Takeda H, Tsuji M, et al. Leptin facilitates learning and memory performance and enhances hippocampal CA1 long-term potentiation and CaMK II phosphorylation in rats. *Peptides*. 2006;27(11):2738-49.
114. Paz-Filho G, Wong ML, Licinio J. The procognitive effects of leptin in the brain and their clinical implications. *Int J Clin Pract*. 2010;64(13):1808-12.
115. Konstantinides S, Schafer K, Koschnick S, Loskutoff DJ. Leptin-dependent platelet aggregation and arterial thrombosis suggests a mechanism for atherothrombotic disease in obesity. *J Clin Invest*. 2001;108(10):1533-40.
116. Singhal A, Farooqi IS, Cole TJ, O'Rahilly S, Fewtrell M, Kattenhorn M, et al. Influence of leptin on arterial distensibility: a novel link between obesity and cardiovascular disease? *Circulation*. 2002;106(15):1919-24.

117. Van Gaal LF, Mertens IL, De Block CE. Mechanisms linking obesity with cardiovascular disease. *Nature*. 2006;444(7121):875-80.
118. Chandran M, Phillips SA, Ciaraldi T, Henry RR. Adiponectin: more than just another fat cell hormone? *Diabetes care*. 2003;26(8):2442-50.
119. Ouchi N, Ohishi M, Kihara S, Funahashi T, Nakamura T, Nagaretani H, et al. Association of hypoadiponectinemia with impaired vasoreactivity. *Hypertension*. 2003;42(3):231-4.
120. Jequier E. Leptin signaling, adiposity, and energy balance. *Ann N Y Acad Sci*. 2002;967:379-88.
121. Banks WA, DiPalma CR, Farrell CL. Impaired transport of leptin across the blood-brain barrier in obesity. *Peptides*. 1999;20(11):1341-5.
122. Caro JF, Kolaczynski JW, Nyce MR, Ohannesian JP, Opentanova I, Goldman WH, et al. Decreased cerebrospinal-fluid/serum leptin ratio in obesity: a possible mechanism for leptin resistance. *Lancet*. 1996;348(9021):159-61.
123. El-Haschimi K, Pierroz DD, Hileman SM, Bjorbaek C, Flier JS. Two defects contribute to hypothalamic leptin resistance in mice with diet-induced obesity. *J Clin Invest*. 2000;105(12):1827-32.
124. Friedman JM, Halaas JL. Leptin and the regulation of body weight in mammals. *Nature*. 1998;395(6704):763-70.
125. Hileman SM, Pierroz DD, Masuzaki H, Bjorbaek C, El-Haschimi K, Banks WA, et al. Characterization of short isoforms of the leptin receptor in rat cerebral microvessels and of brain uptake of leptin in mouse models of obesity. *Endocrinology*. 2002;143(3):775-83.
126. Wilsey J, Zolotukhin S, Prima V, Scarpace PJ. Central leptin gene therapy fails to overcome leptin resistance associated with diet-induced obesity. *Am J Physiol Regul Integr Comp Physiol*. 2003;285(5):R1011-20.
127. Yamashita T, Murakami T, Iida M, Kuwajima M, Shima K. Leptin receptor of Zucker fatty rat performs reduced signal transduction. *Diabetes*. 1997;46(6):1077-80.
128. Farr SA, Banks WA, Morley JE. Effects of leptin on memory processing. *Peptides*. 2006;27(6):1420-5.
129. Bouret SG. Neurodevelopmental actions of leptin. *Brain Res*. 2010;1350:2-9.
130. Morrison CD. Leptin signaling in brain: A link between nutrition and cognition? *Biochim Biophys Acta*. 2009;1792(5):401-8.
131. Zhang F, Wang S, Signore AP, Chen J. Neuroprotective effects of leptin against ischemic injury induced by oxygen-glucose deprivation and transient cerebral ischemia. *Stroke*. 2007;38(8):2329-36.
132. Fulton S, Pissios P, Manchon RP, Stiles L, Frank L, Pothos EN, et al. Leptin regulation of the mesoaccumbens dopamine pathway. *Neuron*. 2006;51(6):811-22.
133. Hayes MR, Skibicka KP, Lechner TM, Guarnieri DJ, DiLeone RJ, Bence KK, et al. Endogenous leptin signaling in the caudal nucleus tractus solitarius and area postrema is required for energy balance regulation. *Cell Metab*. 2010;11(1):77-83.

134. Hommel JD, Trinko R, Sears RM, Georgescu D, Liu ZW, Gao XB, et al. Leptin receptor signaling in midbrain dopamine neurons regulates feeding. *Neuron*. 2006;51(6):801-10.
135. Lu XY, Kim CS, Frazer A, Zhang W. Leptin: a potential novel antidepressant. *Proc Natl Acad Sci U S A*. 2006;103(5):1593-8.
136. Diez JJ, Iglesias P. The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur J Endocrinol*. 2003;148(3):293-300.
137. Ouchi N, Kihara S, Funahashi T, Matsuzawa Y, Walsh K. Obesity, adiponectin and vascular inflammatory disease. *Curr Opin Lipidol*. 2003;14(6):561-6.
138. Villarreal-Molina MT, Antuna-Puente B. Adiponectin: Anti-inflammatory and cardioprotective effects. *Biochimie*. 2012.
139. Qiu G, Wan R, Hu J, Mattson MP, Spangler E, Liu S, et al. Adiponectin protects rat hippocampal neurons against excitotoxicity. *Age (Dordr)*. 2011;33(2):155-65.
140. Qi Y, Takahashi N, Hileman SM, Patel HR, Berg AH, Pajvani UB, et al. Adiponectin acts in the brain to decrease body weight. *Nat Med*. 2004;10(5):524-9.
141. Gardes J, Bouhnik J, Clauser E, Corvol P, Menard J. Role of angiotensinogen in blood pressure homeostasis. *Hypertension*. 1982;4(2):185-9.
142. Kim HS, Kregge JH, Kluckman KD, Hagaman JR, Hodgins JB, Best CE, et al. Genetic control of blood pressure and the angiotensinogen locus. *Proc Natl Acad Sci U S A*. 1995;92(7):2735-9.
143. Ohkubo H, Kawakami H, Kakehi Y, Takumi T, Arai H, Yokota Y, et al. Generation of transgenic mice with elevated blood pressure by introduction of the rat renin and angiotensinogen genes. *Proc Natl Acad Sci U S A*. 1990;87(13):5153-7.
144. Walker WG, Whelton PK, Saito H, Russell RP, Hermann J. Relation between blood pressure and renin, renin substrate, angiotensin II, aldosterone and urinary sodium and potassium in 574 ambulatory subjects. *Hypertension*. 1979;1(3):287-91.
145. Watt GC, Harrap SB, Foy CJ, Holton DW, Edwards HV, Davidson HR, et al. Abnormalities of glucocorticoid metabolism and the renin-angiotensin system: a four-corners approach to the identification of genetic determinants of blood pressure. *J Hypertens*. 1992;10(5):473-82.
146. Yiannikouris F, Karounos M, Charnigo R, English VL, Rateri DL, Daugherty A, et al. Adipocyte-specific deficiency of angiotensinogen decreases plasma angiotensinogen concentration and systolic blood pressure in mice. *Am J Physiol Regul Integr Comp Physiol*. 2012;302(2):R244-51.
147. Banks WA. Denial versus dualism: the blood-brain barrier as an interface of the gut-brain axis. *Endocrinology*. 2006;147(6):2609-10.
148. Pan W, Kastin AJ. Adipokines and the blood-brain barrier. *Peptides*. 2007;28(6):1317-30.
149. Yanai K, Saito T, Kakinuma Y, Kon Y, Hirota K, Taniguchi-Yanai K, et al. Renin-dependent cardiovascular functions and renin-independent blood-brain barrier functions revealed by renin-deficient mice. *J Biol Chem*. 2000;275(1):5-8.

150. Youssef AM, Hamidian Jahromi A, Vijay CG, Granger DN, Alexander JS. Intra-abdominal hypertension causes reversible blood-brain barrier disruption. *The journal of trauma and acute care surgery*. 2012;72(1):183-8.
151. Allen AM, Dampney RA, Mendelsohn FA. Angiotensin receptor binding and pressor effects in cat subretrofacial nucleus. *Am J Physiol*. 1988;255(5 Pt 2):H1011-7.
152. Andreatta SH, Averill DB, Santos RA, Ferrario CM. The ventrolateral medulla. A new site of action of the renin-angiotensin system. *Hypertension*. 1988;11(2 Pt 2):I163-6.
153. Averill DB, Diz DI, Barnes KL, Ferrario CM. Pressor responses of angiotensin II microinjected into the dorsomedial medulla of the dog. *Brain Res*. 1987;414(2):294-300.
154. Jensen LL, Harding JW, Wright JW. Role of paraventricular nucleus in control of blood pressure and drinking in rats. *Am J Physiol*. 1992;262(6 Pt 2):F1068-75.
155. McKinley MJ, Albiston AL, Allen AM, Mathai ML, May CN, McAllen RM, et al. The brain renin-angiotensin system: location and physiological roles. *Int J Biochem Cell Biol*. 2003;35(6):901-18.
156. Simpson JB. The circumventricular organs and the central actions of angiotensin. *Neuroendocrinology*. 1981;32(4):248-56.
157. Thornton SN, Nicolaidis S. Blood pressure effects of iontophoretically applied bioactive hormones in the anterior forebrain of the rat. *Am J Physiol*. 1993;265(4 Pt 2):R826-33.
158. Guerre-Millo M. Adipose tissue and adipokines: for better or worse. *Diabetes Metab*. 2004;30(1):13-9.
159. Soeda S, Koyanagi S, Kuramoto Y, Kimura M, Oda M, Kozako T, et al. Anti-apoptotic roles of plasminogen activator inhibitor-1 as a neurotrophic factor in the central nervous system. *Thromb Haemost*. 2008;100(6):1014-20.
160. Soeda S, Oda M, Ochiai T, Shimeno H. Deficient release of plasminogen activator inhibitor-1 from astrocytes triggers apoptosis in neuronal cells. *Brain Res Mol Brain Res*. 2001;91(1-2):96-103.
161. Jeon H, Kim JH, Kim JH, Lee WH, Lee MS, Suk K. Plasminogen activator inhibitor type 1 regulates microglial motility and phagocytic activity. *J Neuroinflammation*. 2012;9:149.
162. Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nature reviews Neuroscience*. 2007;8(1):57-69.
163. Garden GA, Moller T. Microglia biology in health and disease. *Journal of neuroimmune pharmacology : the official journal of the Society on NeuroImmune Pharmacology*. 2006;1(2):127-37.
164. Graeber MB, Streit WJ. Microglia: biology and pathology. *Acta Neuropathol*. 2010;119(1):89-105.
165. Mallat M, Marin-Teva JL, Cheret C. Phagocytosis in the developing CNS: more than clearing the corpses. *Current opinion in neurobiology*. 2005;15(1):101-7.

166. Woo MS, Park JS, Choi IY, Kim WK, Kim HS. Inhibition of MMP-3 or -9 suppresses lipopolysaccharide-induced expression of proinflammatory cytokines and iNOS in microglia. *J Neurochem*. 2008;106(2):770-80.
167. Miskin R, Masos T. Transgenic mice overexpressing urokinase-type plasminogen activator in the brain exhibit reduced food consumption, body weight and size, and increased longevity. *J Gerontol A Biol Sci Med Sci*. 1997;52(2):B118-24.
168. Rocha VZ, Folco EJ. Inflammatory concepts of obesity. *Int J Inflam*. 2011;2011:529061.
169. Samuelsson AM, Jennische E, Hansson HA, Holmang A. Prenatal exposure to interleukin-6 results in inflammatory neurodegeneration in hippocampus with NMDA/GABA(A) dysregulation and impaired spatial learning. *Am J Physiol Regul Integr Comp Physiol*. 2006;290(5):R1345-56.
170. Balschun D, Wetzel W, Del Rey A, Pitossi F, Schneider H, Zuschratter W, et al. Interleukin-6: a cytokine to forget. *FASEB J*. 2004;18(14):1788-90.
171. Gibertini M, Newton C, Friedman H, Klein TW. Spatial learning impairment in mice infected with *Legionella pneumophila* or administered exogenous interleukin-1-beta. *Brain Behav Immun*. 1995;9(2):113-28.
172. Monje ML, Toda H, Palmer TD. Inflammatory blockade restores adult hippocampal neurogenesis. *Science*. 2003;302(5651):1760-5.
173. Poluektova L, Meyer V, Walters L, Paez X, Gendelman HE. Macrophage-induced inflammation affects hippocampal plasticity and neuronal development in a murine model of HIV-1 encephalitis. *Glia*. 2005;52(4):344-53.
174. Wei H, Zou H, Sheikh AM, Malik M, Dobkin C, Brown WT, et al. IL-6 is increased in the cerebellum of autistic brain and alters neural cell adhesion, migration and synaptic formation. *J Neuroinflammation*. 2011;8:52.
175. Wei H, Chadman KK, McCloskey DP, Sheikh AM, Malik M, Brown WT, et al. Brain IL-6 elevation causes neuronal circuitry imbalances and mediates autism-like behaviors. *Biochim Biophys Acta*. 2012;1822(6):831-42.
176. Milanski M, Degasperi G, Coope A, Morari J, Denis R, Cintra DE, et al. Saturated fatty acids produce an inflammatory response predominantly through the activation of TLR4 signaling in hypothalamus: implications for the pathogenesis of obesity. *J Neurosci*. 2009;29(2):359-70.
177. Ropelle ER, Flores MB, Cintra DE, Rocha GZ, Pauli JR, Morari J, et al. IL-6 and IL-10 anti-inflammatory activity links exercise to hypothalamic insulin and leptin sensitivity through IKKbeta and ER stress inhibition. *PLoS biology*. 2010;8(8).
178. Barger SW, Horster D, Furukawa K, Goodman Y, Kriegstein J, Mattson MP. Tumor necrosis factors alpha and beta protect neurons against amyloid beta-peptide toxicity: evidence for involvement of a kappa B-binding factor and attenuation of peroxide and Ca²⁺ accumulation. *Proc Natl Acad Sci U S A*. 1995;92(20):9328-32.

179. Beattie EC, Stellwagen D, Morishita W, Bresnahan JC, Ha BK, Von Zastrow M, et al. Control of synaptic strength by glial TNF α . *Science*. 2002;295(5563):2282-5.
180. Cacci E, Claassen JH, Kokaia Z. Microglia-derived tumor necrosis factor- α exaggerates death of newborn hippocampal progenitor cells in vitro. *J Neurosci Res*. 2005;80(6):789-97.
181. Pickering M, Cumiskey D, O'Connor JJ. Actions of TNF- α on glutamatergic synaptic transmission in the central nervous system. *Experimental physiology*. 2005;90(5):663-70.
182. Butler MP, O'Connor JJ, Moynagh PN. Dissection of tumor-necrosis factor- α inhibition of long-term potentiation (LTP) reveals a p38 mitogen-activated protein kinase-dependent mechanism which maps to early-but not late-phase LTP. *Neuroscience*. 2004;124(2):319-26.
183. Tancredi V, D'Arcangelo G, Grassi F, Tarroni P, Palmieri G, Santoni A, et al. Tumor necrosis factor alters synaptic transmission in rat hippocampal slices. *Neurosci Lett*. 1992;146(2):176-8.
184. Romanatto T, Cesquini M, Amaral ME, Roman EA, Moraes JC, Torsoni MA, et al. TNF- α acts in the hypothalamus inhibiting food intake and increasing the respiratory quotient--effects on leptin and insulin signaling pathways. *Peptides*. 2007;28(5):1050-8.
185. Selmaj K, Raine CS. Tumor necrosis factor mediates myelin damage in organotypic cultures of nervous tissue. *Ann N Y Acad Sci*. 1988;540:568-70.
186. Marsland AL, Gianaros PJ, Abramowitch SM, Manuck SB, Hariri AR. Interleukin-6 covaries inversely with hippocampal grey matter volume in middle-aged adults. *Biol Psychiatry*. 2008;64(6):484-90.
187. Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation*. 2002;105(9):1135-43.
188. Zadelaar S, Kleemann R, Verschuren L, de Vries-Van der Weij J, van der Hoorn J, Princen HM, et al. Mouse models for atherosclerosis and pharmaceutical modifiers. *Arterioscler Thromb Vasc Biol*. 2007;27(8):1706-21.
189. Brown MS, Goldstein JL. A receptor-mediated pathway for cholesterol homeostasis. *Science*. 1986;232(4746):34-47.
190. Ma Y, Wang W, Zhang J, Lu Y, Wu W, Yan H, et al. Hyperlipidemia and atherosclerotic lesion development in Ldlr-deficient mice on a long-term high-fat diet. *PLoS One*. 2012;7(4):e35835.
191. Radonjic M, Wielinga PY, Wopereis S, Kelder T, Goelela VS, Verschuren L, et al. Differential effects of drug interventions and dietary lifestyle in developing type 2 diabetes and complications: a systems biology analysis in LDLr $^{-/-}$ mice. *PLoS One*. 2013;8(2):e56122.
192. Volkow ND, Wang GJ, Telang F, Fowler JS, Goldstein RZ, Alia-Klein N, et al. Inverse association between BMI and prefrontal metabolic activity in healthy adults. *Obesity (Silver Spring)*. 2009;17(1):60-5.

193. Olde Dubbelink KT, Feliuss A, Verbunt JP, van Dijk BW, Berendse HW, Stam CJ, et al. Increased resting-state functional connectivity in obese adolescents; a magnetoencephalographic pilot study. *PLoS One*. 2008;3(7):e2827.
194. Yau PL, Javier DC, Ryan CM, Tsui WH, Ardekani BA, Ten S, et al. Preliminary evidence for brain complications in obese adolescents with type 2 diabetes mellitus. *Diabetologia*. 2010;53(11):2298-306.
195. Sawyer MG, Harchak T, Wake M, Lynch J. Four-year prospective study of BMI and mental health problems in young children. *Pediatrics*. 2011;128(4):677-84.
196. Tascilar ME, Turkkahraman D, Oz O, Yucel M, Taskesen M, Eker I, et al. P300 auditory event-related potentials in children with obesity: is childhood obesity related to impairment in cognitive functions? *Pediatric diabetes*. 2011;12(7):589-95.
197. Alosco ML, Stanek KM, Galioto R, Korgaonkar MS, Grieve SM, Brickman AM, et al. Body mass index and brain structure in healthy children and adolescents. *Int J Neurosci*. 2014;124(1):49-55.
198. Boitard C, Etchamendy N, Sauvant J, Aubert A, Tronel S, Marighetto A, et al. Juvenile, but not adult exposure to high-fat diet impairs relational memory and hippocampal neurogenesis in mice. *Hippocampus*. 2012;22(11):2095-100.
199. Park HR, Park M, Choi J, Park KY, Chung HY, Lee J. A high-fat diet impairs neurogenesis: involvement of lipid peroxidation and brain-derived neurotrophic factor. *Neurosci Lett*. 2010;482(3):235-9.
200. Wang X, Ge A, Cheng M, Guo F, Zhao M, Zhou X, et al. Increased hypothalamic inflammation associated with the susceptibility to obesity in rats exposed to high-fat diet. *Exp Diabetes Res*. 2012;2012:847246.
201. Gonzalez-Periz A, Horrillo R, Ferre N, Gronert K, Dong B, Moran-Salvador E, et al. Obesity-induced insulin resistance and hepatic steatosis are alleviated by omega-3 fatty acids: a role for resolvins and protectins. *FASEB J*. 2009;23(6):1946-57.
202. Wielinga PY, Harthoorn LF, Verschuren L, Schoemaker MH, Jouni ZE, van Tol EA, et al. Arachidonic acid/docosahexaenoic acid-supplemented diet in early life reduces body weight gain, plasma lipids, and adiposity in later life in ApoE*3Leiden mice. *Mol Nutr Food Res*. 2012;56(7):1081-9.
203. Weisinger HS, Armitage JA, Sinclair AJ, Vingrys AJ, Burns PL, Weisinger RS. Perinatal omega-3 fatty acid deficiency affects blood pressure later in life. *Nat Med*. 2001;7(3):258-9.
204. Brenna JT, Diau GY. The influence of dietary docosahexaenoic acid and arachidonic acid on central nervous system polyunsaturated fatty acid composition. *Prostaglandins, leukotrienes, and essential fatty acids*. 2007;77(5-6):247-50.
205. Davis-Bruno K, Tassinari MS. Essential fatty acid supplementation of DHA and ARA and effects on neurodevelopment across animal species: a review of the literature. *Birth defects research Part B, Developmental and reproductive toxicology*. 2011;92(3):240-50.

206. Kleemann R, van Erk M, Verschuren L, van den Hoek AM, Koek M, Wielinga PY, et al. Time-resolved and tissue-specific systems analysis of the pathogenesis of insulin resistance. *PLoS One*. 2010;5(1):e8817.
207. Jong MC, Hofker MH, Havekes LM. Role of ApoCs in lipoprotein metabolism: functional differences between ApoC1, ApoC2, and ApoC3. *Arterioscler Thromb Vasc Biol*. 1999;19(3):472-84.
208. van Vlijmen BJ, van den Maagdenberg AM, Gijbels MJ, van der Boom H, HogenEsch H, Frants RR, et al. Diet-induced hyperlipoproteinemia and atherosclerosis in apolipoprotein E3-Leiden transgenic mice. *J Clin Invest*. 1994;93(4):1403-10.
209. Kleemann R, Verschuren L, van Erk MJ, Nikolsky Y, Cnubben NH, Verheij ER, et al. Atherosclerosis and liver inflammation induced by increased dietary cholesterol intake: a combined transcriptomics and metabolomics analysis. *Genome biology*. 2007;8(9):R200.
210. Madsen L, Pedersen LM, Liaset B, Ma T, Petersen RK, van den Berg S, et al. cAMP-dependent signaling regulates the adipogenic effect of n-6 polyunsaturated fatty acids. *J Biol Chem*. 2008;283(11):7196-205.
211. Hooijmans CR, Van der Zee CE, Dederen PJ, Brouwer KM, Reijmer YD, van Groen T, et al. DHA and cholesterol containing diets influence Alzheimer-like pathology, cognition and cerebral vasculature in APP^{swe}/PS1^{dE9} mice. *Neurobiol Dis*. 2009;33(3):482-98.
212. Streijger F, Oerlemans F, Ellenbroek BA, Jost CR, Wieringa B, Van der Zee CE. Structural and behavioural consequences of double deficiency for creatine kinases BCK and UbCKmit. *Behav Brain Res*. 2005;157(2):219-34.
213. Janssen CI, Zerbi V, Mutsaers MP, de Jong BS, Wiesmann M, Arnoldussen IA, et al. Impact of dietary n-3 polyunsaturated fatty acids on cognition, motor skills and hippocampal neurogenesis in developing C57BL/6J mice. *J Nutr Biochem*. 2015;26(1):24-35.
214. Bevins RA, Besheer J. Object recognition in rats and mice: a one-trial non-matching-to-sample learning task to study 'recognition memory'. *Nat Protoc*. 2006;1(3):1306-11.
215. Wiesmann M, Jansen D, Zerbi V, Broersen LM, Garthe A, Kiliaan AJ. Improved spatial learning strategy and memory in aged Alzheimer A β PP^{swe}/PS1^{dE9} mice on a multi-nutrient diet. *Journal of Alzheimer's Disease*. 2013;37(1):233-45.
216. Garthe A, Behr J, Kempermann G. Adult-generated hippocampal neurons allow the flexible use of spatially precise learning strategies. *PLoS One*. 2009;4(5):e5464.
217. Aerni-Flessner L, Abi-Jaoude M, Koenig A, Payne M, Hruz PW. GLUT4, GLUT1, and GLUT8 are the dominant GLUT transcripts expressed in the murine left ventricle. *Cardiovascular diabetology*. 2012;11:63.
218. Pleau JM, Esling A, Geutkens S, Dardenne M, Homo-Delarche F. Pancreatic hormone and glutamic acid decarboxylase expression in the mouse thymus: a real-time PCR study. *Biochem Biophys Res Commun*. 2001;283(4):843-8.

219. Lofqvist C, Willett KL, Aspegren O, Smith AC, Aderman CM, Connor KM, et al. Quantification and localization of the IGF/insulin system expression in retinal blood vessels and neurons during oxygen-induced retinopathy in mice. *Investigative ophthalmology & visual science*. 2009;50(4):1831-7.
220. Fruhbeck G, Gomez-Ambrosi J, Martinez JA. Pre- and postprandial expression of the leptin receptor splice variants OB-Ra and OB-Rb in murine peripheral tissues. *Physiol Res*. 1999;48(3):189-95.
221. Schroyen B, Guimaraes EL, Dolle L, Coulon S, Empsen C, Nyssen M, et al. Leptin-mediated reactive oxygen species production does not significantly affect primary mouse hepatocyte functions in vitro. *Eur J Gastroenterol Hepatol*. 2012;24(12):1370-80.
222. Hellstrom A, Perruzzi C, Ju M, Engstrom E, Hard AL, Liu JL, et al. Low IGF-I suppresses VEGF-survival signaling in retinal endothelial cells: direct correlation with clinical retinopathy of prematurity. *Proc Natl Acad Sci U S A*. 2001;98(10):5804-8.
223. Noth R, Lange-Grumfeld J, Stuber E, Kruse ML, Ellrichmann M, Hasler R, et al. Increased intestinal permeability and tight junction disruption by altered expression and localization of occludin in a murine graft versus host disease model. *BMC gastroenterology*. 2011;11:109.
224. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008;3(6):1101-8.
225. Zerbi V, Jansen D, Wiesmann M, Fang X, Broersen LM, Veltien A, et al. Multinutrient diets improve cerebral perfusion and neuroprotection in a murine model of Alzheimer's disease. *Neurobiol Aging*. 2014;35(3):600-13.
226. Zerbi V, Wiesmann M, Emmerzaal TL, Jansen D, Van Beek M, Mutsaers MP, et al. Resting-State Functional Connectivity Changes in Aging apoE4 and apoE-KO Mice. *J Neurosci*. 2014;34(42):13963-75.
227. Zerbi V, Jansen D, Dederen PJ, Veltien A, Hamans B, Liu Y, et al. Microvascular cerebral blood volume changes in aging APPswe/PS1dE9 AD mouse model: a voxel-wise approach. *Brain Structure and Function*. 2012:1-14.
228. Kim SG. Quantification of relative cerebral blood flow change by flow-sensitive alternating inversion recovery (FAIR) technique: application to functional mapping. *Magn Reson Med*. 1995;34(3):293-301.
229. Kwong KK, Chesler DA, Weisskoff RM, Donahue KM, Davis TL, Ostergaard L, et al. MR perfusion studies with T1-weighted echo planar imaging. *Magn Reson Med*. 1995;34(6):878-87.
230. Zerbi V, Kleinnijenhuis M, Fang X, Jansen D, Veltien A, Van Asten J, et al. Gray and white matter degeneration revealed by diffusion in an Alzheimer mouse model. *Neurobiology of aging*. 2012.
231. Shapiro SS, Wilk, M.B. An analysis of variance test for normality (complete samples). *Biometrika*. 1965;52:591-611.

232. Snedecor GW, Cochran WG. Statistical Methods. Iowa State University Presse Med. 1989.
233. Ozanne SE, Lewis R, Jennings BJ, Hales CN. Early programming of weight gain in mice prevents the induction of obesity by a highly palatable diet. *Clin Sci (Lond)*. 2004;106(2):141-5.
234. Winick M, Noble A. Cellular response in rats during malnutrition at various ages. *J Nutr*. 1966;89(3):300-6.
235. Fernandez-Twinn D, Ozanne S. Mechanisms by which poor early growth programs type-2 diabetes, obesity and the metabolic syndrome. *Physiology & Behavior*. 2006;88(3):234-43.
236. Fernandez-Twinn DS, Ozanne SE. Early life nutrition and metabolic programming. *Ann Ny Acad Sci*. 2010;1212(1):78-96.
237. Pacce S, Saure C, Mazza CS, Garcia S, Tomzig RG, Lopez AP, et al. Impact of maternal nutritional status before and during pregnancy on neonatal body composition: A cross-sectional study. *Diabetes & metabolic syndrome*. 2015.
238. Greiner RS, Moriguchi T, Hutton A, Slotnick BM, Salem N, Jr. Rats with low levels of brain docosahexaenoic acid show impaired performance in olfactory-based and spatial learning tasks. *Lipids*. 1999;34 Suppl:S239-43.
239. Ikemoto A, Nitta A, Furukawa S, Ohishi M, Nakamura A, Fujii Y, et al. Dietary n-3 fatty acid deficiency decreases nerve growth factor content in rat hippocampus. *Neurosci Lett*. 2000;285(2):99-102.
240. Kodas E, Vancassel S, Lejeune B, Guilloteau D, Chalon S. Reversibility of n-3 fatty acid deficiency-induced changes in dopaminergic neurotransmission in rats: critical role of developmental stage. *J Lipid Res*. 2002;43(8):1209-19.
241. Li D, Weisinger HS, Weisinger RS, Mathai M, Armitage JA, Vingrys AJ, et al. Omega 6 to omega 3 fatty acid imbalance early in life leads to persistent reductions in DHA levels in glycerophospholipids in rat hypothalamus even after long-term omega 3 fatty acid repletion. *Prostaglandins, leukotrienes, and essential fatty acids*. 2006;74(6):391-9.
242. Litman BJ, Niu SL, Polozova A, Mitchell DC. The role of docosahexaenoic acid containing phospholipids in modulating G protein-coupled signaling pathways: visual transduction. *J Mol Neurosci*. 2001;16(2-3):237-42; discussion 79-84.
243. Pifferi F, Roux F, Langelier B, Alessandri JM, Vancassel S, Jouin M, et al. (n-3) polyunsaturated fatty acid deficiency reduces the expression of both isoforms of the brain glucose transporter GLUT1 in rats. *J Nutr*. 2005;135(9):2241-6.
244. Weisinger HS, Vingrys AJ, Bui BV, Sinclair AJ. Effects of dietary n-3 fatty acid deficiency and repletion in the guinea pig retina. *Investigative ophthalmology & visual science*. 1999;40(2):327-38.
245. Xiao Y, Wang L, Xu RJ, Chen ZY. DHA depletion in rat brain is associated with impairment on spatial learning and memory. *Biomedical and environmental sciences : BES*. 2006;19(6):474-80.

246. Gomez-Pinilla F, Ying Z. Differential effects of exercise and dietary docosahexaenoic acid on molecular systems associated with control of allostasis in the hypothalamus and hippocampus. *Neuroscience*. 2010;168(1):130-7.
247. McMillen IC, Adam CL, Muhlhauser BS. Early origins of obesity: programming the appetite regulatory system. *J Physiol*. 2005;565(Pt 1):9-17.
248. Lizarbe B, Benitez A, Pelaez Brioso GA, Sanchez-Montanes M, Lopez-Larrubia P, Ballesteros P, et al. Hypothalamic metabolic compartmentation during appetite regulation as revealed by magnetic resonance imaging and spectroscopy methods. *Frontiers in neuroenergetics*. 2013;5:6.
249. Tang-Christensen M, Vrang N, Ortmann S, Bidlingmaier M, Horvath TL, Tschöp M. Central administration of ghrelin and agouti-related protein (83-132) increases food intake and decreases spontaneous locomotor activity in rats. *Endocrinology*. 2004;145(10):4645-52.
250. Bernardini R, Chiarenza A, Calogero AE, Gold PW, Chrousos GP. Arachidonic acid metabolites modulate rat hypothalamic corticotropin-releasing hormone secretion in vitro. *Neuroendocrinology*. 1989;50(6):708-15.
251. Davidson TL, Kanoski SE, Schier LA, Clegg DJ, Benoit SC. A potential role for the hippocampus in energy intake and body weight regulation. *Current opinion in pharmacology*. 2007;7(6):613-6.
252. Fehm HL, Born J, Peters A. Glucocorticoids and melanocortins in the regulation of body weight in humans. *Horm Metab Res*. 2004;36(6):360-4.
253. Peters A, Schweiger U, Pellerin L, Hubold C, Oltmanns KM, Conrad M, et al. The selfish brain: competition for energy resources. *Neurosci Biobehav Rev*. 2004;28(2):143-80.
254. Rubio-Aliaga I, Roos B, Sailer M, McLoughlin GA, Boekschoten MV, van Erk M, et al. Alterations in hepatic one-carbon metabolism and related pathways following a high-fat dietary intervention. *Physiol Genomics*. 2011;43(8):408-16.
255. Mells JE, Fu PP, Sharma S, Olson D, Cheng L, Handy JA, et al. Glp-1 analog, liraglutide, ameliorates hepatic steatosis and cardiac hypertrophy in C57BL/6J mice fed a Western diet. *Am J Physiol Gastrointest Liver Physiol*. 2012;302(2):G225-35.
256. Carmiel-Haggai M, Cederbaum AI, Nieto N. A high-fat diet leads to the progression of non-alcoholic fatty liver disease in obese rats. *FASEB J*. 2005;19(1):136-8.
257. Machado MV, Michelotti GA, Xie G, Almeida Pereira T, Boursier J, Bohnic B, et al. Mouse models of diet-induced nonalcoholic steatohepatitis reproduce the heterogeneity of the human disease. *PLoS One*. 2015;10(5):e0127991.
258. Scorletti E, Byrne CD. Omega-3 fatty acids, hepatic lipid metabolism, and nonalcoholic fatty liver disease. *Annual review of nutrition*. 2013;33:231-48.
259. West DB, Boozer CN, Moody DL, Atkinson RL. Dietary obesity in nine inbred mouse strains. *Am J Physiol*. 1992;262(6 Pt 2):R1025-32.
260. Liang W, Tonini G, Mulder P, Kelder T, van Erk M, van den Hoek AM, et al. Coordinated and interactive expression of genes of lipid metabolism and inflammation in adipose tissue and liver during metabolic overload. 2013.

261. Mulder P, Morrison M, Wielinga P, van Duyvenvoorde W, Kooistra T, Kleemann R. Surgical removal of inflamed epididymal white adipose tissue attenuates the development of non-alcoholic steatohepatitis in obesity. *International journal of obesity* (2005). 2015.
262. Gesta S, Tseng YH, Kahn CR. Developmental origin of fat: tracking obesity to its source. *Cell*. 2007;131(2):242-56.
263. Tsukada H, Kakiuchi T, Fukumoto D, Nishiyama S, Koga K. Docosahexaenoic acid (DHA) improves the age-related impairment of the coupling mechanism between neuronal activation and functional cerebral blood flow response: a PET study in conscious monkeys. *Brain Res*. 2000;862(1-2):180-6.
264. Dorrance AM, Matin N, Pires PW. The effects of obesity on the cerebral vasculature. *Current vascular pharmacology*. 2014;12(3):462-72.
265. Selim M, Jones R, Novak P, Zhao P, Novak V. The effects of body mass index on cerebral blood flow velocity. *Clin Auton Res*. 2008;18(6):331-8.
266. Willeumier KC, Taylor DV, Amen DG. Elevated BMI is associated with decreased blood flow in the prefrontal cortex using SPECT imaging in healthy adults. *Obesity* (Silver Spring). 2011;19(5):1095-7.
267. Deng J, Zhang J, Feng C, Xiong L, Zuo Z. Critical role of matrix metalloprotease-9 in chronic high fat diet-induced cerebral vascular remodelling and increase of ischaemic brain injury in micedagger. *Cardiovasc Res*. 2014.
268. Deutsch C, Portik-Dobos V, Smith AD, Ergul A, Dorrance AM. Diet-induced obesity causes cerebral vessel remodeling and increases the damage caused by ischemic stroke. *Microvasc Res*. 2009;78(1):100-6.
269. Lynch CM, Kinzenbaw DA, Chen X, Zhan S, Mezzetti E, Filosa J, et al. Nox2-derived superoxide contributes to cerebral vascular dysfunction in diet-induced obesity. *Stroke*. 2013;44(11):3195-201.
270. Lim S, Suzuki H. Changes in maze behavior of mice occur after sufficient accumulation of docosahexaenoic acid in brain. *J Nutr*. 2001;131(2):319-24.
271. Grayson DS, Kroenke CD, Neuringer M, Fair DA. Dietary omega-3 fatty acids modulate large-scale systems organization in the rhesus macaque brain. *J Neurosci*. 2014;34(6):2065-74.
272. Beilharz JE, Maniam J, Morris MJ. Short exposure to a diet rich in both fat and sugar or sugar alone impairs place, but not object recognition memory in rats. *Brain Behav Immun*. 2014;37:134-41.
273. Lavin DN, Joesting JJ, Chiu GS, Moon ML, Meng J, Dilger RN, et al. Fasting induces an anti-inflammatory effect on the neuroimmune system which a high-fat diet prevents. *Obesity* (Silver Spring). 2011;19(8):1586-94.
274. Boitard C, Cavaroc A, Sauviant J, Aubert A, Castanon N, Laye S, et al. Impairment of hippocampal-dependent memory induced by juvenile high-fat diet intake is associated with enhanced hippocampal inflammation in rats. *Brain Behav Immun*. 2014;40:9-17.

275. Yokum S, Ng J, Stice E. Relation of regional gray and white matter volumes to current BMI and future increases in BMI: a prospective MRI study. *Int J Obes (Lond)*. 2012;36(5):656-64.
276. McNamara RK, Able J, Jandacek R, Rider T, Tso P, Eliassen JC, et al. Docosahexaenoic acid supplementation increases prefrontal cortex activation during sustained attention in healthy boys: a placebo-controlled, dose-ranging, functional magnetic resonance imaging study. *Am J Clin Nutr*. 2010;91(4):1060-7.
277. Alexander AL, Lee JE, Lazar M, Field AS. Diffusion tensor imaging of the brain. *Neurotherapeutics*. 2007;4(3):316-29.
278. Kullmann S, Heni M, Veit R, Ketterer C, Schick F, Haring HU, et al. The obese brain: association of body mass index and insulin sensitivity with resting state network functional connectivity. *Hum Brain Mapp*. 2012;33(5):1052-61.
279. Kringelbach ML, de Araujo IE, Rolls ET. Taste-related activity in the human dorsolateral prefrontal cortex. *Neuroimage*. 2004;21(2):781-8.
280. Perez LM, Pareja-Galeano H, Sanchis-Gomar F, Emanuele E, Lucia A, Galvez BG. 'Adipaging': Aging and obesity share biological hallmarks related to a dysfunctional adipose tissue. *J Physiol*. 2016.
281. Morrison CD, Pistell PJ, Ingram DK, Johnson WD, Liu Y, Fernandez-Kim SO, et al. High fat diet increases hippocampal oxidative stress and cognitive impairment in aged mice: implications for decreased Nrf2 signaling. *J Neurochem*. 2010;114(6):1581-9.
282. Tucsek Z, Toth P, Tarantini S, Sosnowska D, Gautam T, Warrington JP, et al. Aging exacerbates obesity-induced cerebrovascular rarefaction, neurovascular uncoupling, and cognitive decline in mice. *J Gerontol A Biol Sci Med Sci*. 2014;69(11):1339-52.
283. Govindarajan N, Agis-Balboa RC, Walter J, Sananbenesi F, Fischer A. Sodium butyrate improves memory function in an Alzheimer's disease mouse model when administered at an advanced stage of disease progression. *J Alzheimers Dis*. 2011;26(1):187-97.
284. Liang W, Menke AL, Driessen A, Koek GH, Lindeman JH, Stoop R, et al. Establishment of a general NAFLD scoring system for rodent models and comparison to human liver pathology. *PLoS One*. 2014;9(12):e115922.
285. Morrison M, Mulder P, Salic K, Verheij J, Liang W, van Duyvenvoorde W, et al. Intervention with a caspase-1 inhibitor reduces obesity-associated hyperinsulinemia, non-alcoholic steatohepatitis and hepatic fibrosis in LDLR^{-/-}. Leiden mice. *International Journal of Obesity*. 2016.
286. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *PLoS biology*. 2010;8(6):e1000412.
287. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences*. 2011;108(Supplement 1):4516-22.

288. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and environmental microbiology*. 2009;75(23):7537-41.
289. Eren AM, Morrison HG, Lescault PJ, Reveillaud J, Vineis JH, Sogin ML. Minimum entropy decomposition: Unsupervised oligotyping for sensitive partitioning of high-throughput marker gene sequences. *The ISME journal*. 2015;9(4):968-79.
290. Hamady M, Lozupone C, Knight R. Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *The ISME journal*. 2010;4(1):17-27.
291. White JR, Nagarajan N, Pop M. Statistical methods for detecting differentially abundant features in clinical metagenomic samples. *PLoS computational biology*. 2009;5(4):e1000352.
292. Arnoldussen IA, Zerbi V, Wiesmann M, Noordman RH, Bolijn S, Mutsaers MP, et al. Early intake of long-chain polyunsaturated fatty acids preserves brain structure and function in diet-induced obesity. *J Nutr Biochem*. 2016;30:177-88.
293. Mulder P, Morrison MC, Wielinga PY, van Duyvenvoorde W, Kooistra T, Kleemann R. Surgical removal of inflamed epididymal white adipose tissue attenuates the development of non-alcoholic steatohepatitis in obesity. *Int J Obes (Lond)*. 2015.
294. Van Zutphen LFMB, V.; Ohl, F. *Handboek Proefdierkunde*. 5th edition ed. Maarssen: Elsevier Gezondheidszorg; 2009.
295. Jakobsdottir G, Xu J, Molin G, Ahrne S, Nyman M. High-fat diet reduces the formation of butyrate, but increases succinate, inflammation, liver fat and cholesterol in rats, while dietary fibre counteracts these effects. *PLoS One*. 2013;8(11):e80476.
296. Bocarsly ME, Fasolino M, Kane GA, LaMarca EA, Kirschen GW, Karatsoreos IN, et al. Obesity diminishes synaptic markers, alters microglial morphology, and impairs cognitive function. *Proc Natl Acad Sci U S A*. 2015;112(51):15731-6.
297. Marcil V, Delvin E, Seidman E, Poitras L, Zoltowska M, Garofalo C, et al. Modulation of lipid synthesis, apolipoprotein biogenesis, and lipoprotein assembly by butyrate. *Am J Physiol Gastrointest Liver Physiol*. 2002;283(2):G340-6.
298. den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud D-J, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Journal of Lipid Research*. 2013;54(9):2325-40.
299. den Besten G, Bleeker A, Gerding A, van Eunen K, Havinga R, van Dijk TH, et al. Short-Chain Fatty Acids Protect Against High-Fat Diet-Induced Obesity via a PPAR γ -Dependent Switch From Lipogenesis to Fat Oxidation. *Diabetes*. 2015;64(7):2398.
300. Kilgore M, Miller CA, Fass DM, Hennig KM, Haggarty SJ, Sweatt JD, et al. Inhibitors of class 1 histone deacetylases reverse contextual memory deficits in a mouse model of Alzheimer's disease. *Neuropsychopharmacology*. 2010;35(4):870-80.
301. Krauss RM, Winston M, Fletcher BJ, Grundy SM. Obesity: impact on cardiovascular disease. *Circulation*. 1998;98(14):1472-6.

302. Weisbrod RM, Shiang T, Al Sayah L, Fry JL, Bajpai S, Reinhart-King CA, et al. Arterial stiffening precedes systolic hypertension in diet-induced obesity. *Hypertension*. 2013;62(6):1105-10.
303. Bink DI, Ritz K, Aronica E, van der Weerd L, Daemen MJ. Mouse models to study the effect of cardiovascular risk factors on brain structure and cognition. *J Cereb Blood Flow Metab*. 2013;33(11):1666-84.
304. Liu D, Andrade SP, Castro PR, Treacy J, Ashworth J, Slevin M. Low Concentration of Sodium Butyrate from Ultrabraid+NaBu suture, Promotes Angiogenesis and Tissue Remodelling in Tendon-bones Injury. *Scientific reports*. 2016;6:34649.
305. Verschuren L, Kooistra T, Bernhagen J, Voshol PJ, Ouwers DM, van Erk M, et al. MIF-deficiency reduces chronic inflammation in white adipose tissue and impairs the development of insulin resistance, glucose intolerance and associated atherosclerotic disease. *Circulation research*. 2009;105(1):99-107.
306. Merat S, Fruebis J, Sutphin M, Silvestre M, Reaven PD. Effect of aging on aortic expression of the vascular cell adhesion molecule-1 and atherosclerosis in murine models of atherosclerosis. *J Gerontol A Biol Sci Med Sci*. 2000;55(2):B85-94.
307. Thirumangalakudi L, Prakasam A, Zhang R, Bimonte-Nelson H, Sambamurti K, Kindy MS, et al. High cholesterol-induced neuroinflammation and amyloid precursor protein processing correlate with loss of working memory in mice. *J Neurochem*. 2008;106(1):475-85.
308. Vieira EL, Leonel AJ, Sad AP, Beltrao NR, Costa TF, Ferreira TM, et al. Oral administration of sodium butyrate attenuates inflammation and mucosal lesion in experimental acute ulcerative colitis. *J Nutr Biochem*. 2012;23(5):430-6.
309. Nilsson NE, Kotarsky K, Owman C, Olde B. Identification of a free fatty acid receptor, FFA2R, expressed on leukocytes and activated by short-chain fatty acids. *Biochem Biophys Res Commun*. 2003;303(4):1047-52.
310. Nohr MK, Egerod KL, Christiansen SH, Gille A, Offermanns S, Schwartz TW, et al. Expression of the short chain fatty acid receptor GPR41/FFAR3 in autonomic and somatic sensory ganglia. *Neuroscience*. 2015;290:126-37.
311. Tazoe H, Otomo Y, Karaki S, Kato I, Fukami Y, Terasaki M, et al. Expression of short-chain fatty acid receptor GPR41 in the human colon. *Biomed Res*. 2009;30(3):149-56.
312. Zhang C, Zhang M, Pang X, Zhao Y, Wang L, Zhao L. Structural resilience of the gut microbiota in adult mice under high-fat dietary perturbations. *The ISME journal*. 2012;6(10):1848-57.
313. Rajilić-Stojanović M, de Vos WM. The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS microbiology reviews*. 2014;38(5):996-1047.
314. Gilabert-Juan J, Castillo-Gomez E, Perez-Rando M, Molto MD, Nacher J. Chronic stress induces changes in the structure of interneurons and in the expression of molecules related to neuronal structural plasticity and inhibitory neurotransmission in the amygdala of adult mice. *Exp Neurol*. 2011;232(1):33-40.

315. Fitzpatrick AL, Kuller LH, Lopez OL, Diehr P, O'Meara ES, Longstreth WT, Jr., et al. Midlife and late-life obesity and the risk of dementia: cardiovascular health study. *Archives of neurology*. 2009;66(3):336-42.
316. Rosenberg GA. Vascular cognitive impairment: Biomarkers in diagnosis and molecular targets in therapy. *J Cereb Blood Flow Metab*. 2016;36(1):4-5.
317. Tucsek Z, Toth P, Sosnowska D, Gautam T, Mitschelen M, Koller A, et al. Obesity in aging exacerbates blood-brain barrier disruption, neuroinflammation, and oxidative stress in the mouse hippocampus: effects on expression of genes involved in beta-amyloid generation and Alzheimer's disease. *J Gerontol A Biol Sci Med Sci*. 2014;69(10):1212-26.
318. Kim KW, Seo H, Kwak MS, Kim D. Visceral obesity is associated with white matter hyperintensity and lacunar infarct. *Int J Obes (Lond)*. 2017;41(5):683-8.
319. Attwell D, Buchan AM, Charkpak S, Lauritzen M, Macvicar BA, Newman EA. Glial and neuronal control of brain blood flow. *Nature*. 2010;468(7321):232-43.
320. McGavock JM, Victor RG, Unger RH, Szczepaniak LS. Adiposity of the heart, revisited. *Ann Intern Med*. 2006;144:517-24.
321. Scotece M, Conde J, Lopez V, Lago F, Pino J, Gomez-Reino JJ, et al. Adiponectin and leptin: new targets in inflammation. *Basic Clin Pharmacol Toxicol*. 2014;114(1):97-102.
322. van Norden AG, de Laat KF, Gons RA, van Uden IW, van Dijk EJ, van Oudheusden LJ, et al. Causes and consequences of cerebral small vessel disease. The RUN DMC study: a prospective cohort study. Study rationale and protocol. *BMC Neurol*. 2011;11:29.
323. methodology WCCfda. ATC/DDD Index 2018 Oslo, Norway: World Health Organization, ; 2017 [Available from: https://www.whocc.no/atc_ddd_index/].
324. van Leijsen EMC, van Uden IWM, Ghafoorian M, Bergkamp MI, Lohner V, Kooijmans ECM, et al. Nonlinear temporal dynamics of cerebral small vessel disease: The RUN DMC study. *Neurology*. 2017;89(15):1569-77.
325. van Uden IW, Tuladhar AM, van der Holst HM, van Leijsen EM, van Norden AG, de Laat KF, et al. Diffusion tensor imaging of the hippocampus predicts the risk of dementia; the RUN DMC study. *Hum Brain Mapp*. 2016;37(1):327-37.
326. Wardlaw JM, Smith EE, Biessels GJ, Cordonnier C, Fazekas F, Frayne R, et al. Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. *Lancet Neurol*. 2013;12(8):822-38.
327. Ghafoorian M, Karssemeijer N, van Uden IW, de Leeuw FE, Heskes T, Marchiori E, et al. Automated detection of white matter hyperintensities of all sizes in cerebral small vessel disease. *Med Phys*. 2016;43(12):6246.
328. Fazekas F, Chawluk JB, Alavi A, Hurtig HI, Zimmerman RA. MR signal abnormalities at 1.5 T in Alzheimer's dementia and normal aging. *AJR Am J Roentgenol*. 1987;149(2):351-6.
329. Hervé D, Mangin J-F, Molko N, Bousser M-G, Chabriat H. Shape and volume of lacunar infarcts: a 3D MRI study in cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy. *Stroke*. 2005;36(11):2384-8.

330. Cordonnier C, Van der Flier W, Sluimer J, Leys D, Barkhof F, Scheltens P. Prevalence and severity of microbleeds in a memory clinic setting. *Neurology*. 2006;66(9):1356-60.
331. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189-98.
332. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry*. 1998;59 Suppl 20:22-33;quiz 4-57.
333. Hochstenbach J, Mulder T, van Limbeek J, Donders R, Schoonderwaldt H. Cognitive decline following stroke: a comprehensive study of cognitive decline following stroke. *J Clin Exp Neuropsychol*. 1998;20(4):503-17.
334. American Psychiatric A. Diagnostic and statistical manual of mental disorders : DSM-IV-TR. 4th ed., text revision. ed. Washington, DC :: American Psychiatric Association; 2000.
335. Vermeer SE, Prins ND, den Heijer T, Hofman A, Koudstaal PJ, Breteler MM. Silent brain infarcts and the risk of dementia and cognitive decline. *N Engl J Med*. 2003;348(13):1215-22.
336. Lee JH, Reed DR, Price RA. Leptin resistance is associated with extreme obesity and aggregates in families. *International Journal Of Obesity*. 2001;25:1471.
337. Franx BAA, Arnoldussen IAC, Kiliaan AJ, Gustafson DR. Weight Loss in Patients with Dementia: Considering the Potential Impact of Pharmacotherapy. *Drugs Aging*. 2017;34(6):425-36.
338. Gustafson D, Steen B, Skoog I. Body mass index and white matter lesions in elderly women. An 18-year longitudinal study. *Intl J Psychogeriatrics*. 2004;16:327-36.
339. Kwon HM, Park JH, Park JH, Jeong HY, Lim JS, Jeong HG, et al. Visceral Fat Is an Independent Predictor of Cerebral Microbleeds in Neurologically Healthy People. *Cerebrovasc Dis*. 2016;42(1-2):90-6.
340. Lioutas VA, Beiser A, Himali J, Aparicio H, Romero JR, DeCarli C, et al. Lacunar Infarcts and Intracerebral Hemorrhage Differences: A Nested Case-Control Analysis in the FHS (Framingham Heart Study). *Stroke*. 2017;48(2):486-9.
341. Higuchi S, Kabeya Y, Kato K. Visceral-to-subcutaneous fat ratio is independently related to small and large cerebrovascular lesions even in healthy subjects. *Atherosclerosis*. 2017;259:41-5.
342. Yamada S, Satow T, Fukuda A, Ito M, Saiki M. Severe underweight and cerebral microbleeds. *J Neurol*. 2012;259(12):2707-13.
343. Birdsill AC, Oleson S, Kaur S, Pasha E, Ireton A, Tanaka H, et al. Abdominal obesity and white matter microstructure in midlife. *Hum Brain Mapp*. 2017.
344. Brooks SJ, Benedict C, Burgos J, Kempton MJ, Kullberg J, Nordenskjöld R, et al. Late-life obesity is associated with smaller global and regional gray matter volumes: a voxel-based morphometric study. *International Journal of Obesity (2005)*. 2013;37(2):230-6.

345. Garcia-Garcia I, Michaud A, Dadar M, Zeighami Y, Neseliler S, Collins DL, et al. Neuroanatomical differences in obesity: meta-analytic findings and their validation in an independent dataset. *Int J Obes (Lond)*. 2018.
346. Gunstad J, Paul RH, Cohen RA, Tate DF, Spitznagel MB, Grieve S, et al. Relationship between body mass index and brain volume in healthy adults. *Int J Neurosci*. 2008;118(11):1582-93.
347. Sala M, de Roos A, van den Berg A, Altmann-Schneider I, Slagboom PE, Westendorp RG, et al. Microstructural brain tissue damage in metabolic syndrome. *Diabetes Care*. 2014;37(2):493-500.
348. Isaac V, Sim S, Zheng H, Zagorodnov V, Tai ES, Chee M. Adverse Associations between Visceral Adiposity, Brain Structure, and Cognitive Performance in Healthy Elderly. *Front Aging Neurosci*. 2011;3:12.
349. Widya RL, de Roos A, Trompet S, de Craen AJ, Westendorp RG, Smit JW, et al. Increased amygdalar and hippocampal volumes in elderly obese individuals with or at risk of cardiovascular disease. *Am J Clin Nutr*. 2011;93(6):1190-5.
350. Figlewicz DP, Woods SC. Adiposity signals and brain reward mechanisms. *Trends Pharmacol Sci*. 2000;21(7):235-6.
351. Pannacciulli N, Le DSNT, Chen K, Reiman EM, Krakoff J. Relationships between plasma leptin concentrations and human brain structure: a voxel-based morphometric study. *Neuroscience letters*. 2007;412(3):248-53.
352. Yokum S, Stice E. Initial body fat gain is related to brain volume changes in adolescents: A repeated-measures voxel-based morphometry study. *Obesity (Silver Spring)*. 2017;25(2):401-7.
353. Smucny J, Cornier MA, Eichman LC, Thomas EA, Bechtell JL, Tregellas JR. Brain structure predicts risk for obesity. *Appetite*. 2012;59(3):859-65.
354. Kantorova E, Chomova M, Kurca E, Sivak S, Zelenak K, Kucera P, et al. Leptin, adiponectin and ghrelin, new potential mediators of ischemic stroke. *Neuro Endocrinol Lett*. 2011;32(5):716-21.
355. Narita K, Kosaka H, Okazawa H, Murata T, Wada Y. Relationship between plasma leptin level and brain structure in elderly: a voxel-based morphometric study. *Biol Psychiatry*. 2009;65(11):992-4.
356. Bonda DJ, Stone JG, Torres SL, Siedlak SL, Perry G, Kryscio R, et al. Dysregulation of leptin signaling in Alzheimer disease: evidence for neuronal leptin resistance. *J Neurochem*. 2014;128(1):162-72.
357. Meek TH, Matsen ME, Dorfman MD, Guyenet SJ, Damian V, Nguyen HT, et al. Leptin action in the ventromedial hypothalamic nucleus is sufficient, but not necessary, to normalize diabetic hyperglycemia. *Endocrinology*. 2013;154(9):3067-76.
358. Sakai T, Kusakabe T, Ebihara K, Aotani D, Yamamoto-Kataoka S, Zhao M, et al. Leptin restores the insulinotropic effect of exenatide in a mouse model of type 2 diabetes with increased adiposity induced by streptozotocin and high-fat diet. *American journal of physiology Endocrinology and metabolism*. 2014;307(8):E712-9.

359. Davis C, Mudd J, Hawkins M. Neuroprotective effects of leptin in the context of obesity and metabolic disorders. *Neurobiol Dis.* 2014;72 Pt A:61-71.
360. Zeki Al Hazzouri A, Stone KL, Haan MN, Yaffe K. Leptin, mild cognitive impairment, and dementia among elderly women. *J Gerontol A Biol Sci Med Sci.* 2013;68(2):175-80.
361. Larsen MM, Madsen PH, Markenvard JD. [Thrombolysis in treatment of cardiac arrest caused by pulmonary embolism]. *Ugeskr Laeger.* 2008;170(36):2803.
362. Flegal KM, Shepherd JA, Looker AC, Graubard BI, Borrud LG, Ogden CL, et al. Comparisons of percentage body fat, body mass index, waist circumference, and waist-stature ratio in adults. *Am J Clin Nutr.* 2009;89(2):500-8.
363. Kiliaan AJ, Arnoldussen IA, Gustafson DR. Adipokines: a link between obesity and dementia? *Lancet Neurol.* 2014;13(9):913-23.
364. Li C, Ford ES, McGuire LC, Mokdad AH. Increasing trends in waist circumference and abdominal obesity among US adults. *Obesity (Silver Spring).* 2007;15(1):216-24.
365. Toth MJ, Tchernof A, Sites CK, Poehlman ET. Menopause-related changes in body fat distribution. *Annals of the New York Academy of Sciences.* 2000;904:502-6.
366. NIH/NHLBI. Classification of Overweight and Obesity by BMI, Waist Circumference, and Associated Disease Risks 2017 [Available from: https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi_dis.htm].
367. Taki Y, Kinomura S, Sato K, Inoue K, Goto R, Okada K, et al. Relationship between body mass index and gray matter volume in 1,428 healthy individuals. *Obesity (Silver Spring).* 2008;16(1):119-24.
368. Gustafson D, Lissner L, Bengtsson C, Bjorkelund C, Skoog I. A 24-year follow-up of body mass index and cerebral atrophy. *Neurology.* 2004;63(10):1876-81.
369. Whelan EA, Sandler DP, McConaughy DR, Weinberg CR. Menstrual and reproductive characteristics and age at natural menopause. *American journal of epidemiology.* 1990;131:625-32.
370. Markou A, Duka T, Prelevic GM. Estrogens and brain function. *Hormones (Athens).* 2005;4(1):9-17.
371. Backman K, Joas E, Waern M, Ostling S, Guo X, Blennow K, et al. 37 Years of Body Mass Index and Dementia: Effect Modification by the APOE Genotype: Observations from the Prospective Population Study of Women in Gothenburg, Sweden. *J Alzheimers Dis.* 2015;48(4):1119-27.
372. Joas E, Backman K, Gustafson D, Ostling S, Waern M, Guo X, et al. Blood pressure trajectories from midlife to late life in relation to dementia in women followed for 37 years. *Hypertension.* 2012;59(4):796-801.
373. Kivipelto M, Solomon A. Cholesterol as a risk factor for Alzheimer's disease - epidemiological evidence. *Acta Neurol Scand Suppl.* 2006;185:50-7.
374. Gustafson DR, Karlsson C, Skoog I, Rosengren L, Lissner L, Blennow K. Mid-life adiposity factors relate to blood-brain barrier integrity in late life. *J Intern Med.* 2007;262(6):643-50.

375. Barrett-Connor E, Edelstein SL, Corey-Bloom J, Wiederholt WC. Weight loss precedes dementia in community-dwelling older adults. *J Am Geriatr Soc.* 1996;44:1147-52.
376. Gustafson D, Bäckman K, Joas E, Waern M, Östling S, Guo X, et al. A 37-year longitudinal follow-up of body mass index and dementia in women. *Journal of Alzheimer's disease : JAD.* 2012;28:162-71.
377. Erkinjuntti T. Subcortical vascular dementia. *Cerebrovasc Dis.* 2002;13 Suppl 2:58-60.
378. Roman GC, Erkinjuntti T, Wallin A, Pantoni L, Chui HC. Subcortical ischaemic vascular dementia. *Lancet Neurol.* 2002;1(7):426-36.
379. Lear SA, Humphries KH, Kohli S, Birmingham CL. The use of BMI and waist circumference as surrogates of body fat differs by ethnicity. *Obesity (Silver Spring).* 2007;15(11):2817-24.
380. Besser LM, Gill DP, Monsell SE, Brenowitz W, Meranus D, Kukull W, et al. Body mass index, weight change, and clinical progression in mild cognitive impairment and Alzheimer's disease. *Alzheimer disease and associated disorders.* 2014;28(1):36.
381. Johnson DK, Wilkins CH, Morris JC. Accelerated weight loss may precede diagnosis in Alzheimer disease. *Arch Neurol.* 2006;63(9):1312-7.
382. Johnston JM, Hu WT, Fardo DW, Greco SJ, Perry G, Montine TJ, et al. Low plasma leptin in cognitively impaired ADNI subjects: gender differences and diagnostic and therapeutic potential. *Curr Alzheimer Res.* 2014;11(2):165-74.
383. Littlejohns TJ, Kos K, Henley WE, Cherubini A, Ferrucci L, Lang IA, et al. Serum leptin and risk of cognitive decline in elderly italians. *J Alzheimers Dis.* 2015;44(4):1231-9.
384. van Himbergen TM, Beiser AS, Ai M, Seshadri S, Otokozawa S, Au R, et al. Biomarkers for insulin resistance and inflammation and the risk for all-cause dementia and alzheimer disease: results from the Framingham Heart Study. *Arch Neurol.* 2012;69(5):594-600.
385. Oania R, McEvoy LK. Plasma leptin levels are not predictive of dementia in patients with mild cognitive impairment. *Age Ageing.* 2015;44(1):53-8.
386. Peiser C, McGregor GP, Lang RE. Binding and internalization of leptin by porcine choroid plexus cells in culture. *Neurosci Lett.* 2000;283(3):209-12.
387. Schulz C, Paulus K, Lehnert H. Adipocyte-brain: crosstalk. Results and problems in cell differentiation. 2010;52:189-201.
388. Morash B, Li A, Murphy PR, Wilkinson M, Ur E. Leptin gene expression in the brain and pituitary gland. *Endocrinology.* 1999;140(12):5995-8.
389. Wiesner G, Vaz M, Collier G, Seals D, Kaye D, Jennings G, et al. Leptin is released from the human brain: influence of adiposity and gender. *J Clin Endocrinol Metab.* 1999;84(7):2270-4.
390. Friedman JM. Leptin, leptin receptors, and the control of body weight. *Nutr Rev.* 1998;56(2 Pt 2):s38-46; discussion s54-75.

391. Lissner L, Karlsson C, Lindroos AK, Sjöström L, Carlsson B, Carlsson L, et al. Birth weight, adulthood BMI, and subsequent weight gain in relation to leptin levels in Swedish women. *Obes Res*. 1999;7(2):150-4.
392. Sinha MK, Songer T, Xiao Q, Sloan JH, Wang J, Ji S, et al. Analytical validation and biological evaluation of a high molecular-weight adiponectin ELISA. *Clinical chemistry*. 2007;53(12):2144-51.
393. Drolet ea. Fat depot-specific impact of visceral obesity on adipocyte adiponectin release in women. *Obesity (Silver Spring)*. 2008;17:424-30.
394. Steen B, Djurfeldt H. The gerontological and geriatric population studies in Gothenburg, Sweden. *Z Gerontol*. 1993;26:163-9.
395. Bengtsson C, Blohme G, Hallberg L, Hällström T, Isaksson B, Korsan-Bengtson K, et al. The study of women in Gothenburg 1968-79. A population study. General design, purpose and sampling results. *Acta Med Scand*. 1973;193:311-8.
396. Bengtsson C, Ahlquist M, Andersson K, Björkelund C, Lissner L, Söderström M. The Prospective Population Study of Women in Gothenburg, Sweden, 1968-69 to 1992-93. A 24-year follow-up study with special reference to participation, representativeness, and mortality. *Scand J Primary Health Care*. 1997;15:214-9.
397. Skoog I, Nilsson L, Palmertz B, Andreasson LA, Svanborg A. A population-based study of dementia in 85-year-olds. *N Engl J Med*. 1993;328(3):153-8.
398. Skoog I, Lernfelt B, Landahl S, Palmertz B, Andreasson LA, Nilsson L, et al. 15-year longitudinal study of blood pressure and dementia. *Lancet*. 1996;347(9009):1141-5.
399. Jorm AF, Korten AE, Henderson AS. The prevalence of dementia: a quantitative integration of the literature. *Acta Psychiatr Scand*. 1987;76(5):465-79.
400. Åsberg M, Perris C, Schalling D, Sedvall G. The CPRS - development and applications of a psychiatric rating scale. *Acta Psychiatr Scand*. 1978;suppl 271.
401. Bengtsson C, Hallberg L, Noppa H, Tibblin E. Anthropometric data in middle-aged women. The population study of women in Goteborg 1968-1969. *Acta morphologica Neerland-Scandinavica*. 1979;17(2):133-43.
402. Blennow K, Ricksten A, Prince JA, Brookes AJ, Emahazion T, Wasslavik C, et al. No association between the alpha2-macroglobulin (A2M) deletion and Alzheimer's disease, and no change in A2M mRNA, protein, or protein expression. *J Neural Transm*. 2000;107:1065-79.
403. Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry*. 1979;134:382-9.
404. Kerwin DR, Gaussoin SA, Chlebowski RT, Kuller LH, Vitolins M, Coker LH, et al. Interaction between body mass index and central adiposity and risk of incident cognitive impairment and dementia: results from the Women's Health Initiative Memory Study. *J Am Geriatr Soc*. 2011;59(1):107-12.
405. National Heart LaBI. Classification of Overweight and Obesity by BMI, Waist Circumference, and Associated Disease Risks: National Institutes of Health; 2017 [Available from: https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi_dis.htm].

406. Gustafson D, Rothenberg E, Blennow K, Steen B, Skoog I. An 18-year follow-up of overweight and risk of Alzheimer disease. *Arch Intern Med.* 2003;163(13):1524-8.
407. Gustafson DR, Mielke MM, Keating SA, Holman S, Minkoff H, Crystal HA. Leptin, adiponectin and cognition in middle-aged HIV-infected and uninfected women. The Brooklyn Women's Interagency HIV Study *J Gerontol Geriatric Res.* 2015;4(5).
408. Gustafson DR, Mielke MM, Tien PC, Valcour V, Cohen M, Anastos K, et al. Anthropometric measures and cognition in middle-aged HIV-infected and uninfected women. The Women's Interagency HIV Study. *Journal of neurovirology.* 2013;19(6):574-85.
409. ADRD summit 2016 committee. Alzheimer's disease-related dementias. 2016.
410. Wancata J, Borjesson-Hanson A, Ostling S, Sjogren K, Skoog I. Diagnostic criteria influence dementia prevalence. *Am J Geriatr Psychiatry.* 2007;15(12):1034-45.
411. Zettergren A, Kern S, Gustafson D, Gudmundsson P, Sigstrom R, Ostling S, et al. The ACE Gene Is Associated with Late-Life Major Depression and Age at Dementia Onset in a Population-Based Cohort. *Am J Geriatr Psychiatry.* 2017;25(2):170-7.
412. Wu YT, Fratiglioni L, Matthews FE, Lobo A, Breteler MM, Skoog I, et al. Dementia in western Europe: epidemiological evidence and implications for policy making. *Lancet Neurol.* 2016;15(1):116-24.
413. Morley JE. Nutrition and the Brain. *Clinics in geriatric medicine.* 2010;26(1):89-98.
414. Holden KF, Lindquist K, Tylavsky FA, Rosano C, Harris TB, Yaffe K, et al. Serum leptin level and cognition in the elderly: Findings from the Health ABC Study. *Neurobiol Aging.* 2009;30(9):1483-9.
415. Zeki Al Hazzouri A, Haan MN, Whitmer RA, Yaffe K, Neuhaus J. Central obesity, leptin and cognitive decline: the Sacramento Area Latino Study on Aging. *Dement Geriatr Cogn Disord.* 2012;33(6):400-9.
416. Finger EC. Increased heart rate and energy expenditure in frontotemporal dementia. *Brain.* 2017;140(1):10-2.
417. Ahmed RM, Latheef S, Bartley L, Irish M, Halliday GM, Kiernan MC, et al. Eating behavior in frontotemporal dementia: Peripheral hormones vs hypothalamic pathology. *Neurology.* 2015;85(15):1310-7.
418. Magalhaes CA, Carvalho MG, Sousa LP, Caramelli P, Gomes KB. Leptin in Alzheimer's disease. *Clin Chim Acta.* 2015;450:162-8.
419. OECD. OECD OBESITY UPDATE 2017 2017 [updated June 2017; cited 2018 30-01-2018]. Available from: <http://www.oecd.org/health/obesity-update.htm>.
420. Brown E, Umino Y, Loi T, Solessio E, Barlow R. Anesthesia can cause sustained hyperglycemia in C57/BL6J mice. *Visual neuroscience.* 2005;22(5):615-8.
421. Lee HY, Jeong KH, Choi CS, International Mouse Phenotyping C. In-depth metabolic phenotyping of genetically engineered mouse models in obesity and diabetes. *Mamm Genome.* 2014;25(9-10):508-21.

422. Schreyer SA, Vick C, Lystig TC, Mystkowski P, LeBoeuf RC. LDL receptor but not apolipoprotein E deficiency increases diet-induced obesity and diabetes in mice. *Am J Physiol Endocrinol Metab.* 2002;282(1):E207-14.
423. Fox JG. *Laboratory animal medicine*: Elsevier; 2015.
424. Gorelick PB, Scuteri A, Black SE, Decarli C, Greenberg SM, Iadecola C, et al. Vascular contributions to cognitive impairment and dementia: a statement for healthcare professionals from the american heart association/american stroke association. *Stroke.* 2011;42(9):2672-713.
425. Zlokovic BV. Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nature reviews Neuroscience.* 2011;12(12):723-38.
426. Landsberg L, Aronne LJ, Beilin LJ, Burke V, Igel LI, Lloyd-Jones D, et al. Obesity-related hypertension: pathogenesis, cardiovascular risk, and treatment--a position paper of the The Obesity Society and The American Society of Hypertension. *Obesity (Silver Spring).* 2013;21(1):8-24.
427. Fasshauer M, Bluher M. Adipokines in health and disease. *Trends in pharmacological sciences.* 2015;36(7):461-70.
428. Pelgrim CE, Franx BAA, Snabel J, Kleemann R, Arnoldussen IAC, Kiliaan AJ. Butyrate Reduces HFD-Induced Adipocyte Hypertrophy and Metabolic Risk Factors in Obese LDLr-/-Leiden Mice. *Nutrients.* 2017;9(7).
429. Holland WL, Adams AC, Brozinick JT, Bui HH, Miyauchi Y, Kusminski CM, et al. An FGF21-adiponectin-ceramide axis controls energy expenditure and insulin action in mice. *Cell Metab.* 2013;17(5):790-7.
430. Beltowski J. Adiponectin and resistin--new hormones of white adipose tissue. *Medical science monitor : international medical journal of experimental and clinical research.* 2003;9(2):RA55-61.
431. Ouchi N, Kihara S, Funahashi T, Nakamura T, Nishida M, Kumada M, et al. Reciprocal association of C-reactive protein with adiponectin in blood stream and adipose tissue. *Circulation.* 2003;107(5):671-4.
432. Weyer C, Funahashi T, Tanaka S, Hotta K, Matsuzawa Y, Pratley RE, et al. Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab.* 2001;86(5):1930-5.
433. Cnop M, Havel PJ, Utzschneider KM, Carr DB, Sinha MK, Boyko EJ, et al. Relationship of adiponectin to body fat distribution, insulin sensitivity and plasma lipoproteins: evidence for independent roles of age and sex. *Diabetologia.* 2003;46(4):459-69.
434. Luchsinger JA, Gustafson DR. Adiposity and Alzheimer's disease. *Curr Opin Clin Nutr Metab Care.* 2009;12(1):15-21.
435. Albanese E, Davis B, Jonsson PV, Chang M, Aspelund T, Garcia M, et al. Overweight and Obesity in Midlife and Brain Structure and Dementia 26 Years Later: The AGES-Reykjavik Study. *Am J Epidemiol.* 2015;181(9):672-9.

436. Loef M, Walach H. Midlife obesity and dementia: meta-analysis and adjusted forecast of dementia prevalence in the United States and China. *Obesity* (Silver Spring). 2013;21(1):E51-5.
437. Nepal B, Brown LJ, Anstey KJ. Rising midlife obesity will worsen future prevalence of dementia. *PLoS One*. 2014;9(9):e99305.
438. Palmer BF, Clegg DJ. The sexual dimorphism of obesity. *Mol Cell Endocrinol*. 2015;402:113-9.
439. Van Sinderen ML, Steinberg GR, Jorgensen SB, Honeyman J, Chow JD, Herridge KA, et al. Effects of Estrogens on Adipokines and Glucose Homeostasis in Female Aromatase Knockout Mice. *PLoS One*. 2015;10(8):e0136143.
440. Luchsinger JA, Patel B, Tang MX, Schupf N, Mayeux R. Measures of adiposity and dementia risk in elderly persons. *Arch Neurol*. 2007;64(3):392-8.
441. Tang LL, Tang XH, Li X, Yu HB, Xie ZG, Liu XY, et al. Effect of high-fat or high-glucose diet on obesity and visceral adipose tissue in mice. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao*. 2014;36(6):614-9.
442. Cordner ZA, Tamashiro KL. Effects of high-fat diet exposure on learning & memory. *Physiol Behav*. 2015;152(Pt B):363-71.
443. Neschen S, Moore I, Regittnig W, Yu CL, Wang Y, Pypaert M, et al. Contrasting effects of fish oil and safflower oil on hepatic peroxisomal and tissue lipid content. *Am J Physiol-Endoc M*. 2002;282(2):E395-E401.
444. Ikemoto S, Takahashi M, Tsunoda N, Maruyama K, Itakura H, Ezaki O. High-fat diet-induced hyperglycemia and obesity in mice: differential effects of dietary oils. *Metabolism-Clinical and Experimental*. 1996;45(12):1539-46.
445. Costantini L, Molinari R, Farinon B, Merendino N. Impact of Omega-3 Fatty Acids on the Gut Microbiota. *International journal of molecular sciences*. 2017;18(12).
446. Xu J, Chen X, Yu S, Su Y, Zhu W. Effects of Early Intervention with Sodium Butyrate on Gut Microbiota and the Expression of Inflammatory Cytokines in Neonatal Piglets. *PLoS One*. 2016;11(9):e0162461.
447. Zhang T, Ding C, Zhao M, Dai X, Yang J, Li Y, et al. Sodium Butyrate Reduces Colitogenic Immunoglobulin A-Coated Bacteria and Modifies the Composition of Microbiota in IL-10 Deficient Mice. *Nutrients*. 2016;8(12).
448. Lien EL, Richard C, Hoffman DR. DHA and ARA addition to infant formula: Current status and future research directions. *Prostaglandins, leukotrienes, and essential fatty acids*. 2018;128:26-40.
449. Mori TA. Omega-3 fatty acids and cardiovascular disease: epidemiology and effects on cardiometabolic risk factors. *Food & function*. 2014;5(9):2004-19.
450. Zehr KR, Walker MK. Omega-3 polyunsaturated fatty acids improve endothelial function in humans at risk for atherosclerosis: A review. *Prostaglandins & other lipid mediators*. 2018;134:131-40.

451. Brahe LK, Astrup A, Larsen LH. Is butyrate the link between diet, intestinal microbiota and obesity-related metabolic diseases? *Obes Rev.* 2013;14(12):950-9.
452. van der Beek CM, Bloemen JG, van den Broek MA, Lenaerts K, Venema K, Buurman WA, et al. Hepatic Uptake of Rectally Administered Butyrate Prevents an Increase in Systemic Butyrate Concentrations in Humans. *J Nutr.* 2015;145(9):2019-24.

Dankwoord | Acknowledgements

Ha Lieverds,

Aan een PhD project werk je zeker niet alleen, niet alleen in het onderzoek maar ook buiten werktijd hebben velen hun steentje bijgedragen aan het tot stand komen van dit proefschrift en nog belangrijker de vorming van mij als persoon. Dus hieronder een paar pagina's van mijn dankbaarheid aan een ieder die mij heeft geholpen gedurende deze mooie jaren.

Allereerst mijn begeleiders, jullie hebben mij alle kansen geboden om dit proefschrift prachtig te voltooien. Co-promotoren, dr. Amanda Kiliaan en dr. Maximilian Wiesmann. **Lieve Amanda**, allereerst dank je wel voor al het vertrouwen en vele kansen die je me hebt gegeven. Daarnaast voor al het geduld met het doorlezen van de talloze versies van manuscripten, op rare zinnen zonder werkwoord, vage beschrijvingen en vreemde typefouten.... ('Dit is geen zin!') Je hebt me enorm veel laten leren en inzien, niet alleen op wetenschappelijk gebied, maar vooral ook op samenwerkings- en werelddniveau. Mijn grootste dank spreek ik uit dat je er voor mijn gevoel dat je er altijd bent, er is altijd tijd en ruimte voor vragen. Bewonderingswaardig is hoe je vecht met zoveel passie als een leeuw voor je welpjes; onze wetenschaps-mama! Dank dat ik vanaf jouw schouders mocht meekijken en leren.

Lieve **Max**, Meister Max.....Tsjaa Max, we worden oud, we kennen elkaar nu al 11 jaar! Onze eerste kennismaking tijdens introweek van onze studie biologie was bijzonder; je kon me echt niet verstaan met m'n snelle, Limburgse gebrabbel, maar toch bleef je lief lachen en knikken. Maar nu 11 jaar later, lees je mijn gezicht als geen ander! Het is typerend dat we, zonder af te spreken, tegelijkertijd het dansje van de achtergrondanseressen van Helene Fischer inzetten (ook nog allebei op rechts...:D). Dank voor al die jaren vriendschap en dat er nog vele mogen volgen!

Mijn promotoren Prof. dr. Deborah Gustafson en Prof. dr. Otto Boerman. **Deborah**, thanks for the great opportunity to come to NYC to learn, and subsequently all the skype-meetings (you are an Early Brooklyn bird)! Especially, thank you for your bright light on our latest 'artwork', you guided me out of the dark. I learned so much!

Otto, bedankt dat je mijn promotor wilt zijn als ad interim afdelingshoofd van de Afdeling Anatomie. Jij begeleidt de afdeling Anatomie dit komende jaar naar een nieuwe, mooie fase!

Geachte leden van de manuscriptcommissie en de corona: **Prof. dr. Leo Joosten, Prof. dr. Susanne la Fleur, dr. Esther Aarts, Prof. dr. Arend Heerschap, Prof. dr. Ellen Blaak, Prof. dr. Niels Riksen en dr. Peter Wielinga**, hartelijk dank dat jullie mijn proefschrift hebben willen beoordelen, er vandaag zijn om dit te bediscussiëren en hopelijk ook deze mijlpaal te vieren.

Mijn twee paranimfen, Tim en Debby. **Timmie** (mister Adobe, template-Tim), Je hebt echt een geweldige humor en enorm sterk relativiseringsvermogen. Ondanks (vele) tegenslagen ben jij in staat om wind in je zeilen te houden en je koers te behouden! Timmie, jij kunt sfeer in de kamer maken met geweldige opmerkingen of door iemand even lekker uitdagen met een strak gezicht. ('Oww halve dagen vandaag?', 'Ilse die appel is nog niet op' of 'Is dit mijn probleem?!') Lieve Tim, dank dat je bij mijn promotie wilt zijn, succes met het afronden van je proefschrift bovenal een hele mooie tijd toegewenst in de Mayo Clinic.

Lieve **Debby**, klein maar heel fijn! Je kwam een masterstage volgen en ik mocht je begeleiden en na een tijdje leek het alsof je mijn gedachte kon lezen. (Soms had je al iets uitgevoerd, voordat het überhaupt de gedachte in mijn hoofd was ontstaan. ☺) Je slingert nu hier en daar wat behoorlijk rake opmerking door de kamer (en je lijkt zo lief), die de sfeer verhogen. Je gaat een prachtige, grootse studie opzetten, het is nu heel hard werken maar het gaat je zeker vruchten opleveren. Alleen nog even zorgen dat je deze vruchten vangt en dan kun je achterover leunen tot je pensioen. ☺(maar pas op, dan zou je je kunnen gaan vervelen). Dank je wel dat jij mijn paranimf wilt zijn!

Collega's-Anatomie, lief en leed en veel (slechte) muziek werd er gedeeld in onze oio-/aio-kamer. Lieve **Anouk**, we zijn allebei in precies dezelfde status na een weekje MRI-experimenten. En hoe fijn is het dan als jij lekker die droge, directe humor op me loslaat, ik lach er dagen later vaak nog een keer om. Ik heb enorm veel bewondering voor jou, met name hoe jij het al 3 jaar regelt om heen en weer te reizen tussen Voorburg en Nijmegen, zeker in de context van een jong gezin. Dank dat ik, als uitzondering, Micha met een zachte-g mag uitspreken en voor het geweldige Haagse woord: Kikkaah (kikker). **Jeroen, Teun, Bart en Dylan** dank voor alle gezelligheid en gezellige werksfeer. **Bram, Vivienne en Jos**, als analisten is jullie energie en zijn jullie ideeën van groot belang geweest voor mijn onderzoek. Dank voor al jullie tijd, inspiratie en hulp. **Bram**, jouw no-nonsense instelling en nuchterheid is bijna helend! (Soms is mijn motto: Stay calm and think like Bram!) **Vivienne**, met jou zat ik vaak uren gezellig te kletsen bij een aantal (stinkende) muizen voor verschillende doeleinden. Jouw 'Fingerspitzengefühl' m.b.t. dissectie of bloed afnemen, was je een essentieel onderdeel van de muizenstudies in dit proefschrift. **Jos**, als nestor van de afdeling weet je precies wat er bij iedereen speelt

en leef je mee. Je bent de rust zelve en legt in alle rust de fijne kneepjes van de techniek uit. Maar natuurlijk zijn er nog veel mensen die al die jaren bijgedragen hebben aan de prettige werksfeer binnen Anatomie, alle collega's van Anatomie dank voor alle hulp, gesprekken en vooral gezelligheid bij de Dagjes-uit en Kerstborrels. **Prof. dr. Dirk Ruiter**, dank voor je betrokkenheid bij mijn promotietraject en vooral dat je vandaag als rector bij mijn promotie aanwezig wilt zijn. **Prof. dr. Tamás Kozicz, Tamás**, a man of many bright and original ideas! Respect how you manage to travel around the world (in less than 80-days.. ☺) Good luck in the Mayo Clinic!

Het is altijd leuk erachter te komen dat: 'It's a small world after all', zeker als je ex-collega's onverwachts tegenkomt op bijvoorbeeld de jaarlijkse NeuroVacuaire borrel, een congres, of tijdens de begeleiding van studenten. **Diane, Maartje, Rick, Valerio, Nienke en Carola** dank voor alle hulp, kennis, inspiratie, fijne sfeer en natuurlijk tot ziens! **Carola** dank dat ik je paranimf mocht zijn en de vriendschap die is gebleven.

Samenwerkingen zijn essentieel binnen een promotietraject en deze kan ik dan ook zeker niet onbenoemd laten. **Peter, Robert, Martine, Teake, Wim, Nicole, Ivana, Cindy, Bart en Lars (TNO Leiden en Zeist)** dank voor al het harde werken, aanleveren van de muizen, data en resultaten en natuurlijk de vele sparmomenten! Dank voor de enorm fijne samenwerking!!

Gabriele, Marieke, en Ric (Mead Johnson, RB, Nijmegen) dank voor alle feedback, het doorlezen van onze manuscripten, posters en sparmomenten tijdens de meetings. Bedankt voor de fijne samenwerking!

Prof. dr. Frank-Erik de Leeuw en **dr. Esther Leijsen** dank voor alle kansen, input en de talloze versies lezen van het RUN DMC-artikel. Esther, speciaal dat we in dezelfde week onze verdediging hebben. Tot ziens!

Wat doe je als opnieuw opstarten van Paravision niet lukt, of loskoppelen van de stekker de situatie niet verbeterd, of je MRI-signaal wel heel apart is? Dan bel je gewoon **Super Andor** (52287). Andor komt het namelijk zo snel mogelijk voor je maken, bij een kopje koffie en misschien een snoepje (geleend bij de biotechnici van PRIME). Andor, je bent altijd bereikbaar, denkt 'out of the box' en oplossingsgericht, waardoor het altijd weer op z'n pootjes terecht komt met hulp van Super Andor. Dank voor het 'opvrolijken' van mijn scandagen en hulp bij de gekste problemen. **Sjaak van Asten**, voor de handliggend zou zijn: 'Sjaak bedankt voor alle hulp rondom de Software bij de Biospec'. Maar naast alle hulp met betrekking tot software, bedank ik je het meest voor de positieve rust die je uitstraalt. Die rust is gelukkig besmettelijk. Dus het is zo fijn om jou op een hectische dag tegen te komen, dan kun je de rest van de dag er weer compleet 'zen' tegenaan.

Biotechnici van PRIME, **Bianca, Iris, Wilma, Mike, Kitty en Karin** met jullie hulp werden de vervelendste momenten van een experiment juist vaak heel gezellig! Dank voor al jullie hulp rondom de muizenstudies de afgelopen jaren! Owja, ik heb nog 1 vraag? ☺

Ook wil ik graag de studenten, **Ilse, Rikko, Joost, Simone, Eline, Charlotte, Bart, Debby, David, Daphne, Milly, Mieke, Sophie, Lotte, Lonneke en Natalia** bedanken voor jullie inzet en hulp bij de gedragstesten, immunohistochemie, qRT-PCRs en het uitwerken van alle data. Ik wens jullie een prachtige (wetenschappelijke) carrière, maar vooral veel geluk, liefde en gezondheid. Tot ziens!

Mijn familie en vrienden hebben veel betekend voor me de afgelopen jaren en mogen daarom niet onbenoemd blijven in dit dankwoord.

Petra, Mayke, Gunda, Yvet, Jannah, Marita, Elise, ...oftewel HET TEAM. Bij jullie kon ik de bergen aan 'appelmoeshoofden' kwijt. Het is moeilijk te beschrijven, maar het is mooi wat jullie als team voor mij hebben betekend binnen sport, kampioenschappen en vriendschappen. Proost!!

HomieZZ, Laura, Tanja, Saskia, en Marleen bedankt voor al jullie jaren aan vriendschap vanaf de basisschool/middelbare school. Voor mij voelt het nog steeds als datzelfde, zeer vertrouwde vriendengroepje.....alleen ietsje ouder.. ☺ Ik vergeet niet meer dat de moeder van Sas zei: 'Ow, maar in 't bejaardenhuis zitten jullie samen te bridgen hoor! Jullie raken elkaar niet meer kwijt.' Dus daar gaan we dan maar gewoon vanuit, dus ik hoop nog lang van deze vriendschap te mogen genieten!

Lieve **Dominique**, wat kan ik lachen met jou! Dank voor je hulp met het eerste epidemiologische artikel! Daarnaast voor al je energie, enthousiasme en interesse! Proost!

'**Dat doe je niet!**' Lieve **Harry**(of was het nu **Harrie**), **Liëtte, Gerhard, Bernadette, Remco en Suzan**, bedankt voor alle gezellige uitjes, lachstuipen en interesse in de muizen en studies.

Lieve **Kemmasothic**, dank voor alle gezelligheid, alle zinnige maar vooral onzinnige gesprekken tijdens de carnaval, vriendenweekenden of andere feestjes. Vooral dank voor het omtoveren van een kikker tot mijn prins, me redden met soep eten, lekker high tea-en, filterloze-momentjes en de kennis over **DE** jaren 90 (vooral van vóór 98). Er gaan zeker nog vele gezellige jaren volgen!

Lieve **familie en vrienden**, jullie hebben me van jongs af aan op vele manieren gesteund en voor me betekend, zonder jullie was dit proefschrift nooit tot stand

gekomen. Dus heel veel dank voor alle hulp rondom bijvoorbeeld mijn studie, het huis, tuin, voorzien van lekkere maaltijden, alle gastvrijheid, vooral alle mogelijkheden en er gewoon zijn op die belangrijke momenten. Met name **Paul** en **May** dank voor de enorme steun, gastvrijheid en enorm lekkere maaltijden (dat is wel even iets anders dan mijn 'pakjes' maaltijden). **Mayke, Sanne, Gijs** en **Peter**, wat ben ik enorm trots op jullie! Jullie zijn **HET** ultieme voorbeeld dat opgegeven geen optie is. De positieve, levenslustige energie waarmee jullie leven is bewonderenswaardig en misschien zelfs een inspiratie. **Mayke, Sanne** en **Gijs** dank je wel dat jullie onze twee kleine aapjes willen helpen met groot worden! ☺

Thijs en **Steffi**, mooi om jullie voor elkaar te zien vechten en samen sterk te zien. Lief 'klein' broertje, jij bent krachtig en liefdevol tegelijk, dank voor de fijne onbezorgde jeugd samen met de Duplo!

Pap en **mam**, dank voor alle kansen en mogelijkheden waar jullie zo hard voor gewerkt hebben. **Mam**, jij hebt ons geleerd en laten zien wat volharding en toewijding is. Lieve mama, ik hoop dat je nu ook ruimte maakt voor jouw mooie dromen. Lieve **pap**, op jou ben ik gewoon heel erg trots. Jij hebt laten zien dat je met humor, liefde, positiviteit, gecombineerd met realiteitszin een heel eind komt in het leven. Het allermooiste wat je ons hebt laten zien is dat je altijd moet doen wat je leuk vindt, niet alles in één keer kan en familie je fundering is voor al je dromen. Pap, mocht er dan toch één kleine kans zijn dat Petrus werkelijk aan de poort staat, kan hij je vandaag dan even een dagje vrij geven?

Roel, Eva en **Pleun**, jullie laat ik niet meer los! **Roel**, jij bent als een rots voor mij en dank je wel dat je me een andere mogelijkheid kan laten zien als dat nodig is. **Eva** en **Pleun**, wat een schatjes zijn jullie en wat ben ik trots op jullie! **Lieve schatjes** dankjewel dat jullie mij laten zien en voelen waar het leven werkelijk om draait. Houd van jullie!

List of Publications

1. Kiliaan AJ, **Arnoldussen IAC**, Gustafson DR.
Adipokines: a link between obesity and dementia?
Lancet Neurology. 2014. (5-year impact factor: 28.0).
2. **Arnoldussen IAC**, Kiliaan AJ, Gustafson DR.
Obesity and dementia: adipokines interact with the brain.
European Neuropsychopharmacology. 2014. (5-year impact factor: 4.1).
3. **Arnoldussen IAC**, Kiliaan AJ.
Impact of DHA on metabolic diseases from womb to tomb.
Marine Drugs. 2014. (5-year impact factor: 4.2).
4. **Arnoldussen IAC**, Zerbi V, Wiesmann M, Noordman RH, Bolijn S, Mutsaers MP, Dederen PJ, Kleemann R, Kooistra T, van Tol EA, Gross G, Schoemaker MH, Heerschap A, Wielinga PY, Kiliaan AJ.
Early intake of long-chain polyunsaturated fatty acids preserves brain structure and function in diet-induced obesity.
The Journal of Nutritional Biochemistry. 2016. (5-year impact factor: 4.7).
5. **Arnoldussen IAC**, Wiesmann M, Pelgrim CE, Wielemaker EM, van Duyvenvoorde W, Amaral-Santos PL, Verschuren L, Keijser BJ, Heerschap A, Kleemann R, Wielinga PY, Kiliaan AJ.
Butyrate restores HFD-induced adaptations in brain function and metabolism in mid-adult obese mice.
International Journal Obesity. 2017. (5-year impact factor: 5.1).
6. **Arnoldussen IAC**, Sundh V, Bäckman K, Kern S, Östling S, Blennow K, Zetterberg H, Skoog I, Kiliaan AJ, Gustafson DR.
A 10-Year Follow-Up of Adiposity and Dementia in Swedish Adults Aged 70 Years and Older.
Journal of Alzheimers Disease. 2018. (5-year impact factor: 3.5).
7. **Arnoldussen IAC**, Leijssen EM, Gustafson DR, de Leeuw FE, Kiliaan AJ.
Body mass index is inversely associated with gray matter volumes. The RUN DMC study
Submitted to *Neurology*. 2018. (5-year impact factor: 8.3).
8. Pelgrim CE, Franx BAA, Snabel J, Kleemann R, **Arnoldussen IAC**, Kiliaan AJ.
Butyrate Reduces HFD-Induced Adipocyte Hypertrophy and Metabolic Risk Factors in Obese LDLr^{-/-}.Leiden Mice.
Nutrients. 2017. (5-year impact factor: 4.6).
9. Franx BA, **Arnoldussen IAC**, Kiliaan AJ, Gustafson DR.
Weight Loss in Patients with Dementia: Considering the Potential Impact of Pharmacotherapy.
Drugs Aging. 2017. (5-year impact factor: 2.4).

10. van der Doelen RHA, Robbroch B, **Arnoldussen IAC**, Schulpen M, Homb, Kozicz T.
Serotonin and urocortin 1 in the dorsal raphe and Edinger-Westphal nuclei after early life stress in serotonin transporter knockout rats.
Neuroscience. 2017. (5-year impact factor: 3.4).
11. van der Doelen RH, **Arnoldussen IAC**, Ghareh H, van Och L, Homberg JR, Kozicz T.
Early life adversity and serotonin transporter gene variation interact to affect DNA methylation of the corticotropin-releasing factor gene promoter region in the adult rat brain.
Development and Psychopathology. 2015. (5-year impact factor: 4.3).
12. Jansen D, Zerbi V, **Arnoldussen IAC**, Wiesmann M, Rijpma A, Fang XT, Dederen PJ, Mutsaers MP, Broersen LM, Lütjohann D, Miller M, Joosten LA, Heerschap A, Kiliaan AJ.
Effects of specific multi-nutrient enriched diets on cerebral metabolism, cognition and neuropathology in A β PPswe-PS1dE9 mice.
PLoS One. 2013. (5-year impact factor: 3.5).
13. Janssen CI, Zerbi V, Mutsaers MP, de Jong BS, Wiesmann M, **Arnoldussen IAC**, Geenen B, Heerschap A, Muskiet FA, Jouni ZE, van Tol EA, Gross G, Homberg JR, Berg BM, Kiliaan AJ.
Impact of dietary n-3 polyunsaturated fatty acids on cognition, motor skills and hippocampal neurogenesis in developing C57BL/6J mice.
The Journal of Nutritional Biochemistry. 2015. (5-year impact factor: 4.7).
14. Rijpma A, Jansen D, **Arnoldussen IAC**, Fang XT, Wiesmann M, Mutsaers MP, Dederen PJ, Janssen CI, Kiliaan AJ.
Sex Differences in Presynaptic Density and Neurogenesis in Middle-Aged ApoE4 and ApoE Knockout Mice.
Journal of Neurodegenerative Diseases. 2013. (5-year impact factor: 0.6).
15. Vreeken D, Wiesmann M, Deden LN, **Arnoldussen IAC**, Aarts E, Kessels, RPC, Kleemann R, Hazebroek EJ, Aarts EO, Kiliaan AJ.
The BARICO study: A longitudinal, prospective observational study to evaluate the effects of weight loss after bariatric surgery on brain function and structure. Study rationale and protocol.
Submitted to BMJ Open. 2018. (5-year impact factor: 2.3).
16. Jacobs SA, Wekking L, Vreeken D, Franx BA, Wiesmann M, Verweij V, Geenen B, Dederen PJ, van Tol EA, Gross G, Morrison M, Kleemann R, **Arnoldussen IAC**, Kiliaan AJ.
Sex specific alterations in brain structure, function and metabolism in juvenile HFD-induced obese LDLr-/- Leiden mice.
In preparation. 2018.

Curriculum Vitae

Ilse Arnoldussen werd geboren op 9 november 1988 te Gennep. Ze groeide op in Ottersum en behaalde haar VWO-diploma aan het Elzendaal college te Boxmeer in 2007. In datzelfde jaar begon zij aan de studie Biologie aan de Radboud Universiteit te Nijmegen. Vanwege haar grote interesse in het ontstaan van neurologische aandoeningen koos zij na het behalen van haar bachelordiploma voor een masteropleiding medische biologie met als focus neurowetenschappen.

Tijdens haar masteropleiding liep Ilse twee stages. Tijdens haar eerste masterstage onderzocht zij de rol van een multi-nutrienten dieet op het brein in muismodellen voor de ziekte van Alzheimer onder begeleiding van dr. Diane Jansen, dr. Valerio Zerbi en dr. Amanda J. Kiliaan. In haar tweede stage onderzocht Ilse de invloeden van stress in het vroege leven in ratten, waarbij zij vooral breinweefsel onderzocht op eiwit- en mRNA niveau. In deze tweede stage werd zij begeleid door dr. Rick van der Doelen en dr. Tamás Kozicz. Beide stages vergrootte haar interesse in neurowetenschappen en voeding, daarnaast haar voorliefde om *in vivo* experimenten op te volgen tot in groot detail in *ex vivo* hersenweefsel.

In oktober 2012 begon Ilse haar promotieonderzoek op de afdeling Anatomie van het Radboudumc te Nijmegen onder begeleiding van copromotoren: dr. Amanda J. Kiliaan en dr. Maximilian Wiesmann, en promotoren: prof. dr. Deborah R. Gustafson en dr. Otto Boerman. Tijdens haar promotie beschreven in dit proefschrift voerde zij een aantal studies uit in muismodellen in samenwerking met Mead Johnson (RB) en TNO Leiden. In de tweede fase van haar promotieonderzoek reisde zij naar New York city om onder begeleiding van haar promotor, Professor Gustafson zich verder te verdiepen in epidemiologisch onderzoek. Tijdens haar promotietraject begeleidde zij een tiental studenten gedurende hun stages. Tevens presenteerde zij haar bevindingen op verschillende nationale en internationale bijeenkomsten in de vorm van posters en mondelinge presentaties. Heden is Ilse werkzaam als Post-doc op het EFRO BriteN project aan de afdeling Anatomie, Radboudumc te Nijmegen.

Radboud Alzheimer Centrum Series

1. Van Horssen, J. (2005). *Heparan sulfate proteoglycans and vascular pathology in Alzheimer's disease*
2. Wilhelmus, M.M.M. (2006). *Small heat shock proteins and apolipoprotein E in Alzheimer's disease*
3. Zuidema, S.U. (2008). *Neuropsychiatric symptoms in Dutch nursing home patients with dementia*
4. Graff, M.J.L. (2008). *Effectiveness and efficiency of community occupational therapy for older people with dementia and the caregivers*
5. Claassen, J.A.H.R. (2008). *Cerebral hemodynamics in aging: the interplay between blood pressure, cerebral perfusion, and dementia*
6. Dado- Van Beek, H.E.A. (2010). *The regulation of cerebral perfusion in patients with Alzheimer's disease*
7. De Jong, D. (2010). *Anti-inflammatory therapy and cerebrospinal fluid diagnosis in Alzheimer's disease*
8. Persoon, J.W.B. (2010). *Development and validation of the Nurses' Observation Scale for Cognitive Abilities – NOSCA*
9. Meulenbroek, O.V. (2010). *Neural correlates of episodic memory in healthy aging and Alzheimer's disease*
10. Timmer, N.M. (2011). *The interaction of heparan sulfate proteoglycans with the amyloid b protein*
11. Schölzel-Dorenbos, C.J.M. (2011). *Quality of life in dementia: From concept to practice*
12. Bruinsma, I.B. (2011). *Amyloidogenic proteins in Alzheimer's disease and Parkinson's disease: interaction with chaperones and inflammation*
13. Perry, M. (2011). *Development and evaluation of a Dementia Training Programme for Primary care*
14. Derksen, E.W.C. (2011). *Diagnostic disclosure: a basic intervention in dementia care*
15. Wetzels, R.B. (2011). *Neuropsychiatric symptoms in institutionalized residents with dementia: Course and interplay with cognition, quality of life and psychotropic drug use.*
16. Voight-Radloff, S. (2012). *Cross-national transfer of community occupational therapy in dementia.*

17. Spies, P.E. (2012). *The reflection of Alzheimer disease in CSF*.
18. Joosten-Weyn Banning, E.W.A. (2012). *Learning to live with Mild Cognitive Impairment: development and evaluation of a psychological intervention for patients with Mild Cognitive Impairment and their significant others*.
19. Vasse, E. (2012). *A stepwise process for developing and implementing quality indicators to improve psychosocial dementia care in European countries*.
20. Slats, D. (2012). *CSF biomarkers of Alzheimer's disease; serial sampling analysis and the study of circadian rhythmicity*.
21. Leontjevas, R. (2012). *Act in case of Depression! Validation and effectiveness of a multidisciplinary depression care program in nursing homes*.
22. Bakker, C. (2013). *Young onset dementia: care needs & service provision*.
23. Meeuwssen, E.J. (2013). *Towards efficient dementia care : a comparison of memory clinics and general practitioners*.
24. Spijker, G.J.A.A (2013). *Systematic care for caregivers of people with dementia in community mental health services*.
25. Janssen, D. (2013). *The role of nutrition in Alzheimer's disease : a study in transgenic mouse models for Alzheimer's disease and vascular disorders*.
26. Zerbi, V. (2013). *Impact of nutrition on brain structure and function : a magnetic resonance imaging approach in Alzheimer mouse models*.
27. Herbert, M. (2014). *Facing uncertain diagnosis: the use of CSF biomarkers for the differential diagnosis of neurodegenerative diseases*.
28. Ven, G. van de (2014). *Effectiveness and costs of Dementia Care Mapping intervention in Dutch nursing homes*
29. Döpp, C.M.E. (2015). *Making the jump-The translation of research evidence into clinical occupational therapy practice*.
30. Elsen, G. van den (2016). *Tetrahydrocannabinol in the treatment of neuropsychiatric symptoms in dementia*.
31. Vermeij, A. (2016). *Cognitive plasticity in normal aging and mild cognitive impairment: Shedding light on prefrontal activation*.
32. Müller, M. (2016). *Footprints of Alzheimer's disease. Exploring proteins and microRNAs as biomarkers for differential diagnosis*.
33. Bruggink, K.A. (2016). *Amyloid- β and amyloid associated proteins in the pathology and diagnosis of Alzheimer's disease*.

34. Aalbers, T. (2016). *eHealth in the primary prevention of cognitive decline; The Brain Aging Monitor study.*
35. Van der Maaden, T. (2016). *Improving discomfort in nursing home residents with dementia and pneumonia. Development, implementation and evaluation of a practice guideline for optimal symptom relief.*
36. Millenaar, J. (2016). *Young onset dementia, towards a better understanding of care needs and experiences.*
37. Rijpma, A. (2017). *Multi-nutrient interventions and brain metabolism in Alzheimer's disease: a spectrum of effects.*
38. Wiesmann, M. (2017). *Vascular risk factors and Alzheimer's disease. Therapeutic approaches in mouse models.*
39. Jong, D.L.K. de (2018). *Regulation of cerebral perfusion in Alzheimer's disease: from seconds to months.*
40. Donkers, H.W. (2018). *Social Participation Dilemma's in dementia.*
41. Richters, A. (2018). *Network-based care for people with dementia: a complex transition.*
42. Spek, K. van der (2018). *Appropriate psychotropic drug use in institutionalized people with dementia. The PROPER-study.*
43. Arnoldussen, I.A.C. (2018). *Adiposity and the Brain.*

Donders Graduate School for Cognitive Neuroscience

For a successful research Institute, it is vital to train the next generation of young scientists. To achieve this goal, the Donders Institute for Brain, Cognition and Behaviour established the Donders Graduate School for Cognitive Neuroscience (DGCN), which was officially recognised as a national graduate school in 2009. The Graduate School covers training at both Master's and PhD level and provides an excellent educational context fully aligned with the research programme of the Donders Institute.

The school successfully attracts highly talented national and international students in biology, physics, psycholinguistics, psychology, behavioral science, medicine and related disciplines. Selective admission and assessment centers guarantee the enrolment of the best and most motivated students.

The DGCN tracks the career of PhD graduates carefully. More than 50% of PhD alumni show a continuation in academia with postdoc positions at top institutes worldwide, e.g. Stanford University, University of Oxford, University of Cambridge, UCL London, MPI Leipzig, Hanyang University in South Korea, NTNU Norway, University of Illinois, North Western University, Northeastern University in Boston, ETH Zürich, University of Vienna etc.. Positions outside academia spread among the following sectors: specialists in a medical environment, mainly in genetics, geriatrics, psychiatry and neurology. Specialists in a psychological environment, e.g. as specialist in neuropsychology, psychological diagnostics or therapy. Positions in higher education as coordinators or lecturers. A smaller percentage enters business as research consultants, analysts or head of research and development. Fewer graduates stay in a research environment as lab coordinators, technical support or policy advisors. Upcoming possibilities are positions in the IT sector and management position in pharmaceutical industry. In general, the PhDs graduates almost invariably continue with high-quality positions that play an important role in our knowledge economy.

For more information on the DGCN as well as past and upcoming defenses please visit:

<http://www.ru.nl/donders/graduate-school/phd/>